



FOOD STANDARDS
Australia New Zealand
Te Mana Kounga Kai – Ahitereiria me Aotearoa

09/03
21 May 2003

DRAFT ASSESSMENT REPORT

APPLICATION A472

D-TAGATOSE AS A NOVEL FOOD

DEADLINE FOR PUBLIC SUBMISSIONS to the Authority in relation to this matter:

2 July 2003

(See “Invitation for Public Submissions” for details)


FOOD STANDARDS AUSTRALIA NEW ZEALAND

FSANZ's role is to protect the health and safety of people in Australia and New Zealand through the maintenance of a safe food supply. FSANZ is a partnership between ten Governments: the Commonwealth; Australian States and Territories; and New Zealand. It is a statutory authority under Commonwealth law and is an independent, expert body.

FSANZ is responsible for developing, varying and reviewing standards and for developing codes of conduct with industry for food available in Australia and New Zealand covering labelling, composition and contaminants. In Australia, FSANZ also develops food standards for food safety, maximum residue limits, primary production and processing and a range of other functions including the coordination of national food surveillance and recall systems, conducting research and assessing policies about imported food.

The FSANZ Board approves new standards or variations to food standards in accordance with policy guidelines set by the Australia and New Zealand Food Regulation Ministerial Council (Ministerial Council) made up of Commonwealth, State and Territory and New Zealand Health Ministers as lead Ministers, with representation from other portfolios. Approved standards are then notified to the Ministerial Council. The Ministerial Council may then request that FSANZ review a proposed or existing standard. If the Ministerial Council does not request that FSANZ review the draft standard, or amends a draft standard, the standard is adopted by reference under the food laws of the Commonwealth, States, Territories and New Zealand. The Ministerial Council can, independently of a notification from FSANZ, request that FSANZ review a standard.

The process for amending the *Australia New Zealand Food Standards Code* is prescribed in the *Food Standards Australia New Zealand Act 1991* (FSANZ Act). The diagram below represents the different stages in the process including when periods of public consultation occur. This process varies for matters that are urgent or minor in significance or complexity.



INVITATION FOR PUBLIC SUBMISSIONS

The Authority has prepared a Draft Assessment Report of Application A472 – D-tagatose as a novel food; and prepared draft variations to the *Australia New Zealand Food Standards Code*.

The Authority invites public comment on this Draft Assessment Report based on regulation impact principles and the draft variations to the *Australia New Zealand Food Standards Code* for the purpose of preparing an amendment to the *Australia New Zealand Food Standards Code* for approval by the FSANZ Board.

Written submissions are invited from interested individuals and organisations to assist the Authority in preparing the Draft Assessment/Final Assessment for this application/proposal. Submissions should, where possible, address the objectives of the Authority as set out in section 10 of the *Food Standards Australia New Zealand Act 1991* (FSANZ Act). Information providing details of potential costs and benefits of the proposed change to the *Australia New Zealand Food Standards Code* from stakeholders is highly desirable. Claims made in submissions should be supported wherever possible by referencing or including relevant studies, research findings, trials, surveys etc. Technical information should be in sufficient detail to allow independent scientific assessment.

The processes of the Authority are open to public scrutiny, and any submissions received will ordinarily be placed on the public register of the Authority and made available for inspection. If you wish any information contained in a submission to remain confidential to the Authority, you should clearly identify the sensitive information and provide justification for treating it as commercial-in-confidence. Section 39 of the FSANZ Act requires the Authority to treat in-confidence, trade secrets relating to food and any other information relating to food, the commercial value of which would be, or could reasonably be expected to be, destroyed or diminished by disclosure.

Submissions must be made in writing and should clearly be marked with the word “Submission” and quote the correct project number and name. Submissions may be sent to one of the following addresses:

Food Standards Australia New Zealand
PO Box 7186
Canberra BC ACT 2610
AUSTRALIA
Tel (02) 6271 2222 Fax (02) 6271 2278
www.foodstandards.gov.au

Food Standards Australia New Zealand
PO Box 10559
The Terrace WELLINGTON 6036
NEW ZEALAND
Tel (04) 473 9942 Fax (04) 473 9855
www.foodstandards.govt.nz

Submissions should be received by the Authority by: **2 July 2003**. Submissions received after this date may not be considered unless the Project Manager has given prior agreement for an extension. Submissions may also be sent electronically through the FSANZ website using the [Standards Development](#) tab and then through [Documents for Public Comment](#). Questions relating to making submissions or the application process can be directed to the Standards Liaison Officer at the above address or by emailing slo@foodstandards.gov.au.

Further Information

Further information on this Proposal and the assessment process should be addressed to the FSANZ Standards Liaison Officer at one of the following addresses:

Food Standards Australia New Zealand
PO Box 7186
Canberra BC ACT 2610
AUSTRALIA
Tel (02) 6271 2222
www.foodstandards.gov.au

Food Standards Australia New Zealand
PO Box 10559
The Terrace WELLINGTON 6036
NEW ZEALAND
Tel (04) 473 9942
www.foodstandards.govt.nz

Assessment reports are available for viewing and downloading from the FSANZ website www.foodstandards.gov.au or alternatively paper copies of reports can be requested from the Authority's Information Officer at info@foodstandards.gov.au including other general enquiries and requests for information.

CONTENTS

EXECUTIVE SUMMARY AND STATEMENT OF REASONS	7
1. INTRODUCTION.....	9
2. REGULATORY PROBLEM.....	9
3. OBJECTIVE	9
4. BACKGROUND	10
4.1 PROPERTIES OF D-TAGATOSE.....	10
4.2 PROPOSED USES	10
4.3 APPROVAL IN OTHER COUNTRIES	11
5. ISSUES ADDRESSED DURING ASSESSMENT	11
5.1 SAFETY ISSUES.....	11
5.2 TECHNOLOGICAL FUNCTION	11
5.3 DIETARY CONSIDERATIONS.....	11
5.4 RISK ASSESSMENT OF D-TAGATOSE	12
5.5 NUTRITIONAL CONSIDERATIONS.....	13
5.6 ENERGY VALUE FOR D-TAGATOSE	13
5.7 ISSUES RAISED IN PUBLIC SUBMISSIONS.....	14
6. REGULATORY OPTIONS.....	14
7. IMPACT ANALYSIS	14
7.1 OPTION 1. NOT PERMIT THE USE OF D-TAGATOSE	15
7.2 OPTION 2. PERMIT THE USE OF D-TAGATOSE	15
8. CONSULTATION	15
8.1 PUBLIC CONSULTATION	15
8.2 WORLD TRADE ORGANIZATION (WTO) NOTIFICATION.....	15
9. CONCLUSIONS AND RECOMMENDATION.....	16
ATTACHMENTS	16
ATTACHMENT 1	17
DRAFT VARIATIONS TO THE AUSTRALIA NEW ZEALAND FOOD STANDARDS CODE	17
ATTACHMENT 2	18
DRAFT SAFETY ASSESSMENT REPORT	18
ATTACHMENT 3	30
FOOD TECHNOLOGY REPORT	30
ATTACHMENT 4	33

DIETARY EXPOSURE ASSESSMENT REPORT	33
ATTACHMENT 5	44
ENERGY FACTOR FOR D-TAGATOSE.....	44
ATTACHMENT 6	58
SUMMARY OF FIRST ROUND PUBLIC SUBMISSIONS.....	58

Executive Summary and Statement of Reasons

Food Standards Australia New Zealand (FSANZ) received an Application from Aria Food Ingredients a/s (Denmark) on 17 July 2002 seeking to amend Standard 1.5.1 of the Food Standards Code to permit the use of D-tagatose as a novel food ingredient. This application is at the draft assessment stage under section 15 of the *Food Standards Australia New Zealand Act 1991* (FSANZ Act). The work on this application was commenced on 08 August 2002.

Under the current food standards, novel foods are required to undergo a pre-market safety assessment, as per Standard 1.5.1 - Novel Foods. D-Tagatose can be considered a 'sugar' as defined in the Standard 2.8.1 in the *Australia New Zealand Food Standard Code* (the Code) and hence a food. It is also considered a non-traditional food because it has no history of significant human consumption in Australia or New Zealand. Its safety in the context in which it is presented has not yet been determined within the context of the Australian and New Zealand diet. For these reasons, D-tagatose is considered to be a novel food and is accordingly considered under Standard 1.5.1.

The objective of this assessment is to determine whether it is appropriate to amend the Code to permit the use of D-tagatose as a novel food. Such an amendment would need to be consistent with the section 10 objectives of the FSANZ Act.

A range of issues was considered during the assessment of the application. An analysis of the safety and nutritional impact of the use of D-tagatose indicate that there are no public health and safety concerns at the anticipated levels of dietary exposure. D-tagatose has technological properties similar to traditional sugars such as glucose and fructose and can be used as a reducing sugar as it caramelises at elevated temperatures. However, in contrast to traditional sugars, it is only partially absorbed by the body resulting in reduced energy value.

The only regulatory options identified were to approve or not approve the use of D-tagatose as a novel food. The regulatory impact analysis shows that benefits accrue to industry and Governments, in terms of enhanced market opportunities and trade (under Australia and New Zealand's requirements under the World Trade Organization), respectively, and to consumers through potentially a greater choice of foods.

Three submissions were received during the consultation period. One submitter supported the proposal to amend the *Food Standards Code* to permit D-tagatose. One argued that D-tagatose is a traditional food and therefore should not be regulated, while another was undecided on whether permission should be given or not because of insufficient information in the Initial Assessment Report.

Draft Statement of Reasons

FSANZ recommends the approval of the use of D-tagatose as a novel food for the following reasons:

- there are no public health and safety concern associated with the use of D-tagatose as proposed;
- D-tagatose has technological properties similar to traditional sugars but its energy value is lower;

- the proposed changes to the Code are consistent with the section 10 objectives of the FSANZ Act; and
- the Regulatory Impact Statement indicates that, for the preferred option, namely, to approve the use of D-tagatose as a novel food, the benefits of the proposed amendment outweigh the costs.

The proposed drafting for amendment to Standard 1.5.1 is at **Attachment 1** of the Draft Assessment Report.

1. INTRODUCTION

An application was received from Aria Food Ingredients a/s (Denmark) on 17 July 2002 seeking to amend Standard 1.5.1 of the Code to permit the use of D-tagatose as a novel food ingredient. After consultations with FSANZ, the applicant has now withdrawn a written request for considering the Net Metabolisable Energy (NME) value for their product in Standard 1.2.8 in the assessment of the application for approval of D-tagatose. This application is at the draft assessment stage under section 15 of the FSANZ Act.

2. REGULATORY PROBLEM

Under the current food standards, novel foods are required to undergo a pre-market safety assessment, as per Standard 1.5.1 – Novel Foods. The purpose of Standard 1.5.1 is to ensure that non-traditional foods that have features or characteristics that may raise safety concerns will undergo a risk-based safety assessment before they are offered for retail sale in Australia or New Zealand.

Novel Food is defined in the Standard as:

A non-traditional food or food ingredient for which there is insufficient knowledge in the broad community to enable safe use in the form or context in which it is presented, taking into account;

- (a) the composition or structure of the product;
- (b) levels of undesirable substances in the product;
- (c) the potential for adverse effects in humans;
- (d) traditional preparation and cooking methods; or
- (d) patterns and levels of consumption of the product.

Non-traditional food means a food which does not have a history of significant human consumption by the broad community in Australia or New Zealand.

D-Tagatose is considered a ‘sugar’ as defined in the Standard 2.8.1 in the Code and hence a food ingredient. It is also considered a non-traditional food ingredient because it has no history of significant human consumption in Australia or New Zealand. D-Tagatose is found in sterilised milk and milk powder as well as in other dairy products (between 2 to 800 ppm), but its safety in larger quantities (> 1%, 10000 ppm) and the context in which it is presented have not yet been determined within the context of the Australian and New Zealand diet. In these circumstances, D-Tagatose is considered to be a novel food ingredient and should be considered under Standard 1.5.1.

3. OBJECTIVE

The objective of this assessment is to determine whether it is appropriate to amend the Code and permit the use of D-tagatose. Such an amendment to the Code will need to be consistent with the section 10 objectives of FSANZ Act.

In developing or varying a food standard, FSANZ is required by its legislation to meet three primary objectives which are set out in section 10 of the FSANZ Act. These are:

- the protection of public health and safety;
- the provision of adequate information relating to food to enable consumers to make informed choices; and
- the prevention of misleading or deceptive conduct.

In developing and varying standards, FSANZ must also have regard to:

- the need for standards to be based on risk analysis using the best available scientific evidence;
- the promotion of consistency between domestic and international food standards;
- the desirability of an efficient and internationally competitive food industry;
- the promotion of fair trading in food; and
- any written policy guidelines formulated by the Ministerial Council.

4. BACKGROUND

4.1 Properties of D-tagatose

D-Tagatose is a naturally occurring monosaccharide and an enantiomer of D-fructose (inversion at C-4). It is an odourless white crystalline solid, with a sweet taste. It is produced from lactose in a two-step process involving enzymatic hydrolysis of lactose to galactose, followed by isomerisation under basic conditions.

D-Tagatose has technological properties similar to traditional sugars such as glucose and fructose and can be used as a reducing sugar as it caramelises at elevated temperatures. However, it is different from traditional sugars in that it is only partially absorbed by the body resulting in reduced energy value. The major fraction of D-tagatose reaches the large intestine unabsorbed (where it undergoes fermentation). Thus it offers health advantages as a low energy sugar with low glycaemic index.

4.2 Proposed uses

The substance is proposed to be used in the following foods:

- breakfast cereals (ready to eat);
- carbonated diet soft drinks;
- non-carbonated diet soft drinks;
- low fat/fat free ice cream;
- low fat frozen dairy desserts;
- diet/health bars;
- diet soft confectionery;
- hard confectionary;
- icings/frostings; and
- special purpose foods/meal replacements.

4.3 Approval in other countries

D-Tagatose is available in the USA as a Generally Recognised As Safe (GRAS) dietary ingredient (GRAS Notice No. GRN 000078).

There are no Codex standards in relation to D-tagatose.

5. ISSUES ADDRESSED DURING ASSESSMENT

5.1 Safety issues

D-Tagatose is found naturally in dairy products (between 2 to 800 ppm), but its safety in larger quantities (> 1%, 10000 ppm) and the context in which it is presented needed to be determined within the context of the Australian and New Zealand diet prior to approval of its use as a food ingredient in Australia and New Zealand.

In its safety assessment FSANZ considered the recent evaluation conducted by Joint FAO/WHO Expert Committee on Food Additives (JECFA) as well as reviewing more recent studies. The safety assessment undertaken by JECFA of D-tagatose for certain specified uses concluded that D-tagatose was a substance of low toxicity, which did not represent a hazard to human health. JECFA allocated an Allowable Daily Intake (ADI) of '80 mg/kg bw/day'¹ for D-tagatose based on the concern about its potential to induce glycogen deposition and hypertrophy in the liver and to increase the concentrations of uric acid in serum.

However new data available to FSANZ indicates that there is no evidence of any public health and safety concern associated with consumption of D-tagatose up to 15 g/d (0.25 g/kg bw/d for a 60 kg adult) even in susceptible groups, hyperuricemic and gouty subjects. Mild gastrointestinal effects such as nausea, diarrhoea and flatulence occurred in some individuals at bolus doses of 30g/d and above. FSANZ's toxicology report (**Attachment 2**) contains a summary of JECFA's evaluation as well as an assessment of the new additional data from carcinogenicity study in rats and the human toleration studies. Based on these toxicological data, FSANZ concluded that there are no public health and safety concerns associated with the use of D-tagatose in food up to 15g/day (0.25g/kg bw/d) although mild GI effects may be observed at higher dose levels.

5.2 Technological function

The Food Technology Report (**Attachment 3**) concludes that D-tagatose is a naturally occurring sweet-tasting monosaccharide that may be used in a wide range of foods. It could be used in any application permitted by regulation.

5.3 Dietary considerations

Based on the US food consumption data (1994-1996, 1998) JECFA has estimated that the mean estimated intake for D-tagatose from all its proposed uses excluding chewing gums and formula diets was approximately 5 grams per day and for the 90th percentile group was about 11 grams per day. These estimates are based on the use of D-tagatose in all proposed uses at the highest levels. The realistic dietary intake would be expected to be much lower.

¹ FAO/WHO (2002) Report of fifty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives, Evaluation of Certain Food Additives and Contaminants, D-Tagatose, p29.

During the Draft Assessment dietary modelling was conducted by FSANZ to estimate the potential dietary intake of D-tagatose in Australia and New Zealand that may result from permitting its use in the foods specified in the application.

Estimated chronic dietary exposures to D-tagatose were all below 15g/person/day for all population groups. The estimated mean exposure for consumers of D-tagatose for the Australian population aged 2 years and above is 2.7g/day. The estimated mean exposure for consumers of D-tagatose for Australian toddlers aged 2 – 4 years is 1.9g/day and for Australian children aged 5 – 12 years it is 3.0g/day. The estimated mean exposure for consumers of D-tagatose for the New Zealand population aged 15 years and above is 1.9g/day. The estimated 95th percentile exposures for consumers of D-tagatose for the total Australian and New Zealand populations are 9g/day and 7.4g/day respectively. The 95th percentile exposures in children 2-4 years and 5-12 years were 6.9 and 10.5g/day respectively. Estimates of dietary exposures in this case have not been expressed as a per bodyweight figure as it is the bolus dose rather than the per bodyweight dose which is relevant when considering reversible and minor discomforts such as laxation effects.

Estimated short-term exposures were calculated for individual food groups to assess whether gastro-intestinal effects were likely. All estimated short-term exposures were below the dose level when mild laxation effects are observed in some individuals. One exception is individuals consuming large amounts of formulated dietary foods or formulated supplementary sports foods where conservative dietary modelling indicated potential for higher levels of exposure. The full dietary modelling report is attached (**Attachment 4**).

5.4 Risk assessment of D-tagatose

The data support the safety of D-tagatose at the level of exposure that would be achieved by addition of D-tagatose to a range of foods at the maximum levels provided by the Applicant. Exposure for all ages for Australian and New Zealand populations is below the level at which mild gastro-intestinal effects were observed, even at the 95th percentile (high consumer) exposure level. The highest potential exposure by age group was Australian children 5-12 years at the 95th percentile exposure level (10.5g/day), which is still within the level at which mild, gastro-intestinal effects would be observed (**Attachment 4**).

The potential to exceed the threshold dose for mild gastrointestinal effects is considered low if D-tagatose is consumed over a 24-hour period. This is partly due to the conservative nature of the assumptions used in the modelling; namely:

- D-tagatose would have to be added to all the foods proposed which is extremely unlikely;
- all the foods would have to contain D-tagatose at the maximum level proposed;
- the data used for modelling is a 24-hour food recall which overestimates food consumption for high consumers (the use of multiple day records tends to significantly reduce predicted high consumer exposure); and, that
- the modelling for short term exposure assumes that the individual foods eaten over 24 hours were in fact eaten at one time not over a period of two or more occasions during the day, which is more likely (e.g. confectionery).

The data reflecting a single eating occasion exposure, suggested that Australian adults consuming large amounts of formulated dietary foods, or formulated supplementary sports foods (95th percentile consumption level) could potentially have a single consumption of 59g D-tagatose/day on one eating occasion, however, these products are likely to be infrequently consumed. The estimates are a worst-case scenario based on conservative assumptions.

In conclusion there are no public health and safety concerns associated with the use of D-tagatose in foods as proposed. As better information become available on the actual use levels of D-tagatose in various foods, more accurate estimates of dietary exposures will be possible and the public health and safety concerns reassessed, for high intake consumers.

5.5 Nutritional considerations

About 20-25% of ingested D-tagatose is absorbed from the small intestine. The absorbed fraction is metabolised via the same pathway as fructose, with the remaining unabsorbed fraction being available for fermentation in the large bowel.

No scientific evidence has been identified linking D-tagatose consumption to a potential decrease in the bioavailability or intake of essential micronutrients. There has been some suggestion though, that D-tagatose consumption could influence the absorption of fructose due to similarities in their chemical structures. However, scientific evidence reviewed at the 55th meeting of JECFA² indicated that a high intake of D-tagatose does not significantly affect the intestinal absorption of fructose.

On the basis of this limited data, it is concluded that the main nutritional effect of D-tagatose consumption will be a reduction in dietary energy intakes, the intended purpose for using a reduced energy sugar substitute. Therefore, the use of D-tagatose as a novel food ingredient is unlikely to have a significant impact on any other nutritional aspects when considered in the context of the overall diet.

5.6 Energy value for D-tagatose

Because of its poor absorption, D-tagatose potentially has lower energy value, which may make it a useful food ingredient in low-energy food products. The Standard 1.2.8- Nutritional Information Requirements lists energy values of approved food ingredients in Table 1 to subclause 2(2). Currently it has no provision for using Net Metabolisable Energy (NME) value in this Table. After consultations with FSANZ, the Applicant has now withdrawn his initial written request for considering the inclusion of NME value in Table 1 to subclause 2(2) of the Standard 1.2.8 - Nutritional Information Requirements.

The supporting experimental evidence for a lower energy value of D-tagatose was considered at Draft Assessment (**Attachment 5**). The metabolisable energy factor for D-tagatose is best reflected by a value of 11 kJ/g. It is therefore recommended that this energy factor be inserted into Table 2 of subclause 2(2) of Standard 1.2.8.

² FAO/WHO (2001) Report of fifty-fifth report of the Joint FAO/WHO Expert Committee on Food Additives, Evaluation of Certain Food Additives and Contaminants, D-Tagatose, p13.

5.7 Issues Raised In Public Submissions

Comment: *Australian Food and Grocery Council* argues that FSANZ needs to determine before accepting the application as a novel food, that it is a non-traditional food and that there is not sufficient knowledge in the broad community to enable its safe use; further, if FSANZ finds that it is safe for human consumption, then it cannot be listed as a novel food.

Response: The object of the Novel Food Standard is to assess the safety of non-traditional food for which there is ‘there is insufficient knowledge to enable safe use’ in the broader community. Prior to the application, D-tagatose at the proposed levels of use had not undergone a safety assessment in the context of the Australian and New Zealand diets. There was therefore insufficient knowledge in the broad community to ensure safe use in the form in which it is presented. The safety of D-tagatose, in the context of Australia and New Zealand, has been assessed by FSANZ as a consequence of D-tagatose being classified as a novel food. The safety assessment showed subsequently that D-tagatose is safe for human consumption, but without the due process required for novel food assessment, it would not have been possible for FSANZ to determine its safety.

Comment: *Department of Health, WA*, is ‘disappointed that FSANZ had taken significant direction from the USFDA assessment and approval of D-tagatose, and that there was no apparent effort to establish the relevance to the Australian setting and population’.

Response: The Initial Assessment Report merely states the issues and current regulatory status of the product around the world relating to the application. A complete assessment of the product is carried out by FSANZ only after gathering and assessing comments from all sources through the public submission process. The Draft Assessment Report describes toxicological and other evaluations carried out FSANZ staff.

6. REGULATORY OPTIONS

FSANZ is required to consider the impact of various regulatory (and non-regulatory) options on all sectors of the community, which includes consumers, the food industry and governments in both Australia and New Zealand. The benefits and costs associated with the proposed amendment to the Code will be analysed in a Regulatory Impact Assessment.

The regulatory options currently under consideration are:

Option 1. Not permit the use of D-tagatose.

Option 2. Permit the use of D-tagatose.

7. IMPACT ANALYSIS

Parties possibly affected by the options outlined include:

1. Food industry wishing to promote food products with D-tagatose.
2. Consumers who may benefit from the use of D-tagatose -containing products.
3. Government agencies enforcing the food regulations.

The draft regulatory options are as follows:

7.1 Option 1. Not permit the use of D-tagatose

On the basis of this Draft Assessment, there are no perceived benefits to government, consumers or industry by maintaining the *status quo* and not giving specific permission in the Code for the use of this ingredient.

On the basis of this Draft Assessment, there is no perceived cost for the government, however lack of approval in Australia or New Zealand may be construed as a non-tariff barrier to trade unless it is based on public health and safety considerations. Industry may also suffer from the non-availability of this ingredient.

Parties potentially disadvantaged by not permitting this substance, are the manufacturers of D-tagatose and producers who wish to use it in the manufacture of their final food products and consumers who may benefit from its health effects.

7.2 Option 2. Permit the use of D-tagatose

On the basis of this Draft Assessment, industry and consumers would benefit from this option. This option would result in no cost to government, industry or consumers, if its safety can be ensured.

Approval of D-tagatose would promote international trade in food products.

8. CONSULTATION

8.1 Public Consultation

Three submissions were received during the consultation period. One submitter supported the proposal to amend the *Food Standards Code* to permit D-tagatose. One argued that D-tagatose is a traditional food and therefore should not be regulated, while another was undecided because of insufficient information in the Initial Assessment Report. The names of submitters and issues raised in their submissions are provided in **Attachment 6**.

FSANZ is now seeking public comment in order to assist in assessing this application at Final Assessment. Comments that would be useful could cover:

- scientific aspects of this application;
- other issues including labelling of foods and food products containing D-tagatose.

8.2 World Trade Organization (WTO) Notification

Australia and New Zealand are members of the WTO and are bound as parties to WTO agreements. In Australia, an agreement developed by the Council of Australian Governments (COAG) requires States and Territories to be bound as parties to those WTO agreements to which the Commonwealth is a signatory.

Under the agreement between the Governments of Australia and New Zealand on Uniform Food Standards, FSANZ is required to ensure that food standards are consistent with the obligations of both countries as members of the WTO.

In certain circumstances Australia and New Zealand have an obligation to notify the WTO of changes to food standards to enable other member countries of the WTO to make comment. Notification is required in the case of any new or changed standards that may have a significant trade effect and which depart from the relevant international standard (or where no international standard exists).

Amending the *Food Standards Code* to allow foods to contain D-tagatose will not be notified to the WTO under either the Technical Barrier to Trade (TBT) or Sanitary and Phytosanitary Measure (SPS) agreements as the permission is unlikely to significantly effect trade. The potential uses for D-tagatose as a carrier in food products is limited. Typically only very small amounts <1% are used and the range of potential suitable processed food products is limited. Suppliers of food products are not required to take up permissions granted through amendments to the *Code*.

9. CONCLUSIONS AND RECOMMENDATION

FSANZ recommends the approval of the use of D-tagatose as a novel food for the following reasons:

- there are no public health and safety concern associated with the use of D-tagatose as proposed;
- D-tagatose has technological properties similar to traditional sugars but its energy value is lower;
- the proposed changes to the Code are consistent with the section 10 objectives of the FSANZ Act; and
- the Regulatory Impact Statement indicates that, for the preferred option, namely, to approve the use of D-tagatose as a novel food, the benefits of the proposed amendment outweigh the costs.

ATTACHMENTS

1. Draft variation to the *Australia New Zealand Food Standards Code*
2. Draft Safety Assessment Report
3. Food Technology Report
4. Dietary Modelling Report
5. Nutrition Report
6. Summary of public submissions

ATTACHMENT 1

DRAFT VARIATIONS TO THE *AUSTRALIA NEW ZEALAND FOOD STANDARDS CODE*

To commence: On gazettal

[1] *Standard 1.5.1 of the Australia New Zealand Food Standards Code is varied by inserting in the Table to clause 2 -*

D-Tagatose	May only be added to food according to Standard 1.3.4.
------------	--

[2] *Standard 1.2.8 of the Australia New Zealand Food Standards Code is varied by inserting in the Table 2 to subclause 2(2) -*

D-Tagatose	11
------------	----

DRAFT SAFETY ASSESSMENT REPORT

D-TAGATOSE AS A NOVEL FOOD INGREDIENT

INTRODUCTION

Structure & Properties

D-Tagatose is a stereoisomer (epimer) of D-fructose differing in the spatial configuration of the hydroxyl group at C-4. D-Tagatose is produced from enzymatic hydrolysis of food-grade lactose to D-galactose followed by chemical isomerisation reaction by calcium hydroxide to form D-tagatose, which is then purified by mineralisation, ion exchange chromatography and re-crystallisation.

Natural occurrence

D-Tagatose is found in sterilised milk and milk powder as well as in other dairy products (between 2 to 800 ppm).

Evaluation by JECFA

D-Tagatose was evaluated in 2000 and 2001 by the Joint FAO/WHO Expert Committee on Food Additives (JECFA)^{1,2} which allocated an ADI of 80 mg/kg bw/d based on concerns about its potential to induce glycogen deposition and hypertrophy in the liver and to increase the concentrations of uric acid in serum.

Current evaluation based on JECFA review plus additional data

FSANZ has not reviewed all studies again in detail – instead FSANZ has reviewed key studies used to establish ADI.

Condensed versions of the two reviews by JECFA are given below followed by a detailed toxicological assessment of the new data submitted to FSANZ.

REPORT FROM JECFA 55

D-Tagatose is an epimer Of D-fructose and is produced from D-galactose by isomerisation under alkaline conditions in the presence of calcium. It is used as a sweetener, texturiser, stabilizer, humectant and formulation aid. D-Tagatose has not been evaluated previously by the Committee.

In a study in rats adapted to consumption of D-tagatose, 80-90% of an oral dose was absorbed.

¹ FAO/WHO (2001) Report of fifty-fifth report of the Joint FAO/WHO Expert Committee on Food Additives, Evaluation of Certain Food Additives and Contaminants, D-Tagatose, p13.

² FAO/WHO (2002) Report of fifty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives, Evaluation of Certain Food Additives and Contaminants, D-Tagatose, p29.

The predicted daily intake of D-tagatose was determined on the basis of data on food consumption in the USA and the assumption that all foods in the categories being considered contain the additive at the maximum technological level. For the population of the USA, intake of this sugar from all proposed uses (except chewing-gum, dietary supplements and meal replacements) was predicted to be 9g/day for consumers with mean intakes and 18g/day for those with intakes at the 90th percentile. Intake from chewing gum was predicted to be 4g/day for consumers with mean intakes and 8g/day for those with intakes at the 90th percentile. Similar results were obtained for the predicted intakes of young people aged 3-5 years, 6-12 years and 13-19 years. The estimated intakes of D-tagatose from dietary supplements and meal replacements were 3 g and 5 g per eating occasion, respectively. An analysis based on the same assumptions, combined with available data on food consumption from Australia and the European Union, showed that the predicted intake of D-tagatose would be similar in these regions.

D-Tagatose was tested in Sprague-Dawley rats in a series of short-term toxicity studies. The observed increases in liver weights and liver hypertrophy were found to be due, at least in part, to glycogen accumulation. The hepatic changes were partially reversed after exclusion of D-tagatose from the diet. Recovery from the induced liver hypertrophy took longer than recovery from glycogen accumulation. Data from short-term studies of the mechanism of glycogen accumulation suggest that the hepatic changes are due to physiological changes in Sprague-Dawley rats and that Wistar rats are less sensitive to expression of these effects.

The precise metabolic pathway of D-tagatose that leads to gluconeogenesis has not been established. D-Tagatose is metabolized more slowly than fructose. A similar biochemical effect characterized by glycogen accumulation occurs in patients with hereditary fructose intolerance, and this reaction can increase the rates of purine breakdown and accumulation of uric acid. D-Tagatose is more effective than fructose in increasing the concentration of uric acid in serum.

In two studies of developmental toxicity in Sprague-Dawley rats, minimal effects were observed in dams, including reduced food consumption at doses greater than 12g/kg of body weight per day and initial depression of weight gain, which returned to normal later in the study. A dose-related, statistically significant increase in liver weight was found, but histological examination of the livers revealed no abnormalities. No effects were found in either study on reproductive or developmental parameters.

The results of tests for genotoxicity in vitro and in vivo were consistently negative.

A number of studies of gastrointestinal effects have been conducted in healthy human volunteers and in patients with type 2 diabetes³. Nausea and adverse gastrointestinal effects were reported in healthy adults given D-tagatose at high doses. Studies in which baseline serum concentrations of insulin and glucose were investigated showed no effect following administration of single or multiple doses, but a decreased glycaemic response was observed when D-tagatose was given before a glucose tolerance test.

Elevated serum uric acid concentrations were reported in three out of six studies in which this parameter was measured; in two of these studies, the values exceeded the normal range.

³ Donner TW, Wilber JF, Ostrowski D, *D-Tagatose, a novel hexose: acute effects on carbohydrate tolerance in subjects with and without type 2 diabetes*, **Diabetes Obes Metab** 1999 Sep;1(5):285-91.

In the three studies in which parameters indicative of liver function or hepatic changes were measured, no effects were observed.

On the basis of the available data, the Committee concluded that D-tagatose is not genotoxic, embryotoxic or teratogenic. The Committee noted that the increased liver weights and hepatocellular hypertrophy seen in Sprague-Dawley rats occurred concurrently with increased glycogen deposition; however, the reversal of increased glycogen storage after removal of D-tagatose from the feed occurred more rapidly than regression of the liver hypertrophy.

Although the gastrointestinal symptoms seen in adult humans with the expected daily intake of D-tagatose were minor, the Committee was concerned about the increased serum uric acid concentrations observed in a number of studies in humans following administration of either single or repeated doses of D-tagatose. Similar increases are seen with other sugars, such as fructose, but D-tagatose appears to be a more potent inducer of this effect. The Committee noted that the effect of D-tagatose has not been studied in people prone to high serum uric acid concentrations.

The Committee concluded that an ADI could not be allocated to D-tagatose because of concern about its potential to induce liver glycogen deposition and hypertrophy and to increase serum uric acid concentrations. Two studies in Sprague-Dawley and Wistar rats were submitted that might help to resolve the relevance of the induction of liver glycogen deposition and hypertrophy, but the reports were received in draft form and were not suitable for consideration at the present meeting. Before reviewing the compound again, the Committee would wish to evaluate the final reports of these studies and data to clarify the extent, mechanism and toxicological consequences of the increased serum uric acid concentrations observed in humans exposed to D-tagatose.

A toxicological monograph and new specifications were prepared.

REPORT FROM JECFA57

D-Tagatose is a keto-hexose, an epimer of D-fructose inverted at C-4, with a sweet taste. It is obtained from D-galactose by isomerisation under alkaline conditions in the presence of calcium.

D-Tagatose was evaluated by the Committee at its fifty-fifth meeting (Annex 1, reference 149), when it concluded that the available data indicated that D-tagatose is not genotoxic, embryotoxic or teratogenic. It noted that the increased liver weights and hepatocellular hypertrophy seen in Sprague-Dawley rats occurred concurrently with increased glycogen deposition; however, the reversal of increased glycogen storage after removal of D-tagatose from the feed was more rapid than regression of the liver hypertrophy. Although the gastrointestinal symptoms seen in adult humans with the expected daily intake of D-tagatose were minor, the Committee was concerned about the increased serum uric acid concentrations observed in a number of studies in humans after administration of either single or repeated doses of D-tagatose. Similar increases were observed with other sugars, such as fructose, but D-tagatose appeared to be a more potent inducer of this effect. The Committee also noted that this effect of D-tagatose had not been studied in persons prone to high serum uric acid concentrations. The Committee concluded that an ADI could not be allocated for D-tagatose because of concern about its potential to induce glycogen deposition in the liver and liver hypertrophy and to increase the serum concentration of uric acid.

Two studies of up to 7 days' duration in Wistar and Sprague-Dawley rats given repeated doses of D-tagatose were submitted to the Committee at its fifty-fifth meeting, but the reports were received only in draft form and were not suitable for consideration at that time. The Committee therefore asked for the final reports and for further data to clarify the extent, mechanism and toxicological consequences of the increased serum uric acid concentrations observed in humans exposed to D-tagatose. At its present meeting, the Committee reviewed the reports of the two studies in rats, the results of a study in volunteers (on the relevance of the glycogen deposition and liver hypertrophy) and some published studies on the increased uric acid concentrations in serum after intake of D-tagatose, other sugars and other food components.

Biological data: Review of the results of the studies considered by the Committee at its fifty-fifth meeting and comparisons with the data reviewed at the present meeting revealed a difference in sensitivity between Wistar and Sprague-Dawley rats. Sprague-Dawley rats given D-tagatose at a concentration of 50g/kg of diet for 28 days showed increased hepatic glycogen only when they had not been fasted the night before necropsy, and this effect was not associated with any microscopic changes in the liver. In a 90-day study in which Sprague-Dawley rats were killed after fasting overnight, administration of D-tagatose at a concentration of 50 g/kg of diet had no adverse effect on the liver. In a 6-month study in Wistar rats in which the animals were killed after fasting 7, 14 and 28 days and 3 and 5 months after treatment, administration of D-tagatose at concentrations of up to 100g/kg of diet had no adverse effects. Wistar rats are therefore less susceptible to the hepatic effects of D-tagatose than Sprague-Dawley rats. As D-tagatose stimulated glycogen deposition to a similar degree in the two rat strains in short-term studies, the difference is likely to occur at a later stage, during glycogen-induced or other stimulation of liver growth.

The authors suggested that the increase in normal liver mass seen in fasted rats fed diets containing 100 or 200g/kg D-tagatose is triggered by increased postprandial storage of liver glycogen resulting from simultaneous feeding of D-tagatose and glucose equivalents. In order to test this hypothesis, the effects of separate and simultaneous administration of D-tagatose and glycogen precursors on liver weight and glycogen level were investigated in Wistar and Sprague-Dawley rats. The results neither supported nor invalidated the hypothesis.

As several studies have been performed in healthy volunteers and in patients with diabetes, the number of persons varying from 4 to 73, the Committee based its toxicological evaluation on the data from these studies. The length of these studies varied from several days to several weeks; one study of 12 months' duration included only a limited number of patients with type 2 diabetes. The toxicological aspects investigated included gastrointestinal effects, increased serum uric acid concentrations and hepatic effects.

Mild gastrointestinal symptoms were reported in only one study, in 3 of 10 patients with type 2 diabetes receiving D-tagatose at 10 g/day for several days, whereas in other studies diarrhoea was observed only in patients receiving 25 g three times daily for 8 weeks. In healthy individuals, administration of a single dose of 30 g induced diarrhoea in some persons only, whereas other studies showed no laxative effect of single doses of D-tagatose as high as 75 g.

The serum or plasma concentration of uric acid was increased transiently in some studies, but the increased uric acid concentration was above the normal range for a number of days in only one study of persons receiving 75 g/day. The other studies showed either no increase or a transient increase in serum uric acid concentrations within the normal range.

In a 28-day study in which 15 g Of D-tagatose or 15 g of sucrose were given three times daily to volunteers, magnetic resonance imaging was used to determine liver volume, and glycogen concentrations and several clinical chemical parameters were measured⁴. The results did not reveal any relevant effect on the liver. In addition, no diarrhoea and no increase in serum uric acid concentration were observed. Therefore, the NOEL was 45 g/person per day, equivalent to 0.75 g/kg of body weight per day (for a person weighing 60 kg).

The Committee considered the 28-day study in which humans received a daily dose of 45 g of D-tagatose or sucrose in three divided doses as most representative of human dietary intake and therefore most relevant for assessing the acceptable intake of D-tagatose accurately. While effects were observed after administration of a single dose of 75 g, no effects were seen following administration of three daily doses of 15 g of D-tagatose, equivalent 0.75 g/kg of body weight per day. The Committee established an ADI of 0-80 mg/kg of body weight on the basis of this NOEL and a safety factor of 10.

Assessment of intake: D-tagatose is proposed for use as a bulk sweetener in low-energy foods, such as edible ices (at a concentration of 3 g/kg), chewing-gum and confectionery (at 15 g/kg), breakfast cereals (at 15 g/kg) and soft drinks (at 1 g/kg). At its present meeting, the Committee considered that the predicted intakes of D-tagatose determined at the fifty-fifth meeting, which were based on the manufacturers' proposed levels of use and individual dietary records in several countries, were conservative. This was because use had been assumed in the entire food category rather than only in the low-energy food component. The mean consumer intakes of D-tagatose from all proposed uses (except chewing-gum, dietary supplements and meal replacements) predicted for Australia, the Member States of the European Union and the USA ranged from 3 to 9 g/day (63-190% of the ADI), and the predicted intakes by persons at high percentiles of consumption were up to 18 g/day (375% of the ADI). On the basis of the information on possible uses, the Committee concluded that the ADI for D-tagatose may be exceeded by some groups of the population.

A toxicological monograph was prepared. The specifications prepared by the Committee at its fifty-fifth meeting were maintained.

REVIEW OF THE NEW TOXICOLOGY DATA

Chronic toxicity and carcinogenicity study with D-tagatose and fructose in Wistar rats.
Report Number V 4533 by Lina BAR and Kuper CF. TNO Nutrition and Food Research, The Netherlands, August 2002.

Test materials:	D-tagatose; 20% fructose
Control materials	pregelatinised potato starch
Test Species:	Young (5-6 weeks old) Wistar albino rats (HsdCpb:WU), 50 male and 50 female per group dosed by administration in diet.
Dose:	0, 2.5, 5 or 10% w/w in diet for 24 months.
Guidelines:	USFDA 1982-Toxicological Guidelines, OECD Guideline for Testing of Chemicals 451, May 1981 and EC Guideline 88/302/EC.

⁴ Boesch C, Ith M, Jung B, Bruegger K, Erban S, Diamantis I, Kreis R, Bar A, *Effect of oral D-tagatose on liver volume and hepatic glycogen accumulation in healthy male volunteers*, **Regul Toxicol Pharmacol** 2001 **Apr;33(2):257-67**

Test articles and control material

D-tagatose and fructose were obtained as a crystalline white powders with > 99% purity. The control substance was a pregelatinised potato starch as a coarse white powder.

Study conduct

Four groups of rats (50/sex/group) were treated with D-tagatose in the diet at 2.5, 5, 10 or 10% D-tagatose + 10% fructose (equivalent to 1.0, 2.0 or 4.0 g/kg bw/day for males and 1.2, 2.5 and 4.9 g/kg bw/day for females). The test substance was incorporated in the feed at the expense of 20% barley. The control, low-dose and mid-dose diets were compensated by adding respectively 20%, 17.5%, 15% and 10% pregelatinised potato starch.

Clinical observations were recorded daily and bodyweight were measured weekly for the first 13 weeks and subsequently once every month. From 6 months after the start of the study until the end of the study, the animals were palpated weekly to detect palpable masses. Food consumption was assessed on a cage basis, by weighing the feeders, over each 1-week period during the first 13 weeks and subsequently over 1-week periods every month. Blood samples for haematology were taken from all rats after 12 and 24 months. At the end of the study, all animals were sacrificed and a complete necropsy performed (gross examination, organ weights and tissue sampling). Histopathology was performed on adrenals, liver and kidneys of all rats.

Results

The highest average intake of the test substance was 4.0g/kg bw/day in males and 5.0g/kg bw/day in females of the highest dose group. There were no dose related differences in appearance, general condition or behaviour among the groups. The incidence of grossly visible or palpable masses was not considered treatment-related effect. The test substance did not affect mortality rate.

Mean body weights decreased in males and females at the highest dose. The decrease was statistically significant but there was no dose-response relationship. There were no noticeable differences in overall food intake.

Haematology showed that haemoglobin concentration and packed cell volume tended to be decreased in females fed 10% D-tagatose throughout the study. There were no significant differences in clinical chemistry variables among the groups.

There was a statistically significant increase in the relative weight of the liver in females given the highest doses of D-tagatose and fructose. The relative weights of the kidneys, adrenals and filled and empty caecums were also increased in females and males to various degrees at higher doses of D-tagatose and fructose.

Macroscopic examination at necropsy showed a relatively high incidence of enlargement of the adrenals in both sexes in all groups including the fructose groups. Microscopic examination did not reveal any treatment-related histopathological changes. There were no observable differences between the treated and control animals in terms of liver tumour incidence.

Conclusions

Based on this 2-year study in which the rats were administered with D-tagatose in the diet for up to 20%, there was no evidence that the test substance is carcinogenic to these animals.

There were increases in the relative weights of liver in both D-tagatose and fructose groups but this could be ascribed to additional growth of hepatocytes in reaction to increased glycogen deposition, and is not considered to represent an adverse effect. All other clinical and haematological parameters remained normal. Macroscopic and microscopic examinations did not reveal any treatment-related gross or histopathological changes. Based on the above findings, it is concluded that long-term exposure to D-tagatose at dietary levels of up to 10% (4g/kg/bw for males and 5g/kg/bw for females) does not lead to adverse long-term consequences on the liver of rats. Based on this study a NOEL of 4g/kg/bw/d

Effect of an oral 30-g dose of D-tagatose on the plasma uric acid levels of healthy male volunteers

by Diamantis I and Bar A. Department of Internal Medicine, University of Crete, Heraklion, Greece, *November 2001.*

Test materials:	D-tagatose
Subjects:	6 male volunteers between 22 and 24 years of age and having body weights of 62-65 Kg.
Dose:	marmalade containing 30g D-tagatose for breakfast.
Guidelines:	in accordance with the declaration of Helsinki approved by the Ethics Committee of the University Hospital of Heraklion.

Test articles and control material

D-tagatose was obtained as a powder with > 99% purity. Marmalade was prepared with a D-tagatose content of about 40% and a fruit content of about 60%.

Study conduct

Blood samples were collected from the volunteers 1 hour prior to providing them with a light breakfast and blood collection continued just before and up to 4 hours after breakfast. The subjects were required to avoid alcohol the previous night. Six after the breakfast, clinical examination was performed on the subjects. Urine was collected over a 24-hour period starting from intake of the test breakfast.

Plasma and urine uric acid concentrations were determined. Plasma glucose and plasma inorganic phosphate were also measured within one hour of blood and urine collection.

Results

All 6 subjects completed the study according to protocol. Plasma glucose concentrations showed statistically significant increase at 1 hour after test breakfast. Plasma uric acid concentrations showed statistically significant increase in all 6 subjects at all postprandial times but no correlation to effect of time (1h to 4h) and they remained within the normal range at all times. Postprandial plasma phosphate levels showed statistically significant decrease 1-2 hour after breakfast, but increased towards the end of the observation period. The levels remained within the normal range throughout the observation period.

The 24-hour urinary excretion of uric acid was within the normal range.

No side effects (flatulence, bloated feeling and laxation) were reported up to 24-hour post treatment observation period.

Conclusions

This study confirms earlier findings using animal models that D-tagatose produces an increase in the plasma uric acid concentrations in humans, however the uric acid levels remain well within the normal range after ingestion of 30 g D-tagatose.

Effect of an oral 15-g dose of D-tagatose on the plasma uric acid levels of hyperuricemic male volunteers

by Diamantis I and Bar A. Department of Internal Medicine, University of Crete, Heraklion, Greece, *December 30, 2002.*

Test materials:	D-tagatose
Subjects:	12 male volunteers (7 hyperuricemic and 5 with gout) between 57 and 64 years of age and having body weights of 72-84 Kg.
Dose:	marmalade containing 15g D-tagatose for breakfast.
Guidelines:	in accordance with the declaration of Helsinki approved by the Ethics Committee of the University Hospital of Heraklion.

Test articles and control material

D-tagatose was obtained as a powder with > 99% purity. Marmalade was prepared with a D-tagatose content of about 40% and a fruit content of about 60%.

Study conduct

Blood samples were collected from the volunteers 1 hour prior to providing them with a light breakfast and blood collection continued just before and up to 4 hours after breakfast. The subjects were required to avoid alcohol the previous night. Six after the breakfast, clinical examination was performed on the subjects. Urine was collected over a 24-hour period starting from intake of the test breakfast.

Plasma and urine uric acid concentrations were determined. Plasma glucose, plasma inorganic phosphate, creatinine and lactate were also measured within one hour of blood and urine collection.

Results

All 12 subjects completed the study according to protocol. Plasma glucose concentrations showed statistically significant increase from 0.5 to 2 hours after test breakfast. Plasma uric acid concentrations showed statistically significant increase in all 12 subjects at all postprandial times but they remained small and not different from that observed during the pre-treatment period at all times. Postprandial plasma phosphate levels did not vary in response to the intake of test breakfast.

The 24-hour urinary excretion of uric acid was above the normal value in three subjects.

No side effects (flatulence, bloated feeling and laxation) were reported up to 24-hour post treatment observation period.

Conclusions

This study shows that a single dose of 15 g D-tagatose does not produce a clinically significant increase of plasma uric acid concentrations in hyperuricemic and gouty subjects.

ADDITIONAL PUBLISHED LITERATURE ON THE SAFETY/ METABOLISM OF D-TAGATOSE IN HUMANS

<p>Normen L, Laerke HN, Jensen BB, Langkilde AM, Andersson H, <i>Small-bowel absorption of D-tagatose and related effects on carbohydrate digestibility: an ileostomy study</i>, Am J Clin Nutr 2001 Jan;73(1):105-10.</p>	<p>The aims of this study were to measure the excretion of D-tagatose from the human small bowel, to calculate the apparent absorption of D-tagatose, and to study the effects of D-tagatose on the small-bowel excretion of other carbohydrates. It was concluded that the apparent absorption of 15 g D-tagatose/d was 81%. D-Tagatose had only a minor influence on the apparent absorption of other nutrients.</p>
<p>Buemann B, Gesmar H, Astrup A, Quistorff B, <i>Effects of oral D-tagatose, a stereoisomer of D-fructose, on liver metabolism in man as examined by 31P-magnetic resonance spectroscopy</i>, Metabolism 2000 Oct;49(10):1335-9.</p>	<p>The effect of 30 g D-tagatose or D-fructose administered orally on ketohexose-1-phosphates, ATP, and inorganic phosphate (Pi) levels in the liver was studied by 31P-magnetic resonance spectroscopy (PMRS) in 5 young male volunteers. Blood and urine were collected to detect a possible increased uric acid production. The results suggest that a moderate intake of D-tagatose may affect liver metabolism by phosphate trapping despite the fact that the sugar may only be incompletely absorbed in the gut.</p>
<p>Buemann B, Toubro S, Astrup A., <i>Human gastrointestinal tolerance to D-tagatose</i>, Regul Toxicol Pharmacol 1999 Apr;29(2 Pt 2):S71-7</p>	<p>The gastrointestinal effects of D-tagatose on after the consumption of 29 or 30 g of D-tagatose was investigated. Nausea and diarrhea were reported with an incidence of 15.1 and 31.5%, respectively, in 73 healthy young male subjects in a screening study. Increased flatulence after D-tagatose was frequently reported in all the studies and the flatulence did not decline during a 15-day period with intake of 30g in one dose daily. In most cases, symptoms were reported as light or moderate. The results suggest that 30g taken at one time may be above the recommended dose for ordinary use.</p>
<p>Buemann B, Toubro S, Raben A, Blundell J, Astrup A, <i>The acute effect of D-tagatose on food intake in human subjects</i>, Br J Nutr 2000 Aug;84(2):227-31.</p>	<p>A double-blind randomized crossover study was performed with nineteen normal-weight men to investigate the effect on subsequent ad libitum food intake of replacing 29 g sucrose with 29 g D-tagatose as sweetener to a breakfast meal. The results from this study suggest that D-tagatose may contribute to a reduced energy intake.</p>
<p>Buemann B, Toubro S, Holst JJ, Rehfeld JF, Bibby BM, Astrup A, <i>D-tagatose, a stereoisomer of D-fructose, increases blood uric acid concentration</i>, Metabolism 2000 Aug;49(8):969-76.</p>	<p>The effect of 30 g oral D-tagatose versus D-fructose on plasma uric acid and other metabolic parameters was tested in 8 male subjects by a double-blind crossover design. Serum uric acid concentration was significantly higher after D-tagatose compared with either 30 g D-fructose or plain water. D-Tagatose attenuated the glycaemic and insulinemic responses to a meal that was consumed 255 minutes after its administration. Moreover, both fructose and D-tagatose increased plasma concentrations of cholecystokinin (CCK) and glucagon-like peptide-1 (GLP-1). The metabolic effects of D-tagatose occurred despite its putative poor absorption.</p>

<p>Laerke HN, Jensen BB, Hojsgaard S, <i>In vitro</i> fermentation pattern of D-tagatose is affected by adaptation of the microbiota from the gastrointestinal tract of pigs, J Nutr 2000 Jul;130(7):1772-9.</p>	<p><i>In vitro</i> fermentation experiments with pig intestinal contents and bacteria harvested from the gastrointestinal tract of pigs were used to investigate the degradation of D-tagatose and the formation of fermentation products. Two groups of eight pigs were fed either a control diet containing 150 g/kg sucrose or a diet which had 100 g/kg of the sucrose replaced by D-tagatose. After 18 d the pigs were killed and the gastrointestinal contents collected for <i>in vitro</i> studies. No microbial fermentation of D-tagatose occurred in the stomach or in the small intestine, whereas the sugar was fermented in the caecum and colon. The authors suggest that D-tagatose is not fermented in the upper gastrointestinal tract, and the ability of the large intestinal microbiota to ferment D-tagatose is dependent on adaptation.</p>
---	--

Conclusion of human studies assessed by JECFA, FSANZ and those published in literature

A survey of the literature suggests that the consumption of D-tagatose causes transient increase in uric acid levels but remains within the normal range. No significant toxicological effects were observed in human studies.

JECFA assessment was based on the concern about its potential to induce glycogen deposition and hypertrophy in the liver and to increase the concentrations of uric acid in serum. At that time, the effect of D-tagatose had not been studied in people prone to higher serum uric acid concentrations. However new data available to FSANZ indicates that there is no evidence of any public health and safety concern associated with consumption of D-tagatose up to 15g/d (0.25g/kg bw/d) even in susceptible groups, hyperuricemic and gouty subjects.

Further the application of 10-fold safety factor by JECFA on bulk ingredients is inappropriate, as they cannot be tested above 40-50g/d due to laxation effects. The new data on humans readily demonstrate the safety of D-tagatose consumption by humans up to 15g/person/day. Therefore no further safety factor needs to be applied in determining its ADI.

OVERALL CONCLUSIONS

JECFA established an ADI of 0–80 mg/kg bw (~ 45g/person/day) on the basis of NOEL from human studies and using a safety factor of 10. However bulk ingredients such as D-tagatose cannot be tested above 40-60g/d since low-digestible carbohydrates induce intestinal side effects such as laxation at high doses. Further, assessment of new data submitted by the applicant confirms that there is no evidence of any public health and safety concern associated with consumption of D-tagatose up to 15g/d even in susceptible groups (hyperuricemic and gouty subjects). Another human study demonstrated that D-tagatose produces a transient increase in the plasma uric acid concentrations in humans, however the uric acid levels remain well within the normal range after ingestion of 30 g D-tagatose. Therefore applying a 10-fold safety factor in this case is not appropriate.

There were increases in the relative weights of liver in rats fed D-tagatose but this is because of additional growth of hepatocytes in reaction to increased glycogen deposition, and is not considered a significant toxicological effect. This effect has also been observed with fructose and there were no histopathological changes or adverse long-term consequences of glycogen-induced liver enlargement in these rats. There is some evidence that gastrointestinal effects such as nausea, diarrhoea and flatulence can occur in some individuals at doses 30g/d and above.

It is concluded therefore that D-tagatose is safe for use in food for all individuals up to 15g/day. Mild GI effects may be observed in some individuals at higher dose levels.

FOOD TECHNOLOGY REPORT


D-TAGATOSE AS A NOVEL FOOD

INTRODUCTION

Structure, function and occurrence

D-tagatose is a naturally occurring sweet-tasting monosaccharide. D-tagatose is a ketohexose in which its fourth carbon is chiral and is a mirror image of the respective carbon atom of the common D-sugar, fructose. The CAS number for D-tagatose is 87-81-0. The empirical formula for D-tagatose is $C_6H_{12}O_6$. The molecular weight for D-tagatose is 180.16.

The structural formula of tagatose is:



D-tagatose occurs naturally in *Sterculia setigera* gum, a partially acetylated acidic polysaccharide. D-tagatose is also found in heated cow's milk. Sterilised cow's milk and milk powder were found to contain tagatose at concentrations between 2 to 20 and 800 ppm, respectively (Troyano et al., 1996).

Production

The production process of D-tagatose occurs in a stepwise manner, starting from the raw material lactose (a disaccharide), which is then altered with the use of enzymes and various fractionating, isomerisation, and purification techniques. Food grade lactose is hydrolyzed to galactose and glucose by passing the solution through an immobilized lactase column.

The sugar mixture from the enzyme hydrolysis is fractionated by chromatography. The chromatographic separation of glucose and galactose is essential and is similar to the normal industrial separation of glucose and fructose, using calcium-based cationic resins. The galactose fraction from the chromatography is converted to D-tagatose under alkaline conditions by adding a suspension of $\text{Ca}(\text{OH})_2$ and, optionally, a catalyst CaCl_2 . The reaction is stopped by adding sulphuric acid. D-tagatose is stable under the conditions of the isomerisation process. The resulting filtrate is further purified by means of demineralization and chromatography. The purified D-tagatose solution is then concentrated and crystallized to give a white crystalline product which is more than 99% pure.

Physical properties

D-tagatose is 92% as sweet as sucrose in a 10% aqueous solution (Table 1). D-tagatose is a white crystalline powder with an appearance very similar to sucrose. The main difference is that the crystal form is a tetragonal bipyramid. Crystallisation from aqueous solution results in anhydrous crystals in an α -pyranose form, with a melting point of 134-137 °C.

Table 1. Relative Sweetness of Tagatose and other sugars and polyols.

	Relative Sweetness
Sucrose	1
Fructose	0.8-1.7
Tagatose	0.92
Xylitol	0.8-1.0
Glucose	0.5-0.8
Erythritol	0.50
Sorbitol	0.4-0.7
Lactitol	0.4

D-tagatose is a nonhygroscopic product similar to sucrose. This means that D-tagatose will not absorb water from its surrounding atmosphere under normal conditions and does not require special storage.

D-tagatose exerts a greater osmotic pressure, and hence a lower water activity than does sucrose at equivalent concentrations. The effect of water activity is similar to fructose due to the same molecular weight. Water activity influences product microbial stability and freshness.

D-tagatose is similar to sucrose in its solubility in water. This makes it suitable for applications where it is substituted for sugar. Compared with polyols, tagatose is more soluble than erythritol and less soluble than sorbitol.

D-tagatose solutions are lower in viscosity (180 cP at 70% w/w and 20 °C) than sucrose solutions at the same concentrations but slightly higher than fructose and sorbitol.

Chemical properties

D-tagatose is a reducing saccharide and takes part in Maillard reactions, which leads to a distinct browning effect in food. It also decomposes (caramelizing) more readily than sucrose at high temperatures. At low and high pH, D-tagatose is less stable and converts to various compounds. However, D-tagatose can be used satisfactorily in many different applications at high temperature when process time is kept short.

Applications

D-tagatose is well suited for confectionery products such as chocolate and hard boiled candies, fondants, fudges, and caramels due to its similar sweetness to sucrose, good ability to crystallize and low caloric value. It could also have applications in ice-cream, soft drinks, and breakfast cereals.

Table 1 Conditions of Use Proposed by Applicant	
Food Category	<i>Use Level in Food (%)</i>
Breakfast cereals (ready to eat)	4-20 (depending on serving size)
Carbonated diet soft drinks	1
Non-carbonated diet drinks	1
Low fat/fat free Ice-cream	3
Low fat frozen dairy desserts	3
Diet/Health bars	10
Diet soft confectionery	10
Hard confectionery	10-15
Chewing gum	60
Icings/frostings	30
Special purpose foods/Meal replacements	15-20g per serving

Conclusion

D-tagatose is a naturally occurring sweet-tasting monosaccharide that may be used in a wide range of foods.

References

Lyn O'Brien Nabors (Ed). Alternative Sweeteners. Marcel Dekker, Inc., New York, 3rd Edition 2001.

Troyano E., Villamiel M., Olano A., Sanz J. and Martinez-Castro, I. (1996) Monnosaccharides and myo-inositol in commercial milks. J. Agric. Food Chem. 44: 815-817.

DIETARY EXPOSURE ASSESSMENT REPORT

D-TAGATOSE AS A NOVEL FOOD

An application was received by FSANZ in July 2002 requesting approval of the monosaccharide D-tagatose for use as a novel food ingredient in a range of low fat, fat free, diet and reduced energy type and special purpose processed foods. A dietary exposure assessment was conducted to predict exposure of Australians and New Zealanders to D-tagatose when used in foods in accordance with those specified by the applicant. The applicant requested a general permission at GMP to permit its use in all food groups and not just for those specific foods listed in the application.

Summary

Estimated dietary exposures were calculated on a chronic and single eating occasion basis for the Australian and New Zealand populations and Australian children aged 2-4 years and 5-12 years.

Estimated mean chronic dietary exposures to D-tagatose for consumers were 2.7 g/d for the Australian population (2 years and above), 1.9 g/d for Australian toddlers 2-4 years, 3.0 g/d for Australian children aged 5-12 years and 1.9 g/d for the New Zealand population (15 years and above). Estimated 95th percentile exposures to D-tagatose for consumers were 9.0 g/d for the Australian population, 6.9 g/d for children 2-4 years, 10.5 g/d for children 5-12 years and 7.4 g/d for the New Zealand population. Ninety-fifth percentile exposures are overestimated as 24-hour food consumption data exaggerates habitual exposures. These exposures are also based on maximum proposed use levels, also producing an overestimate of exposure.

Estimated short term exposures for high consumers in a single eating occasion were calculated for individual food groups to assess whether gastro-intestinal effects were likely. All estimated short term exposures were 24 grams or less for any food for any population group, except for the Australian population consuming large amounts of formulated dietary foods (59 g D-tagatose/day), or formulated supplementary sports foods (56 g D-tagatose/day).

Dietary Exposure Assessment provided by the applicant

The applicant provided a detailed dietary exposure assessment for D-tagatose, based on food groups and levels of use similar to those being proposed for Australia and New Zealand. However this assessment was not considered to be sufficient for assessing the safety of potential exposure to D-tagatose in Australia and New Zealand as the assessment, although detailed, was based on United States food consumption data (US Department of Agriculture Continuing Survey of Food Intakes by Individuals, 1994-96). Therefore FSANZ conducted a dietary exposure assessment.

The exposure assessment submitted by the applicant indicated that mean daily exposure to D-tagatose for consumers of foods containing added D-tagatose (excluding chewing gum and formulated diet foods) was 5 g/day and 11 g/day for the 90th percentile consumer.

Expressed in terms of grams per kilogram body weight per day (g/kg bw/d), the highest exposure is projected for preschool children aged 2-5 years with mean and 90th percentile exposures of 0.17 and 0.33 g/kg bw/d respectively. For the consumers of all age groups combined the estimated exposure is 0.08 and 0.18 g/kg bw/d for the mean and 90th percentile respectively. The additional exposure from chewing gum is estimated at 0.06 – 0.07 g/kg bw/d for the average consumer (assuming a use level of D-tagatose of 60% in the gum). The average daily exposure to D-tagatose from formulated diet foods was considered separately by the applicant as such products are intended to be consumed as meal replacements or as a snack in the context of weight loss or weight control diet. The average daily exposure to D-tagatose from this use is estimated at 0.08 – 0.16 g/kg bw/d for the 20+ year old consumer who consumes one or two servings of a formulated diet food per day. US food consumption data are based on two days of food records. Use of multiple day records tends to significantly reduce the predicted high consumer exposure (Rutishauser, 2000).

The USFDA estimated the dietary exposure to D-tagatose under the intended conditions of use to be 7.5 g/day for the mean consumer and 15 g/day for the 90th percentile.

The applicant indicated that these estimated exposure levels are likely to be an overestimate since it was presumed that D-tagatose would be used in all foods at the maximum use levels.

The exposure assessment submitted in the application did not estimate exposure to D-tagatose per single eating occasion (e.g. per meal or per snack). Additional information submitted by the applicant did contain data on exposure ‘per eating occasion’ (e.g. breakfast, lunch, dinner and snack). Per eating occasion for the population (2 years and above), mean exposures were 4.3 grams and 90th percentile exposures were 7.2 grams.

Dietary Modelling

The FSANZ dietary exposure assessment was conducted using dietary modelling techniques that combine food consumption data with food chemical concentration data to estimate the exposure to the food chemical from the diet. The dietary exposure assessment was conducted using FSANZ’s dietary modelling computer program, DIAMOND.

$$\boxed{\text{Dietary exposure} = \text{food chemical concentration} \times \text{food consumption}}$$

The exposure was estimated by combining usual patterns of food consumption, as derived from national nutrition survey (NNS) data, with proposed levels of use of D-tagatose in foods.

The chronic dietary exposure assessment is based on a range of food sources of added D-tagatose in the diet. That is, it includes where consumers may have only eaten one food containing D-tagatose or many foods containing D-tagatose to assess chronic exposures. The exposure assessment also assessed individual foods to determine the likelihood of short term effects of exposure.

Naturally occurring sources of D-tagatose were not included in the exposure assessment as it was assumed that they would not contribute substantially to the total exposure to D-tagatose.

Dietary Survey Data

DIAMOND contains dietary survey data for both Australia and New Zealand; the 1995 NNS from Australia that surveyed 13 858 people aged 2 years and above, and the 1997 New Zealand NNS that surveyed 4 636 people aged 15 years and above. Both of these surveys used a 24-hour food recall methodology.

The dietary exposure assessment was conducted for both Australian and New Zealand populations. Modelling was conducted for the whole population, as well as for 2 – 4 year olds and 5 – 12 year olds (for Australia only). An exposure assessment was conducted on these younger age groups because children generally have higher exposures due to their smaller body weight, they consume more food per kilogram of body weight compared to adults and because they tend to consume a lot of the food types that are proposed to contain D-tagatose, such as ready-to-eat breakfast cereals, chewing gum, ice creams, gum (soft) type sweets, confectionery and beverages.

D-Tagatose Concentration levels

The levels of D-tagatose in foods that were used in the exposure assessment were from the application (Table to part 2 (c)). The list of foods and the proposed levels of use, as provided by the applicant, are shown below in Table 1. However, this may not be an extensive list of all foods that D-tagatose could be used in if a general permission for use is given, as D-tagatose can be used as a replacement for other sugars in many foods.

Alternatively sweetened food categories were used, where available, for the exposure assessment. There was no food consumption data available from the NNS for alternatively sweetened ready-to-eat processed breakfast cereals. Based on information supplied by the applicant, that D-tagatose is only likely to be used in reduced fat / reduced-energy type foods, it is unrealistic to assume that D-tagatose would be used in all breakfast cereals. Therefore, FSANZ used the Australian market share for ‘healthy’ breakfast cereals (11%) (Retail World, 2002) as a more realistic estimate of the amount of cereals that could contain D-tagatose. Where a range of concentrations for a food was reported by the applicant, the highest one was used in the exposure assessment in order to assume a worst-case scenario. The concentration of D-tagatose used in the exposure assessment is shown in Table 2.

Table 1: Proposed use of D-Tagatose in foods and levels of use

Food Name	Proposed use level in food	Concentration Level (mg/kg)
Low fat / Fat-free ice cream	3 %	30 000
Chewing gum	60 %	600 000
Diet soft confectionery	10 %	100 000
Hard confectionery	10 – 15 %	100 000 – 150 000
Icings / Frostings	30 %	300 000
Breakfast cereals ready-to-eat	4 – 20% (depending on serve size)	40 000 – 200 000
Special purpose foods / Meal replacements	(15 – 20 g per serve)	*
Non-carbonated diet drinks	1 %	10 000
Carbonated diet soft drinks	1 %	10 000
Low fat frozen dairy desserts	3 %	30 000
Diet / Health bars	10 %	100 000

* depends on serve size assigned. See Table 2.

Table 2: Food groups and concentrations of D-Tagatose used in the FSANZ dietary exposure assessment (DEA)

Food Code	Food Name	Concentration Level (mg/kg)	Foods included	Assumptions
3.1.1.8	Low fat / Fat reduced ice cream	30 000	Carbohydrate modified ice cream products, ice confection	
5.2.1	Bubble gum and chewing gum	600 000		
5.3.2	Gum (soft) type sweets confectionery	100 000	Jubes, jellies, Turkish delight, marshmallow	
5.3.3	Hard boil sugar confectionery	150 000		
5.4	Icings and Frostings	300 000		
6.3	Processed cereal and meal products	22 000	“Healthy” breakfast cereals	DEA assumed market share of 11%
13.3	Formula dietary foods	250 000	Diet bars / biscuits	Based on 80 g serve size and 20 g D-tagatose / serve
13.4	Dietetic formulae for slimming and weight loss	53 000		13.4-13.6 Based on a 375g serve size and 20 g D-tagatose / serve
13.5	Supplementary foods for dietetic uses	53 000	Meal replacement drinks	
13.6	Formulated supplementary sports foods	53 000	Sports drinks, high protein drinks	
14.1.3.3	Water based flavoured drinks, artificially sweetened	10 000	Cordial, soft drink	
20.2.2.1	Dairy desserts, artificially sweetened	30 000	Mousse, puddings	Fat reduced chilled yogurt products were not included
20.2.3	Confectionery	100 000	Health bars, muesli bars, chocolate coated health bars, breakfast bars	

A concentration level of 22,000 mg/kg was used for ready-to-eat processed breakfast cereals. Because FSANZ does not have specific consumption data for alternatively sweetened breakfast cereals, a value of 11% (based on market share data) of the total food category’s level of use was used, i.e. eleven percent of 200,000 mg/kg (proposed maximum level of use as indicated by the applicant for ready-to-eat breakfast cereals). For the special purpose foods / meal replacements food category, a concentration value of 53 000 mg/kg was used for liquid foods and 250 000 mg/kg for solid foods in this category. This is based on a 375 g serve size for liquid foods and an 80 g serve for solid foods and the upper level of use of 20 grams/serve (as indicated by the applicant) of D-tagatose in these types of products.

How the estimated dietary exposures were calculated

The DIAMOND model allows D-tagatose concentrations to be assigned to food groups. The concentration values in Table 2 were assigned to the alternatively sweetened sub-group of the main food category where a sub category existed. For example, only artificially sweetened carbonated and non-carbonated drinks were assigned the concentration value of 10 000 mg/kg. Otherwise, the whole group of the normal counterpart of the food group was assigned the D-tagatose concentration (e.g. all icings and frostings).

Individuals' exposure to the additive was calculated using their individual food records from the dietary survey. The DIAMOND program multiplies the specified concentration of D-tagatose by the amount of food that an individual consumed from that group in order to estimate the exposure to each food. Once this has been completed for all of the foods specified to contain D-tagatose, the total amount of D-tagatose consumed from all foods is summed for each individual. Population statistics (mean and high percentile exposures) are then derived from the individuals' ranked exposures.

Where estimated dietary exposures are expressed per kilogram of body weight, each individuals' total dietary exposure is divided by their own body weight, the results ranked, and population statistics derived.

Percentage contributions of each food group to total estimated exposures are calculated by summing the exposures for a food group from each individual in the population group who consumed a food from that group and dividing this by the sum of the exposures of all individuals who consumed from any of the food groups containing D-tagatose and multiplying this by 100.

Food consumption amounts for each individual take into account where each food in a classification code is consumed alone and as an ingredient in mixed foods. For example, breakfast cereal consumed as is, or as part of a slice, are included in the consumption of breakfast cereal.

Estimated exposures for single food groups were calculated using the 95th percentile consumption amount for the food group derived by DIAMOND and multiplying this by the concentration of D-tagatose in the food.

Assumptions in the dietary modelling

Assumptions made in the dietary modelling include:

- Where a permission is given to a food category, all foods in that category contain D-tagatose at the specified amount;
- All foods in a group will contain D-tagatose at the maximum proposed use level for the group;
- Mean body weight for Australian respondents aged 2 – 4 years is 17 kg;
- Mean body weight for Australian respondents aged 5 – 12 years is 32 kg;
- Mean body weight for Australian respondents aged 2 years and above is 66 kg;
- Mean body weight for New Zealand respondents aged 15 years and above is 71 kg;
- Consumption amounts of foods as recorded in the NNSs represent current consumption patterns; and
- Consumers always selected the product in the food group that contained D-tagatose.

These assumptions (other than those relating to body weight) are likely to lead to a conservative estimate for D-tagatose dietary exposure. This is because it is unlikely that all foods permitted to contain D-tagatose would in fact contain it, and if they did contain D-tagatose it is unlikely to always be added at the maximum amounts outlined in the application. However the application did not contain information on the likely market share of D-tagatose-containing foods.

The applicant indicated that D-tagatose is expensive in comparison to sugar and is unlikely to be used in large quantities in an extensive range of foods.

Limitations of the dietary modelling

A limitation of estimating dietary exposure over a period of time associated with the dietary modelling is that only 24-hour dietary survey data were available, and these tend to over-estimate habitual food consumption amounts for high consumers. Therefore, predicted high percentile exposures are likely to be higher than actual high percentile exposures over a lifetime.

In addition, the NNS data used in DIAMOND do not distinguish between eating occasion (e.g. breakfast, lunch, dinner) so it is not possible to report the amount of D-tagatose eaten in one meal/snack. For example, the 240 g of low fat / reduced fat ice cream reportedly eaten by Australian high consumers (95th percentile, see Table 6 below) could have been eaten at one time or over two or more occasions during the day (lunch and dinner, for example). For the purpose of the exposure assessment it is assumed that this amount is eaten at one meal, which is the worst-case scenario.

RESULTS

Estimated dietary exposures to D-tagatose

The estimated dietary exposures to D-tagatose are shown in Table 4. Results are reported for all respondents (i.e. regardless of whether D-tagatose was consumed or not), and for consumers only (i.e. only those individuals that consumed D-tagatose). Only consumer exposures are discussed below in detail because this is the group of the population FSANZ needs to ensure has safe exposure.

The estimated mean exposure for consumers of D-tagatose for the Australian population aged 2 years and above is 2.7 g/day. The estimated mean exposure for consumers of D-tagatose for Australian toddlers aged 2 – 4 years is 1.9 g/day and for children aged 5 – 12 years it is 3.0 g/day. The estimated mean exposure for consumers of D-tagatose for the New Zealand population aged 15 years and above is 1.9 g/day. The estimated 95th percentile exposures for consumers of D-tagatose for the total Australian and New Zealand populations are 9.0 g/day and 7.4 g/day respectively. The estimated 95th percentile exposures for consumers of D-tagatose for Australian toddlers aged 2 – 4 years and children aged 5 – 12 years are 6.9 g/day and 10.5 g/day respectively.

Table 4: Estimated dietary exposures to D-tagatose

Country	Age group	Number of consumers of D-tagatose	Consumers as a % of total respondents [#]	Grams per day		
				Mean all respondents	Mean consumers only	95 th percentile consumers only
Australia	Whole population (2 years+)	7341	53	1.4	2.7	9.0
	2-4 years	472	81	1.6	1.9	6.9
	5-12 years	1166	78	2.4	3.0	10.5

New Zealand	Whole population (15 years+)	2311	50	0.9	1.9	7.4
--------------------	------------------------------	------	----	-----	-----	-----

Total number of respondents for Australia: whole population = 13 858, 2-4 years = 583, 5-12 years = 1 496; New Zealand: whole population = 4 636.

Major contributing foods to total estimated dietary exposures

The major foods (>5%) that contributed to the total estimated exposures to D-tagatose are displayed in Table 5. These are displayed for the total population as well as for the younger age groups. The major contributor for each population group assessed was breakfast cereals at around 30% contribution. Other major contributors were gum (soft) type sweets, hard boil sugar confectionery, supplementary foods for dietetic use, formulated supplementary sports foods, water based flavoured drinks and health bars.

The results are based on the 24-hour food recall component of the NNS. The food frequency questionnaire (FFQ) from the Australian NNS (on respondents 12 years and above) showed only 30% of adults and 43% of children 12-18 years consume breakfast cereal on a daily basis. Thirty-three percent of 12-18 year olds and adults eat cereal once to 6 times per week. Twenty-five percent of adults never/less than once per month eat breakfast cereal. As breakfast cereals are the major contributing food to D-tagatose exposure, the estimated exposures to D-tagatose could be lower than estimated on most week days over a lifetime.

Table 5: Major contributors to total D-tagatose dietary exposures for Australia and New Zealand, and for different age groups

Country	Age group	Major contributing foods and percent of total D-tagatose exposures
Australia	Whole population (2+ years)	Breakfast cereals (30%) Health bars (14%) Formulated supplementary sports foods (13%) Water based flavoured drinks (11%) Gum (soft) type sweets (9%) Hard boil sugar confectionery (8%) Supplementary foods for dietetic use (5%)
	2-4 years	Breakfast cereals (30%) Health bars (22%) Hard boil sugar confectionery (13%) Gum (soft) type sweets (12%) Supplementary foods for dietetic use (12%) Water based flavoured drinks (8%)
	5-12 years	Breakfast cereals (27%) Health bars (21%) Gum (soft) type sweets (14%) Hard boil sugar confectionery (14%) Water based flavoured drinks (6%)
New Zealand	Whole population (15+ years)	Breakfast cereals (29%) Hard boil sugar confectionery (23%) Health bars (19%) Gum (soft) type sweets (12%) Formula diet foods (5%) Supplementary foods for dietetic use (5%)

Comparison of estimated dietary exposure with other studies of D-tagatose exposure

The estimated dietary exposures calculated by FSANZ are slightly lower on a population basis than those submitted by the applicant. Differences in methodology of calculating the exposures (including a different population, consumer consumption patterns, different NNS methodology and calculations) can result in differences in estimated exposures.

The estimated exposures calculated by FSANZ are slightly lower in grams per day than JECFA estimates of 3-9 g/d from a number of countries. High consumers were estimated to be exposed to up to 18 g/d (WHO 2002). This estimate was higher than what FSANZ predicted for high consumers.

Estimated dietary exposures to D-tagatose for single food groups

The dietary exposures to D-tagatose from individual foods were calculated in order to determine whether a consumer could exceed the bolus dose reference health standard, therefore possibly encountering adverse gastro-intestinal effects. They were calculated by multiplying the 95th percentile food consumption amount for a food group by the specified concentration of D-tagatose as outlined in Table 1.

The estimated exposures are shown in Table 6 and 7 for Australia and New Zealand respectively. These dietary exposures differ from the estimated 95th percentile dietary exposures to D-tagatose referred to earlier in the report, in that the results in this section are for 95th percentile consumption amounts for single foods whereas the results earlier in the report refer to the 95th percentile dietary exposure to D-tagatose from consumption of all foods proposed to contain D-tagatose.

Table 6: Estimated dietary exposure to individual food groups at the 95th percentile level of consumption - Australia

Food code	Description	Level of use (mg/kg)	All consumers 2+ years intake		Toddlers aged 2 – 4 years		Children aged 5 – 12 years	
			P95 food intake per day (g)	Grams D-tagatose / day	P95 food intake per day (g)	Grams D-tagatose / day	P95 food intake per day (g)	Grams D-tagatose / day
3.1.1.8	Reduced fat ice cream	30 000	240	7.2	82	*	235	7.1
5.2.1	Chewing gum	600 000	24.85	14.9	18.97	*	31.97	19.2
5.3.2	Gum (Soft) type sweets	100 000	133.42	13.3	90	9.0	125	12.5
5.3.3	Hard candy	150 000	64.19	9.6	52.56	7.9	62.96	9.4
5.4	Icings/frostings	300 000	57.3	17.2	20	*	40	*
6.3	Ready-to-eat breakfast cereals	200 000	120	24.0	67.5	13.5	112.5	22.5
13.3	Formulated dietary foods	250 000	237	59.3	NC		NC	
13.5	Supplementary foods for dietetic use	53 000	40.66	2.2	48.22	2.6	23.67	1.3
13.6	Formulated supplementary sports foods	53 000	1055.3	55.9	NC		508	*

Food code	Description	Level of use (mg/kg)	All consumers 2+ years intake		Toddlers aged 2 – 4 years		Children aged 5 – 12 years	
			P95 food intake per day (g)	Grams D-tagatose / day	P95 food intake per day (g)	Grams D-tagatose / day	P95 food intake per day (g)	Grams D-tagatose / day
14.1.3.3	Water based flavoured drinks, artificially sweetened	10 000	1242	12.4	971	*	1103.7	11.0
20.2.2.1	Dairy desserts, artificially sweetened	30 000	70.33	*	NC		NC	
20.2.3	Confectionary (health bars)	100 000	85	8.5	77.4	7.7	62	6.2

* <21 respondents therefore not able to be calculated
NC = not consumed

Table 7: Estimated dietary exposure to individual food groups at the 95th percentile level of consumption – New Zealand

Food code	Description	Level of use (mg/kg)	All population (15+ years)	
			P95 food intake per day (g)	Grams D-tagatose / day
3.1.1.8	Reduced fat ice cream	30 000	240	*
5.2.1	Chewing gum	600 000	30	*
5.3.2	Gum (Soft) type sweets	100 000	144.2	14.4
5.3.3	Hard candy	150 000	125.12	18.8
5.4	Icings/frostings	300 000	24.26	*
6.3	Ready-to-eat breakfast cereals	200 000	75	15.0
13.3	Formulated dietary foods	250 000	237.49	*
13.4	Dietetic formula for slimming	53 000	60	*
13.5	Supplementary foods for dietetic use	53 000	25.58	1.4
13.6	Formulated supplementary sports foods	53 000	NC	
14.1.3.3	Water based flavoured drinks	10 000	1270.8	*
20.2.2.1	Dairy desserts	30 000	667.5	*
20.2.3	Confectionary (health bars)	100 000	102.25	10.2

* <21 respondents therefore not able to be calculated
NC = Not consumed

Where there are less than 21 consumers of a food group, no bolus dose exposures have been calculated from the 95th percentile consumption figure since there are insufficient consumers to derive the figure and for the result to be sufficiently robust. Therefore, the discussion below only includes commodity/ age group combinations for which there were more than 21 consumers.

All estimated short term exposures from single eating occasion, for any population group, for any food, are 24 grams or lower, except for the Australian population for formulated dietary foods (59.3 g) and formulated supplementary sports foods (55.9 g). These products are likely to be infrequently consumed, and would mostly be consumed by adults. The estimates are

also based on the FSANZ estimate of serve size and concentration, which assumes a worst case scenario. Manufacturers may add lower amounts of D-tagatose in practice.

References

Retail World 2002. *Retail World Grocery Guide 2002. 12th Edition*. Retail Media, North Parramatta.

Rutishauser 2000. *Getting it right:- how to use the data from the 1995 National Nutrition Survey*. Commonwealth of Australia.

World Health Organisation (WHO) 2002. *WHO Technical Report Series 909. Evaluation of certain food additives and contaminants. Fifty-seventh report of the Joint FAO/WHO Expert Committee on Food Additives*. World Health Organisation, Geneva.

ENERGY FACTOR FOR D-TAGATOSE

D-TAGATOSE AS A NOVEL FOOD

1 Requirements in the *Australia New Zealand Food Standards Code*

Standard 1.2.8 – Nutrition Information Requirements defines ‘energy factor’ as metabolisable energy and lists factors, expressed as kJ/g, for a large number of energy-yielding components. Energy factors are used in the calculation of a food’s energy content for the purposes of nutrition labelling. Those components that contribute to energy intake or substitute for energy-contributing components are required to have an energy factor listed within Standard 1.2.8. As D-Tagatose is proposed for use as a sugar substitute, and is not an unavailable carbohydrate, the default energy factor is 17 kJ/g. However, the applicant has submitted data that enables derivation and listing of a more accurately estimated factor.

Energy factors in Standard 1.2.8 are derived using the following formula for metabolisable energy:

$$ME = GE - FE - UE - GaE - SE$$

Where ME = metabolisable energy
GE = gross energy
FE = energy lost in faeces
UE = energy lost in urine
GaE = energy lost in gases from large intestine fermentation
SE = energy content of waste products lost from surface areas

In deriving the ME, the information submitted by the applicant has been assessed in accordance with the FSANZ Guidelines “Derivation of energy factors for specific food components not already listed in Standard 1.2.8” (FSANZ Guidelines). These Guidelines can be found at <http://www.foodstandards.gov.au/standardsdevelopment/informationforapplicants/energyfactorsforspec1683.cfm>.

2 Conformity of submitted studies to FSANZ guidelines

Eighteen studies were submitted by the applicant providing evidence for use in the determination of both the percentage of D-tagatose absorbed from the upper gastrointestinal tract (intestinal absorption) and the percentage of ingested D-tagatose excreted into the urine. These studies were assessed against the quality criteria established in FSANZ Guidelines; a detailed description of this assessment is provided in the Appendix to this Attachment.

2.1 Excluded Studies

When assessed against FSANZ Guidelines, six of the eighteen submitted studies were excluded from further consideration due to the use of inappropriate subjects. The results from two studies in rats (Saunders et. al. 1999; Whistler et. al. 1974) could not be realistically applied to humans. Four studies were conducted on humans with an ileostomy, and were thus considered unsuitable (Bär 1990; Jenkins et. al. 1994; Langkilde 1994; Normén et. al. 1999).

2.2 Accepted Studies

Three studies on pigs and nine studies on humans were accepted for assessment against FSANZ Guidelines.

Although the three studies on pigs were deemed relevant for the calculation of energy factors, two did not meet all of the criteria established in FSANZ Guidelines. These studies were not published/peer-reviewed, and did not disclose their source of funding.

Four of the nine human studies did not meet all of the criteria established in FSANZ Guidelines by either:

- absence of control measures within the experimental design;
- inadequate dietary management during the study;
- nondisclosure of the source of funding; or
- nondisclosure of the means of obtaining informed consent from participants.

Notwithstanding these deficiencies, the totality of evidence provides a consistent set of results for each species. The twelve studies have therefore been utilised for the calculation of an energy factor as shown in Table A.

Table A: Application of studies on intestinal absorption and UE

Subject Type for Study	Number of Studies	Used for calculation of intestinal absorption	Used for calculation of UE
Pigs	3	Studies 2 and 3 in Table 1 of the Appendix	Studies 1 and 3 in Table 1 of the Appendix
Humans (healthy)	9	Studies 4 – 10 in Table 1 of the Appendix *	Studies 11 and 12 in Table 1 of the Appendix

* These studies provide only urinary excretion data on L-rhamnose; they were supplied by the applicant for the derivation of a percentage for the intestinal absorption of D-tagatose.

3 Calculating the metabolisable energy of D-tagatose

Each of the components that comprise ME (GE, FE, UE, GaE and SE) requires a separate assessment and calculation. Therefore, an assessment of the evidence for these components has been provided below, with the subsequent derivation of the ME for D-tagatose.

3.1 Gross Energy (GE)

GE or heat of combustion is the total quantity of energy available within a substance. This value is best measured by adiabatic bomb calorimetry, which provides very precise estimates.

A value 15.7 kJ/g can be obtained for the GE of D-tagatose from data supplied by the applicant (Levin et. al. 1995; Livesey 1999). None of this information is based on published bomb calorimetry results, however 15.7 kJ/g is similar to known GE values for glucose and fructose (15.5 kJ/g and 15.2 kJ/g respectively).

A value of **15.7 kJ/g ingested D-tagatose** was assigned to GE.

3.2 Percentage of Gross Energy that is Completely Absorbed in the Upper Intestine

Intestinal absorption is an important consideration, as it determines the energy directly available to the human body from D-tagatose, and the amount of D-tagatose remaining in the large bowel for further fermentation. The percentage of D-tagatose available for fermentation will influence the values obtained for FE and GaE.

Two of the three submitted studies on pigs (Jensen and Laue 1998; Lærke and Jensen 1999) supplied information on intestinal absorption. These studies indicate that approximately 25% of ingested D-tagatose is absorbed from the small intestine, leaving approximately 75% available for further fermentation in the large bowel.

Unlike studies on pigs, human studies cannot ethically undertake invasive portal vein measurements to directly determine the intestinal absorption of D-tagatose. Indirect measurements from urinary excretion values are also unsuitable, as D-tagatose is readily metabolised by the human liver upon absorption from the gastrointestinal tract (Buemann et. al. 1998; Livesey 1999). L-rhamnose, however, is incompletely metabolised in humans and has a similar chemical structure to D-tagatose. Therefore, seven of the nine accepted human studies (Bjarnason et. al. 1991; Bjarnason et al. 1991; Delahunty and Hollander 1986; Howden et al. 1991; Maxton et. al. 1986; Menzies et al. 1990; Mooradian et. al. 1986) were supplied as a means of deriving the intestinal absorption of D-tagatose in humans from the urinary excretion of ingested L-rhamnose.

In using figures on ingested L-rhamnose, consideration has been given to the influence of lipophilicity and molecular size on intestinal absorption (Hamilton et. al. 1987). L-rhamnose will thus have a higher (unknown) rate of intestinal absorption than D-tagatose.

The seven human studies on L-rhamnose indicated that approximately 8-17% of ingested L-rhamnose was excreted into the urine. The applicant has used this information to assign a value of 17-24% to the intestinal absorption of L-rhamnose, and subsequently inferred that no more than 20% of D-tagatose will be absorbed from the gastrointestinal tract in humans (i.e. 80% available for fermentation).

As a number of assumptions are required when using the human studies on L-rhamnose, results from both pigs and humans have been given equally weighting in the determination of intestinal absorption values.

The range of **20-25%** of ingested D-tagatose is to be assigned to intestinal absorption. Consequently, **75-80%** of ingested D-tagatose will be assigned to the percentage of D-tagatose available for fermentation.

3.3 Energy Lost in Faeces (FE)

As specified under FSANZ Guidelines, FE can be calculated by dividing it into three components:

- uFE – the energy lost through excretion of the ingested substance in faeces unchanged,
- mFE – the energy lost in microbial mass through fermentation, and
- oFE – the energy lost through short chain fatty acids that escape large intestinal absorption.

As D-tagatose is completely fermented in humans and appears unchanged in the faeces in minute quantities (Livesey, 1999), uFE can be assigned a value of 0% of fermented D-tagatose.

The applicant has also requested that oFE be set at 0% of fermented D-tagatose, consistent with FSANZ Guidelines.

No data has been made available on mFE; instead the applicant has used the default value of mFE as specified in FSANZ Guidelines. This value is stated as 30% of fermented D-tagatose.

With no energy losses from uFE and oFE, the total value for FE is equivalent to mFE, or **30% of fermented D-tagatose.**

3.4 Percentage of D-tagatose Excreted into Urine

The UE is derived using the percentage of ingested D-tagatose excreted into the urine multiplied by GE.

Using results from studies on pigs (Jensen and Laue 1998; Jørgensen and Lærke 1998), the excretion of 3-7% of ingested D-tagatose into urine has been observed. A slightly lower range was obtained from two human studies (Buemann et. al. 1998; Buemann et. al. 1999), with the observation that approximately 1-5% of ingested D-tagatose is excreted into the urine.

As the results from pigs align closely with those from humans, the human values have been preferentially used as the basis for calculating UE.

The range of **1-5% of ingested D-tagatose** is to be assigned to the percentage of D-tagatose excreted into the urine.

3.5 Energy Lost in Gases from Large Intestine Fermentation (GaE) and Energy Content of Waste Products Lost from Surface Areas (SE)

The applicant has supplied no information on GaE or SE to suggest that the default values provided in FSANZ Guidelines are inappropriate.

GaE and SE will be assigned values of **5% of fermented D-tagatose** and **0 kJ/g of ingested D-tagatose** respectively as specified in FSANZ Guidelines.

3.6 Calculation of the Metabolisable Energy for D-Tagatose

Each of the separate components in the equation for ME are derived as follows:

- GE = 15.7 kJ/g ingested D-tagatose
- FE = % ingested D-tagatose available for fermentation x 0.3 (30%) x GE
- UE = % ingested D-tagatose excreted into the urine x GE
- GaE = % ingested D-tagatose available for fermentation x 0.05 (5%) x GE
- SE = 0 kJ/g ingested D-tagatose

As a range of values can be obtained for both the percentage of D-tagatose available for fermentation (75-80%) and the percentage of ingested D-tagatose excreted into the urine (1-5%), the calculation of ME produces a range of values as listed in Table B.

Table B: Calculation of ME using the range of percentages for UE and availability of D-tagatose for fermentation

Combination of Different Percentages	GE	FE	UE	GaE	SE	ME
Available for Fermentation = 80% Urinary Excretion = 1%	15.7	3.77	0.16	0.63	0	11.15
Available for Fermentation = 75% Urinary Excretion = 1%	15.7	3.5	0.16	0.6	0	11.42
Available for Fermentation = 80% Urinary Excretion = 5%	15.7	3.77	0.8	0.63	0	10.52
Available for Fermentation = 75% Urinary Excretion = 5%	15.7	3.5	0.8	0.6	0	10.79

All values are in kJ/g ingested D-tagatose

Despite the variation in percentages for intestinal absorption and urinary excretion of D-tagatose, the overall impact on the calculation of ME is small. Therefore, the ME of D-tagatose falls within the range of 10.52 – 11.42 kJ/g, with 11 kJ/g being the most appropriate value when rounded to a whole number (consistent with the listing of other energy factors in Standard 1.2.8).

4 Conclusion

The metabolisable energy factor for D-tagatose is best reflected by a value of 11 kJ/g. It is therefore recommended that this energy factor be inserted into Table 2 of subclause 2(2) of Standard 1.2.8.

5 Acknowledgements

FSANZ would like to thank Dr. Penny Warwick for reviewing the attachment on the energy factor for D-tagatose prior to its public release. Dr. Warwick has considerable experience in the area of energy factors, and was instrumental in the development of the guidelines: “Derivation of energy factors for specific food components not already listed in Standard 1.2.8”.

6 References

- Bär A (1990) *Factorial Calculation model for the estimation of the physiological caloric value of polyols*, in Hosoya N (ed.) *Proceedings of the International Symposium on Caloric Evaluation of Carbohydrates*; Japanese Association for Dietetic and Enriched Foods, Tokyo; p209-257.
- Bjarnason I, Fehilly B, Smethurst P, Menzies IS, and Lebi AJ (1991) *Importance of local versus systemic effects of non-steroidal anti-inflammatory drugs in increasing small intestinal permeability in man*, *Gut* 32: 275-277.
- Bjarnason I, Maxton D, Reynolds AP, Catt S, Peters TJ, and Menzies IS (1994) *Comparison of four markers of intestinal permeability in control subjects and patients with Coeliac Disease*, *Scandinavian Journal of Gastroenterology* 29: 630-639.
- Buemann B, Toubro S, and Astrup A (1998) *D-tagatose, a stereoisomer of D-fructose, increases hydrogen production in humans without affecting 24-hour energy expenditure or respiratory exchange ratio*, *Journal of Nutrition* 128: 1481-1486.
- Buemann B, Toubro S, Holst JJ, Rehfeld JF, Bibby BM and Astrup A (1999) *D-tagatose, a stereoisomer of D-fructose, increases blood uric acid concentration*, *Metabolism* 49(8): p969-976.
- Delahunty T and Hollander D (1986) *A comparison of intestinal permeability between humans and three common laboratory animals*, *Comparative Biochemistry and Physiology* 86A(3): 565-567.
- Hamilton I, Rothwell J, Archer D, and Axon ATR (1987) *Permeability of the rat small intestine to carbohydrate probe molecules* *Clinical Science* 73: 189-196.
- Howden CW, Robertson C, Duncan A, Morris AJ, and Russell RI (1991) *Comparison of different measurements of intestinal permeability in inflammatory bowel disease*, *American Journal of Gastroenterology* 86(10): 1445-1449
- Jenkins AP, Menzies IS, Nukajam WS, and Cremer B (1994) *The effect of ingested lactulose on absorption and of L-rhamnose, D-xylose, and 3-O-methyl-D-glucose in subjects with ileostomies*, *Scandinavian Journal of Gastroenterology* 29: 820-825.
- Jensen BB and Laue A (1998) *Absorption of tagatose and fermentation products of tagatose from the gastrointestinal tract of pigs*, Danish Institute of Agricultural Science.
- Jørgensen H and Lærke HN (1998) *The influence of D-tagatose on digestibility and energy metabolism in pigs*, Danish Institute of Agricultural Science.
- Lærke HN and Jensen BB (1999) *D-tagatose has low small intestinal digestibility but high intestinal fermentability in pigs*, *Journal of Nutrition* 129: 1002-1009.
- Levin GV, Zehner LR, Saunders JP, and Beadle JR (1995) *Sugar Substitutes: their energy values, bulk characteristics, and potential health benefits*, *American Journal of Clinical Nutrition* 1995(62) Suppl: 1161S-1168S.
- Livesey G (1999) *D-tagatose: critical review and derivation of its food energy value*, unpublished report.
- Langkilde AM (1994) *Digestion and absorption of sorbitol, maltitol and isomalt from the small bowel. A study in ileostomy subjects*, *European Journal of Clinical Nutrition* 48: 768-775.
- Maxton DG, Bjarnason I, Reynolds AP, Catt SD, Peters TJ, Menzies IS (1986) *Lactulose ⁵¹Cr-labelled ethylenediaminetetra-acetate, L-rhamnose and polyethyleneglycol 500 as probe markers for assessment in vivo of human intestinal permeability*, *Clinical Science* 71: 71-80.
- Menzies IS, Jenkins AP, Heduan SD, Catt SD, Segal MB, Creamer B (1990) *The effect of poorly absorbed solute on intestinal absorption*, *Scandinavian Journal of Gastroenterology* 25: 1257-1264.
- Mooradian AD, Morley JE, Levine AS, Prigge WF, Gebhard RL (1986) *Abnormal intestinal permeability to sugars in diabetes mellitus*, *Diabetologia* 29: 221-224.

Normén L, Lærke HN, Jensen BB, Langkilde AM, and Anderson H (1999) *Small bowel absorption of D-tagatose and related effects on carbohydrate digestibility. An ileostomy study*, American Journal of Clinical Nutrition 73: 105-110.

Saunders JP, Zehner LR, and Levin GV (1999) *Disposition of D-[U-¹⁴C]tagatose on the rat*, Regulatory Toxicology and Pharmacology 29(2): S46-S56.

Whistler RL, Singh PP, Lake WC (1974) *D- Psicose metabolism in the rat*, Carbohydrate Research 34: 200-202.

**D-Tagatose
Comparison of Scientific Literature Against FSANZ Criteria**

The applicant supplied eighteen references in support of the calculation of an energy factor. As specified in the FSANZ Guidelines for the “Derivation of Energy Factors for Specific Food Components Not already listed in Standard 1.2.8” (the FSANZ Guidelines), these studies were assessed against a set of quality criteria.

In the preamble to the quality criteria for submitted studies (Section 3 of the FSANZ Guidelines), animal studies must meet four requirements:

1. Data is provided to show comparability between the results of animal studies and human studies of the same or similar compounds;
2. Care is taken to eliminate coprophagy in rat experiments;
3. Experiments are done at ranges of intakes and in circumstances relevant to realistic intakes in humans; and
4. Clinical (human) studies are completed to confirm any preliminary data obtained by in vitro or animal experiments.

On this basis, two rat studies (Saunders et. al. 1999; Whistler et. al. 1974) were excluded from further consideration, as the results could not be compared and applied realistically to humans. The three pig studies supplied by the applicant (Jørgensen and Lærke 1998; Lærke et. al. 1998; Jensen and Laue 1998) met these requirements.

Within the FSANZ Guidelines, sixteen criteria are listed:

Studies must –

1. have been published in peer-reviewed literature with international circulation;
2. have adhered to ethical guidelines for experimentation in animals or humans (as appropriate), including informed consent in humans, and have reported details of that adherence;
3. report details of funding arrangements for the study;
4. report details of study design, analytical methodology, duration and statistical analysis, and that discuss the limitations of methodology used;
5. report details of how the food component was administered and how ME was calculated (e.g. results from single bolus dose with ME content determined by difference, or from a range of doses and ME determined statistically using regression techniques);
6. include administration of the food component orally with meals/diets of known energy and nutritional content;
7. are conducted under controlled conditions where possible;
8. are conducted under conditions as close as possible to the normal physiological state of the animal or human;
9. in humans, use healthy subjects rather than patients with diagnosed disorders;
10. use adequate (and appropriate) experimental controls;
11. show appropriate statistical considerations in study design and data analysis;
12. use statistically appropriate numbers (and types) of subjects;
13. use appropriate study durations;

14. be minimally invasive;
15. provide appropriately described details; and
16. explore other factors that might affect the estimation of the energy factor of the food component such as adaptation of subjects, fasted or non-fasted conditions, ingestion as liquid or solid or with or without meals, single large dose versus multiple smaller doses, any effects of the test substance on absorption or digestion of other dietary components, and vice versa, and effects of a range of different background diets.

On the basis of criterion 9, four human studies involving ileostomates as subjects were excluded from further consideration (Normén et. al. 1999; Langkilde 1994; Bär 1990; Jenkins et. al 1994). The remaining twelve studies were assessed against the criteria as listed below in Table 1. The headings of each column in Table 1 correspond to the quality criteria as follows:

Peer Reviewed	– Criterion 1
No. of Subjects	– Criterion 12
Ethical Approval / Informed Consent	– Criterion 2
Funding Publicised	– Criterion 3
Study Design	– Criteria 4, 10, 11
Methodology Criteria Met	– Criteria 7, 8, 13, 14, 15
Administration	– Criteria 5 (partially), 6
Calculation of ME Provided	– Criterion 5 (partially)
Met Dietary Criteria Including Adaptation	– Criterion 16

Of the twelve studies assessed, six comply with the sixteen criteria (Bjarnason et al 1991; Buemann et. al. 1999; Howden et al. 1991; Lærke and Jensen 1998; Menzies et al 1990; Mooradian et al 1986). All twelve studies were considered within the calculation of an energy factor for D-tagatose as the results were consistent across each species. However, the relevant discrepancies were noted for each of the six non-compliant studies.

Table 1: Evidence for Intestinal Absorption and Urinary Excretion Values Assessed Against FSANZ Guidelines

Study	Peer Reviewed	Subjects	No. of Subjects	Ethical Approval / Informed Consent	Source of Funding	Study Design	Methodology Criteria Met	Administration	Calculation of ME Provided	Met Dietary Criteria Including Adaptation	Results
1. Jørgensen and Lærke (1998)	No – unpublished report	Pigs	n = 6 (2 per diet; rotated through each diet)	Unspecified	Unspecified	Cross-over, randomised trial, unblinded, controlled.	Yes	Diet 1 – 20% sucrose (control diet) Diet 2 – 10% sucrose + 10% D-tagatose, Diet 3 – 20% D-tagatose Each diet was given over a two-week period.	Yes - ME calculated by the following equation: ME = $GEx0.566$ (absorption from the gut) – UE – GaE.	Yes – 1 out of the 2 weeks as an adaptation period for each diet. Composition of diets was provided.	5.2% ingested D-tagatose in urine, no D-tagatose in faeces.
2. Lærke and Jensen (1998)	Yes	Pigs	n = 16 (6 per diet)	Handling of animals documented in detail.	Funding from MD Food Ingredients AMBA	Parallel group, controlled	Yes	Diet 1 – 15% sucrose (control diet) Diet 2 – 5% sucrose, 10% D-tagatose Each diet was given over fourteen days.	No	Yes – 4 out of 18 days for experimental diet	25.8% ingested D-tagatose absorbed, No D-tagatose in faeces.
3. Jensen and Laue (1998)	No – unpublished report	Pigs	n = 17	Yes – complied with Danish legislation	Unspecified	Cross-over, controlled	Yes	Diet 1 – 20% sucrose (control diet) Diet 2 – 20% tagatose (pigs switched to this diet after control diet).	No	Yes – 1 week adaptation to 20% sucrose, 1 week to 20% D-tagatose. Composition of diets was provided.	26-28% ± 7% ingested D-tagatose absorbed, 3.3-7% ingested D-tagatose into urine.

Study	Peer Reviewed	Subjects	Number of Subjects	Ethical Approval / Informed Consent	Source of Funding	Study Design	Methodology Criteria Met	Administration	Calculation of ME Provided	Met Dietary Criteria Including Adaptation	Results
4. Maxton et. al. (1986)	Yes	Humans	n = 6 (IV) n = 10 (oral)	Yes	Donations of materials for the trial were made by several pharmaceutical companies	Factorial design, uncontrolled	Yes	L-Rhamnose given intravenously and orally. Oral administration provided as isomolar, hyperosmolar or centrimide containing solutions following fasting period. Cr-EDTA and lactulose also in solution. Dose was given once and urine collected at 0, 2.5, 5, 10 and 24 hours.	No	Not Applicable	11.7% and 13.2% ingested L-rhamnose in urine at 5 and 10 hours respectively for the isomolar solution.
5. Howden et al (1991)	Yes	Humans	n = 59 17 control, 28 with Crohn's disease, 14 with ulcerative colitis.	Yes	Donations of materials for the trial were made by several pharmaceutical companies	Factorial design, controlled	Yes	0.7% bolus solution of L-rhamnose given (1g dose) along with mannitol, cellobiose, lactulose, and Cr-EDTA. Urine collected over 6-24 hours.	No	Not Applicable	10.1% of ingested L-rhamnose in urine of control group at 6 hours.
6. Bjarnason et. al. (1991)	Yes	Humans	n = 12	Yes	Funding from Beecham Pharmaceuticals	Randomised controlled trial – double blinded	Yes	Test subjects consumed 100 mL bolus solution containing 1g L-rhamnose (control), and the same solution after 7 days of steroid use. Urine was collected over 5 hours following bolus solution.	No	Not Applicable	17.4 ± 2.5% of ingested L-rhamnose in urine for control dose.

Study	Peer Reviewed	Subjects	Number of Subjects	Ethical Approval / Informed Consent	Source of Funding	Study Design	Methodology Criteria Met	Administration	Calculation of ME Provided	Met Dietary Criteria Including Adaptation	Results
7. Bjarnason et al. (1994)	Yes	Humans	n = 37 12 controls 25 with coeliac disease.	Yes	Donations of materials for the trial were made by several pharmaceutical companies	Factorial design, controlled, double blinded	Yes	5g L-rhamnose dissolved into 150 mL and provided as a bolus. Polyethylene glycol, lactulose, and labelled Cr-EDTA were also provided, bringing the solution to 233 mmol/L. Urine collections taken at 0, 2.5, 5 and 10 hours.	No	Not Applicable	11.3 and 13.7% of ingested L-rhamnose excreted into urine at 5 and 10 hours respectively for controls.
8. Mooradian et al (1986)	Yes	Humans	n = 61 13 controls 48 with diabetes	Yes	Unspecified	Factorial design, controlled, blinding unknown	Yes	Subjects consumed oral solution of 1g L-rhamnose in hyperosmolar solution (1505 mOsm/L) along with 20g lactose, 20g sucrose, and 5g lactulose. Urine was collected over 5 hours	No	Not Applicable	84.3 ± 18.4 mg (8-10%) of ingested L-rhamnose was excreted by controls over 5 hours.
9. Delahunty and Hollander (1986)	Yes	Humans, rats, hamsters and guinea pigs	n = 6 (humans)	Unspecified	Funding from Goldsmith foundation, and CCRC research fund	Parallel group (comparison between species), uncontrolled	Yes	Human subjects were given 5 g L-rhamnose, 10 g lactulose, and 1g mannitol in 200 mL solution as a bolus. Urine was collected over 6 hours.	No	Not Applicable	16.3 ± 2.5% of L-rhamnose into urine of humans.

Study	Peer Reviewed	Subjects	Number of Subjects	Ethical Approval / Informed Consent	Source of Funding	Study Design	Methodology Criteria Met	Administration	Calculation of ME Provided	Met Dietary Criteria Including Adaptation	Results
10. Menzies et al (1990)	Yes	Humans	n = 10 (study 1) n = 8 (study 2)	Yes	Funding from the Special Trustees of St Thomas Hospital	Cross-over trial, controlled	Yes	Subjects were given control solutions of 0.2g glucose, 0.5g xylose and 1.0g rhamnose. Subsequent solutions were given with varying concentrations of either mannitol (study 1) or mannitol and lactulose (study 2). Urine was collected after 5 hours and 48 hours was allowed to elapse between the consumption of each solution.	No	Not Applicable	Study 1: 14.7 ± 1.8% of ingested L-rhamnose into urine (control solutions only). Study 2: 16.7 ± 1.3% of ingested L-rhamnose into urine (control solutions only).
11. Buemann et. al. (1999)	Yes	Humans	n = 8	Yes	Funding from MD Food Ingredients AMBA	Cross-over, randomised, double blinded, controlled	Yes	Subjects fasted for 13 hours. Solutions containing either 30g of D-tagatose or 30g fructose were then provided as a bolus, with a 5-33 day washout period between solutions. Urine was collected at 4 and 7 hours after bolus. Energy expenditure measurements and blood tests were also carried out.	No	Yes – a standardised meal was provided during the seven-hour intervention period.	1.5% of the ingested D-tagatose into urine after 7 hours.

Study	Peer Reviewed	Subjects	Number of Subjects	Ethical Approval / Informed Consent	Source of Funding	Study Design	Methodology Criteria Met	Administration	Calculation of ME Provided	Met Dietary Criteria Including Adaptation	Results
12. Buemann et. al. (1998)	Yes	Humans	n = 6	Yes	Funding from MD Food Ingredients AMBA	Cross-over, randomised, double blinded, controlled	Yes	30g of D-tagatose or Sucrose (control diet) were administered each day via cakes that were consumed as part of the normal diet. Subjects acted as own controls and switched between the two diets after a two-month washout period. The sugars were administered over two weeks. Urine samples were taken during this time at three regular periods during the day. Other tests included hydrogen breath tests and energy expenditure tests.	No	No – dietary restrictions were only enforced days of the energy expenditure tests. Otherwise subjects consumed their normal diet.	0.7 – 5.3% of the ingested D-tagatose appeared in the urine. No D-tagatose appeared in the urine when sucrose was consumed. No significant variation between different urine samples was observed.

SUMMARY OF FIRST ROUND PUBLIC SUBMISSIONS

Food Technology Association, Victoria Inc

- Approve the use of D-tagatose.
- Request that they be maintained on the circulation lists for further changes to this application.

Australian Food And Grocery Council

- Argues that FSANZ needs to determine before accepting the application as a novel food, that it is a non-traditional food and that there is not sufficient knowledge in the broad community to enable its safe use;
- Further, if FSANZ finds that it is safe for human consumption, then it cannot be listed as a novel food.

Department of Health, Western Australia

- disappointed that FSANZ had taken significant direction from the USFDA assessment and approval of D-tagatose, and that
- there was no apparent effort to establish the relevance to the Australian setting and population.
- undecided regarding which option to support because of insufficient information in the Initial Assessment Report