

8.5.2 Primary DNA-damage

Primary DNA-damage in bacteria (Table 40)

H₂O₂ induced DNA-repair in *Escherichia coli* (Thielmann and Gersbach, 1978; De Flora *et al.*, 1984), SOS responses in *Salmonella typhimurium* (Nakamura *et al.*, 1987) and *E. coli* (Von der Hude *et al.*, 1988; Zhou *et al.*, 1991). In the presence of metabolic activation, catalase or SOD, no effect was observed in the DNA-repair test or the SOS chromotest using *E. coli* strains (De Flora *et al.*, 1984; Zhou *et al.*, 1991).

DNA-strand breaks in cultured mammalian cells (Table 41)

H₂O₂ induced DNA single-strand breaks in bovine lens epithelial cells (Kleiman *et al.*, 1990), rat intestinal epithelial cells (Grisham and Kviety, 1989), L1210 leukemia cells (Hincks and Gibson, 1988/89), rat hepatocytes (Olson, 1988), L5178 Y mouse lymphoma cells (Garberg *et al.*, 1988), P388D1 cells (Schraufstatter *et al.*, 1986), rat and human lymphocytes (Schraufstatter *et al.*, 1986; Grisham and Kviety, 1989), human bronchial epithelial cells (Saladino *et al.*, 1985), JB6 Cl 21 cells (Gensler and Bowden, 1983) or V79 cells (Bradley *et al.*, 1979; Bradley and Erickson, 1981; Prise *et al.*, 1989).

DNA double-strand break induction was observed in V79 cells (Prise *et al.*, 1989) and in human leucocytes (Cristovão *et al.*, 1991) but not in cultured rat hepatocytes (Olson, 1988). Decreased DNA and RNA synthesis in human bronchial cells was observed (Saladino *et al.*, 1985) after treatment with H₂O₂, but there was no or a relatively insignificant induction of DNA-protein (Bradley *et al.*, 1979; Bradley and Erickson, 1981; Olson, 1988; Kleiman *et al.*, 1990) or DNA-DNA crosslinks (Bradley *et al.*, 1979; Bradley and Erickson, 1981).

Unscheduled DNA Synthesis (UDS) in cultured mammalian cells (Table 42)

Unscheduled DNA Synthesis was observed in rat hepatocytes (Cattley *et al.*, 1988; Beales and Suter, 1989) and human cells (Stich *et al.*, 1978; Coppinger *et al.*, 1983) after *in vitro* treatment with H₂O₂.

Sister Chromatid Exchanges (SCE) in cultured mammalian cells (Table 43)

H₂O₂ induced Sister Chromatid Exchanges (SCE) in V79 cells (Bradley *et al.*, 1979; Speit *et al.*, 1982; Mehnert *et al.*, 1984a,b; Speit, 1986; Tachon and Giacomoni, 1989), CHO cells (Mehnert *et al.*, 1984a,b; Tucker *et al.*, 1989; Wilmer and Natarajan, 1981), human D98/AH2 cells (Estervig and Wang, 1984) and human lymphocytes (Mehnert *et al.*, 1984a,b). In the presence of S9, catalase or peroxidase, SCE-induction was reduced or inhibited (MacRae and Stich, 1979; Speit *et al.*, 1982; Mehnert *et al.*, 1984a,b).

TABLE 40
PRIMARY DNA-DAMAGE IN BACTERIA

Test system	Metabolic activation	Concentration (μ M)	Results (lowest positive response)	Reference
SOS chromotest				
<i>E. coli</i> PQ37	No Catalase SOD	0-500 20 20	+ve (20 μ M) -ve -ve	Zhou <i>et al.</i> , 1991
SOS chromotest				
<i>E. coli</i> PQ37	No	0-1,000	+ve (0.3mM)	Von der Hude <i>et al.</i> , 1988
Umu test				
<i>Salmonella typhimurium</i> TA1535/pSK1002	No	Unknown	+ve (45mg/ml)	Nakamura <i>et al.</i> , 1987
DNA-repair test				
<i>E. coli</i> WP2, WP67 and CM871	No	Unknown	Minimal inhibitory concentration was 0.02 μ g	De Flora <i>et al.</i> , 1984
DNA excision repair	S9	Unknown	-ve	
<i>E. coli</i>	No	0-400,000	+ve	Thielmann and Gersbach, 1978

TABLE 41

PRIMARY DNA-DAMAGE IN CULTURED MAMMALIAN CELLS - INDUCTION OF DNA-STRAND BREAKS

Test system	Metabolic activation	Treatment time/temperature	Concentration (μM)	Results (lowest effective concentration)	Reference
Human leucocytes	No	10min/ Unknown	500-1,000	Dose-dependent induction of DNA double strand breaks	Cristovao <i>et al.</i> , 1991
Bovine lens epithelial cells	No	5min/37°C	10-200	Linear increase of DNA single-strand breaks (50 μM)	Kleiman <i>et al.</i> , 1990
Rat intestinal epithelial cells	No	15min/37°C	100	Induction of DNA-strand breaks	Grisham and Kviety, 1989
Lymphocytes	No	15min/37°C	100	Induction of DNA-strand breaks	Grisham and Kviety, 1989
V79 cells	No	10-20min/0°C	10-1,000	Dose-dependent induction of DNA single-strand breaks (10 μM)	Prise <i>et al.</i> , 1989
V79 cells	No	20min/0°C	10-10,000	Induction of DNA double-strand breaks at concentration higher than 100 μM	Prise <i>et al.</i> , 1989
L1210 leukemia cells	No	1h/0°C	50-200	Dose-dependent formation of DNA strand breaks (50 μM)	Hincks and Gibson, 1988/89
Rat hepatocytes	No	1h/37°C	10-1,000	Concentration-dependent increases in single-strand DNA breaks (10-50 μM) but lack of induction of double-strand DNA breaks	Olson, 1988
Rat hepatocytes	No	1h/37°C	200/500	Lack of H ₂ O ₂ -induced DNA-protein cross linking	Olson, 1988

TABLE 41 (cntd.)

PRIMARY DNA-DAMAGE IN CULTURED MAMMALIAN CELLS - INDUCTION OF DNA-STRAND BREAKS

Test system	Metabolic activation	Treatment time /temperature	Concentration (μM)	Results (lowest effective concentration)	Reference
L5178Y Mouse Lymphoma cells	No	3h/37°C	2,000-2,510	Concentration-related increase in single-strand DNA breaks (2000 μM)	Garberg <i>et al.</i> , 1988
P388D1 cells and human lymphocytes	No	5min/37°C	25-1,000	Concentration-related increase in single-strand DNA breaks (25 μM)	Schraufstatter <i>et al.</i> , 1986
Normal Human Bronchial Epithelial Cells	No	1h/37°C	100 or 120	Decrease DNA and RNA synthesis Induced DNA single-strand breaks	Saladino <i>et al.</i> , 1985
JB6C121 cells	No	1h/37°C	1-100	Concentration-dependent increase in single-strand breaks (10 μM)	Gensler and Bowden, 1983
V79 cells	No	2h/1°C	7.4-73.5	Concentration-dependent increase in single-strand breaks (14.7 μM)	Bradley and Erickson, 1981
	No	1h/1°C	353	No induction of DNA-protein or DNA-DNA crosslinks	Bradley and Erickson, 1981
V79-4 cells	No	Unknown	353	Induction of DNA single-strand breaks and/or alkali-labile lesions No induction of DNA-protein or DNA-DNA crosslinks	Bradley <i>et al.</i> , 1979

TABLE 42
PRIMARY DNA-DAMAGE IN CULTURED MAMMALIAN CELLS
- INDUCTION OF UNSCHEDULED DNA SYNTHESIS (UDS)

Test system	Protocol	Exposure time (h)	Metabolic activation	Concentration (mM)	Results (lowest effective concentration)	Reference
Rat hepatocytes	Liquid scintillation	20	No	0.1-100	Concentration related increase (1mM)	Beales and Suter, 1989
Rat hepatocytes	Auto-radiography	3x1	No	1.9-6.4	Induction of UDS (3.2mM)	Cattley <i>et al</i> , 1988
WI-38 CCL75 cells (human diploid fetal lung cells)	Auto-radiography	3-4	No	0.6-2400 µg/ml	Concentration-related increase (9µg/ml)	Coppinger <i>et al</i> , 1983
	Differential density labeling	3-4	No	0.15-600 µg/ml	Concentration-related increase (9.4µg/ml)	Coppinger <i>et al</i> , 1983
Human fibroblastes	Auto-radiography	3	No	0.1-10	Concentration-related increase (0.6mM)	Stich <i>et al</i> , 1978

**TABLE 43 - PRIMARY DNA-DAMAGE IN MAMMALIAN CELLS
- INDUCTION OF SISTER CHROMATID EXCHANGES (SCE)**

Test system	Metabolic activation	Exposure time (h)	Concentration (μ M)	Cyto-toxicity (IC ₅₀) (μ M)	Results (lowest effective concentration)	Reference
V79 cells	No	1	5-20	>20 in MEM 5-10 in PBS	No SCE induction in MEM culture medium Concentration related induction of SCE (10 μ M) in PBS medium	Tachon and Giacomoni, 1989
CHO cells	No	15-22	40-240	160	Induction of highly significant levels of SCE and endoreduplicated cells (40 μ M and 160 μ M, respectively)	Tucker <i>et al.</i> , 1989
V79 cells	No	3	10-80	20-40	Induction of SCE (20 μ M)	Speit, 1986
Whole human blood or human purified lymphocytes	No	24	20-2,000	Unknown	SCE induction in purified lymphocytes (20 μ M) but not in whole blood culture	Mehner <i>et al.</i> , 1984a
	No	2	80-200	Unknown	SCE induction in purified lymphocytes (80 μ M) but not in whole blood culture	Mehner <i>et al.</i> , 1984a
Human purified lymphocytes	Catalase or peroxidase or S9-mix	2	80-200	Unknown	Reduction of H ₂ O ₂ -induced SCEs	Mehner <i>et al.</i> , 1984a
D98/AH2 human cells	No	24	15-60	>60	3-fold SCE induction at 60 μ M	Estervig and Wang, 1984
CHO cells	No	2	0.1-100mM	10mM	Slight increase of the SCE frequency (0.5mM)	Wilmer and Natarajan, 1981

TABLE 43 (Cntd)

Test system	Metabolic activation	Exposure time (h)	Concentration (μM)	Cyto-toxicity (IC_{50}) (μM)	Results (lowest effective concentration)	Reference
CHO cells	No	24	0.3-7.8	No toxicity	Two-fold increase of SCE (3.9 μM)	MacRae and Stich, 1979
	Catalase	Un-known	5-100	No toxicity	Complete reduction of the H_2O_2 -induced SCE	MacRae and Stich, 1979
V79 and CHO cells	No	1 or 24	10-40	Un-known	Concentration-related induction of SCE (10-20 μM)	Mehnert <i>et al.</i> , 1984b
	S9-mix	1	10-40	Un-known	Inhibition of SCE-induction	Mehnert <i>et al.</i> , 1984b
V79 cells	No	1 or 9	1-800	Un-known	Induction of SCE	Speit <i>et al.</i> , 1982
	Catalase	9	100-800	Un-known	Complete reduction of the H_2O_2 -induced SCE	Speit <i>et al.</i> , 1982
CHO cells	No	2	100-100,000	>10,000 μM	Slight increase of the SCE frequency (500 μM)	Wilmer and Natarajan, 1981
V79 cells	No	Un-known	353	20% survival	Two-fold increase of the SCE frequency	Bradley <i>et al.</i> , 1979
CHO cells	No	24	0.31-7.8	No toxicity	Two-fold increase of the SCE frequency (3.9 μM)	MacRae and Stich, 1979
	Catalase	Un-known	5-100	No toxicity	Complete reductions of the hydrogen peroxide induced SCE's	MacRae and Stich, 1979

8.5.3 Clastogenicity

Chromosomal aberrations in cultured mammalian cells (Table 44)

Chromosome aberrations (mainly chromatid breaks or micronuclei) were induced in V79 cells (Tsuda, 1981; Tachon and Giacomoni, 1989), Don-6 Chinese hamster cells (Sasaki *et al.*, 1980), human embryonic fibroblasts (Oya *et al.*, 1986), human leucocytes (Cristovão *et al.*, 1991), CHO cells (Stich *et al.*, 1978; Tsuda, 1981; Stich and Dunn, 1986), CHL cells (Ishidate *et al.*, 1984), Syrian hamster and Balb/c mouse cells (Tsuda, 1981), but not in D98/AH2 human cells (Estervig and Wang, 1984) after *in vitro* treatment with H_2O_2 .

In vitro exposure of murine splenocytes to H_2O_2 did not increase the frequency of micronucleated cells (Dreosti *et al.*, 1990). This negative result may be due to the use of the culture medium as the solvent for H_2O_2 and inadequate H_2O_2 concentrations (Oya *et al.*, 1986). Combined treatment with ferrous ions and H_2O_2 , led to a synergistic enhancement of the frequency of micronuclei (Dreosti *et al.*, 1990).

Chromosomal aberrations and micronuclei induction in mammals

Forty-eight hours after implantation of ascite-tumours (S2 sarcoma, Ehrlich ascites carcinoma and sarcoma 180), mice were injected i.p. with 0.2ml H_2O_2 solution at concentrations ranging from 0.01 to 0.5M; 48h later, the treated and untreated tumours were examined. A dose-related increase in the number of chromosomal aberrations of the tumour cells was observed in the H_2O_2 treated mice (Schoeneich, 1967).

H_2O_2 did not induce chromosomal aberrations in bone-marrow cells of rats (Kawachi *et al.*, 1980) and was negative in a micronucleus test in mice (Keck *et al.*, 1980). Details of these two studies are lacking.

8.5.4 Genotoxicity in Cultured Mammalian Cells - Other End-points

H_2O_2 caused morphological transformation of C3H/10T1/2 cells (Nassi-calo *et al.*, 1989).

8.5.5 Mechanistic Considerations

According to several authors, DNA appears to be the main cellular target of H_2O_2 , but its cytotoxicity limits the expression of its genotoxic potential *in vitro* (Sanford *et al.*, 1986; Meneghini, 1988; Cantoni *et al.*, 1989; Ruch *et al.*, 1989).

The formation of adenine N-1-oxide (Mouret *et al.*, 1990), 8-hydroxyguanosine (Kasai *et al.*, 1986), thymidine glycol (Frenkel and Chrzan, 1987), H_2O_2 -induced alkali-labile sites (including

TABLE 44

CHROMOSOMAL ABERRATIONS IN CULTURED MAMMALIAN CELLS

Test system	End-point	Exposure time (h)	Metabolic activation	Concentration (μM)	Results (lowest effective concentration)	Reference
Human leucocytes	Chromosomal aberrations	24	No	15 and 20mM	Six fold increase at 20mM	Cristavao <i>et al.</i> , 1991
Murine splenocytes	Cytokinesis-block micronucleus assay	Unknown	No	10 and 20	No increase of the frequency of micronucleated splenocytes	Dreosti <i>et al.</i> , 1990
V79 cells	Micronuclei	Unknown	Ferrous ions	20	Synergistic enhancement in micronucleus frequency	Dreosti <i>et al.</i> , 1990
		1	No	10-20	Concentration-related increase of micronuclei (10μM) in PBS' medium, no effect in MEM' medium	Tachon and Giacomoni, 1989
Human embryonic fibroblasts	Chromosomal aberrations Chromatid aberrations	10 min	No	10-1,000	Concentration-related increase of chromosomal and chromatid aberrations (20μM)	Oya <i>et al.</i> , 1986
CHO cells	Chromosomal aberrations and micronuclei	3	No	<10-25	Concentration-related increase of chromosome aberrations (12.5μM), chromatid translocations (12.5μM) and micronuclei (12.5μM)	Stich and Dunn, 1986
CHL cells	Chromosomal aberrations	24/48	No	0-7.3	Positive at 24h (3.7μM) and 48h (1.8μM)	Ishidate <i>et al.</i> , 1984

PBS, Phosphate-buffered saline; MEM, Eagles Modified Minimal Essential Medium.

TABLE 44 (cntd)
CHROMOSOMAL ABERRATIONS IN CULTURED MAMMALIAN CELLS

Test system	End-point	Exposure time (h)	Metabolic activation	Concentration (μM)	Results (lowest effective concentration)	Reference
D98/AH ₂ human cells	Chromosomal aberrations	24	No	15-60	Negative	Estervig and Wang, 1984
CHO-K1 cells	Chromosomal aberrations	3	No	100-1,000	Concentration-related increase of chromatid breaks (200 μM)	Tsuda, 1981
V79 cells	Chromosomal aberrations	3	No	100-1,000	Concentration-related increase of chromatid breaks (200 μM)	Tsuda, 1981
Syrian hamster cells	Chromosomal aberrations	3	No	100-1,000	Concentration-related increase of chromatid breaks (200 μM)	Tsuda, 1981
Balb/c mouse cells	Chromosomal aberrations	3	No	10-100	Concentration-related increase of chromatid breaks (50 μM)	Tsuda, 1981
CHO cells	Chromosomal aberrations	2	No	100-100,000	Only positive at 10,000 μM	Wilmer and Natarajan, 1981
CHO cells	Chromosomal aberrations	3	No	100-1,000	Concentration related increase of chromosome aberrations (100 μM), toxic at concentrations higher than 300 μM	Stich <i>et al</i> , 1978
Don-6 cells (Chinese hamster cells)	Chromosomal aberrations	3	No	500-2,000	Dose related increase of chromosome aberrations (1,000 μM)	Sasaki and Sugimura, 1980

DNA strand-breaks), SCEs, gene mutations, chromosomal aberrations, DNA-repair and cell transformations are all mediated by hydroxyl radicals ($\cdot\text{OH}$) aimed at the DNA by DNA-bound transition metals (Emerit *et al*, 1982; Oya *et al*, 1986; Joenje, 1989; Tachon, 1990). These effects can be inhibited by the addition of a strong Fe-chelating agent or a hydroxyl radical ($\cdot\text{OH}$) scavenger (Mello Filho and Meneghini, 1984; Oya *et al*, 1986; Cantoni *et al*, 1989; Nassi-Calo *et al*, 1989) and enzymes that catalytically scavenge intermediates of oxygen reduction (O_2 and H_2O_2) (section 7).

No specific defences are known against $\cdot\text{OH}$ radicals and singlet oxygen which react quickly with almost any biologically active molecule.

The *in vivo* data show discrepancies between positive results observed in host-mediated assays (gene mutations and chromosomal aberrations) and negative results in chromosomal aberration tests in rats and micronucleus tests in mice. Several reports confirm that sensitivity to exogenous H_2O_2 is negatively correlated with cellular catalase activity in mammalian cells (Winqvist *et al*, 1984; Vuillaume, 1987; Sawada *et al*, 1988). The low catalase activity of tumour cells used in the host mediated assay (Schoeneich, 1967) and the direct contact with H_2O_2 by i.p. administration may explain the positive results in the host-mediated assays. On the other hand, the absence of chromosomal abnormalities in the bone-marrow of the orally treated animals may be explained by decomposition of H_2O_2 in the bowel before absorption and by the high catalase activity of red blood cells, which can decompose H_2O_2 after its absorption.

8.5.6 Evaluation

The *in vitro* and *in vivo* genotoxic potential of H_2O_2 is summarised in Table 45.

In conclusion, since only hydroxyl radicals and singlet oxygen are capable of damaging DNA directly, the genotoxic potential of H_2O_2 depends on the accessibility of the extremely reactive hydroxyl radical to its target DNA. *In vitro*, the bacteria or other cells come into direct contact with H_2O_2 and genotoxic effects can be induced; in general, the addition of an exogenous metabolic agent or catalase reduces or abolishes the mutagenic response. *In vivo*, many factors contribute to the reduction of the bioavailability of H_2O_2 for systemic genotoxic action. However, the occurrence of genotoxic effects on cells which are in direct contact with H_2O_2 (at the site of application) cannot be excluded.

TABLE 45
EVALUATION OF GENOTOXICITY

End-Point	Test System	Results
<i>IN VITRO</i>		
Gene mutation	<i>Salmonella typhimurium</i>	+ve without activation decrease of the genotoxic potential in presence of exogenous S9 or catalase
	<i>Escherichia coli</i>	+ve without activation
	<i>Saccharomyces cerevisiae</i>	+/-ve without activation
	<i>Bacillus subtilis</i>	+ve without activation
	Mammalian cells	+ve without activation
Primary DNA-damage		
DNA-repair	<i>Salmonella typhimurium</i>	+ve without activation
	<i>Escherichia coli</i>	+ve without activation -ve with activation
DNA-strand break SCE	Mammalian cells	+ve without activation
	Mammalian cells	+ve without activation
	Mammalian cells	+ve without activation
		decrease of the SCE induction in presence of exogenous S9 or catalase
Chromosomal aberration	Mammalian cells	+ve without activation
Morphological transformation	Mammalian cells	+ve without activation
<i>IN VIVO</i>		
Gene mutation	<i>Drosophila melanogaster</i>	-ve
	<i>Salmonella typhimurium</i> (host mediated assay in mice)	+ve
Chromosomal aberration		
Micronucleus Metaphase analysis	Mice	-ve
	Rat	-ve
	Tumour cells (host mediated assay in mice)	+ve

8.6 REPRODUCTIVE TOXICITY/TERATOGENICITY

8.6.1 Male Fertility

The spermicidal activity of H_2O_2 was investigated both *in vivo* and *in vitro*. Washed and unwashed sperm from bull, rabbit, ram, fowl, dog and mouse were exposed to solutions containing .3, 3, 30, 300 and 3,000ppm H_2O_2 . Sperm were evaluated for mobility *in vitro* following washing and when added directly to diluent. There were large species differences in the tolerance to H_2O_2 . Rabbit spermatozoa were the most resistant and bull and fowl were the least resistant to the immobilising effect of H_2O_2 . Although 3ppm had small, but significant effects on bull and fowl spermatozoa, concentrations 10 to 100 times greater were required to immobilise rabbit spermatozoa. In general, washing increased the toxicity of H_2O_2 , but this effect varied with the species tested. Bull and fowl spermatozoa were least affected by washing and rabbit spermatozoa were most affected (Wales *et al*, 1959). Endogenous catalase is removed when spermatozoa are washed; thus, washing the spermatozoa is expected to increase the toxicity of H_2O_2 .

Male albino mice were given 0.33%, 1.0% or 3.0% H_2O_2 solutions in place of drinking water. There were no controls. The mice at the highest dose (3.0%) would not drink the solution and were taken off the study. Mice were mated after 7 and 21d on H_2O_2 . All females became pregnant within a few days and delivered litters of normal size. The concentration, morphology and mobility of the spermatozoa of the male mice and rabbits receiving H_2O_2 in their drinking water over 3 and 6 weeks remained normal. *In vivo*, H_2O_2 has no significant spermicidal action in mice at concentrations up to 1% in solution (Wales *et al*, 1959).

8.6.2 In Vitro Embryotoxicity

The embryotoxicity of 30% H_2O_2 was investigated using 3-day old (72-76h) Leghorn chicken eggs (Korhonen *et al*, 1984). H_2O_2 solutions (0.05, 0.09, 0.19 and 0.37mg/egg) were dropped on the inner shell membrane in the egg's air chamber, focussing on the embryo visible under the membrane. The ED_{50} for all effects observed was 0.9mg/egg. The LD_{50} was 1.2mg/egg. The NOEL was 0.05mg/egg based on embryo-lethality and toxicity. The significance of this type of study to mammalian species is uncertain.

8.6.3 Reproductive Organs

Male and female rats were administered H_2O_2 by gavage at doses of 1/10-1/5 LD_{50} . Several effects were reported: modification of the oestrus cycle in females and a decrease in sperm mobility in males. Male and female rats received daily doses of 0.005-50mg/kg H_2O_2 by gavage for 6 months. At the high dose, females showed modifications of the oestrus cycle and males reduced mobility of spermatozoa. Treated animals were mated. Among high dose

females, only 3/9 produced litters, compared to 7/9 in the control group. In addition, the body weight of the offspring of the high dose females was reduced relative to those of control females (Antonova, 1974).

8.6.4 Teratology

An attempt was made to assess the teratogenic potential of H_2O_2 in the rat. H_2O_2 was mixed with powdered feed to give 0.02, 0.1, 2 and 10% in the diet which was fed to female rats for 1 week during their 'critical period' of gestation. Unfortunately, this study has multiple deficiencies including a poor design which led to severe reduction of food intake, lack of diet analysis and too few animals for a meaningful statistical analysis. The authors acknowledged that H_2O_2 decomposes readily and showed that the level in the 2% diet declined to background level within 3d. Thus no conclusion regarding the teratogenic potential of H_2O_2 can be made from this study. (Moriyama *et al*, 1982).

A brief communication published over 30 years ago described a study in which 3 rat dams were given 0.45% H_2O_2 to drink. After 5 months, the H_2O_2 was replaced with water and they were mated; all 3 produced normal litters (Hankin, 1958). However, no details regarding study design or conduct were given, thus precluding any evaluation of the adequacy of this study.

8.6.5 Evaluation

Data on the teratogenic potential and reproductive toxicity are too limited to allow an evaluation.

SECTION 9. EFFECTS ON MAN

9.1 ACCIDENTAL EXPOSURE INCIDENCE (TABLE 46)

Reports of accidental exposure resulting in treatment at poison control centres have been tabulated by the American Association of Poison Control Centers since 1985 (Litovitz *et al.*, 1990). Similar data on the European situation are not available. In 1989, there were 7,354 cases of accidental exposure due to consumer use of H_2O_2 as a household topical disinfectant (4,293 cases in 1987). There were 8,557 additional cases due to contact with cosmetic/personal care products containing peroxide (7,587 cases in 1987). The combined number in 1989 (15,911) was greater than the total number of cases reported in 1987 (11,871) (Litovitz *et al.*, 1988).

TABLE 46
ACCIDENTAL EXPOSURES IN THE USA IN 1989
(Litovitz *et al.*, 1990)

Total number of exposures	Age Group (y)			Reason		Treated in Health Care Facility
	<6	6-17	>17	Accident	Intention	
Topical use						
7,354 ^a	4,613	590	2,100	7,199	138	487
Cosmetic and personal care use						
8,557 ^b	5,073	669	2,717	8,291	229	727

a Outcomes of exposure included 2,071 cases with no toxicity, 1,395 cases of minor toxicity and 37 cases of moderate toxicity. There were no deaths.

b Outcomes of exposure included 2,619 cases with no toxicity, 1,917 cases of minor toxicity, 75 cases of moderate toxicity and 3 cases of major toxicity. There were no deaths.

9.2 ACUTE ORAL TOXICITY

A 1-year-old child died of respiratory failure within one hour after ingesting an unknown quantity of a 30% H_2O_2 solution (Giusti, 1973). Post-mortem examination revealed extensive hypostasis and petechiae of the thymus, epicardium and duodenum. The stomach was distended with gas and the viscera were congested. There was some necrosis of the gastro-intestinal mucosa.

A 2-year-old boy ingested an unknown amount of a 3% H_2O_2 solution. He was foaming around the mouth and had vomited three times. A gas embolism was visualised in the portal venous system. It was concluded that liberation of oxygen from H_2O_2 in the stomach ascended through the gastric veins to the portal venous system. The child recovered (Rackoff and Merton, 1990).

A 26-month-old girl ingested a mouthful of a 35% H_2O_2 solution. The child vomited spontaneously. Clinical signs included lethargy and bright red-tinged, frothy emesis. She experienced one fainting episode with short cessation of respirations which returned spontaneously. Endoscopy 16h after exposure showed erosion of the cardiac stomach and erythema of the lower oesophageal sphincter and a gastric burn. Follow-up endoscopy 12d later showed normal esophageal and duodenal mucosa with minimal hyperemia of the cardiac stomach and no ulceration or eschar of the gastric mucosa. The child recovered completely within 2 weeks (Humberston *et al*, 1990).

A 3-year old girl died following ingestion of an unknown quantity of 40% H_2O_2 solution. Pathological findings at autopsy included oedema of the lungs and small erosions of the stomach mucosa, probably due to the lavage tube used to wash out the stomach. The report indicates that the probable cause of death was asphyxia due to obstruction of the distal respiratory tract by the foam liberated by the peroxide (oxygen gas) (Zecevic and Gsparec, 1979).

A 6-year old girl died and her four-year old friend was in critical condition with oropharyngeal burns after accidentally ingesting a solution of 35% H_2O_2 (Thompson, 1989).

Five non-fatal poisonings in adults were reported following consumption of about 25-100ml of H_2O_2 (concentration unknown). The victims experienced sharp pains in the abdomen, foaming at the mouth, vomiting, transitory loss of consciousness, sensory and motor impairment and elevated temperature. Microhaemorrhages of the skin and conjunctiva and moderate leukocytosis were also reported. One person exhibited marked visual (including temporary blindness) and neurological symptoms after swallowing 100ml of the H_2O_2 solution. Symptoms were considered to be due to microemboli of generated oxygen. All victims recovered completely in two to three weeks (Budagovskij *et al*, 1971).

A near-fatality occurred when a 33 year-old woman unintentionally ingested the contents of a 1 pint bottle (<500ml) of 35% H_2O_2 . The patient was cyanotic and experienced seizures and foaming at the mouth. The abdomen was slightly distended. Neurological signs included lack of spontaneous eye movement, no verbal response and withdrawal from noxious stimuli. Laparotomy revealed air bubbles in the stomach but no perforations. Recovery was complete following treatment (Giberson *et al*, 1989).

9.2.1 Evaluation

Young children are particularly vulnerable to accidental ingestion of H_2O_2 . Deaths have been reported from ingestion of unknown quantities of H_2O_2 solutions of 30-40% concentration.

9.3 INTENTIONAL EXPOSURE: MISUSE

There are a number of unofficial reports that some individuals self-administer H_2O_2 under the impression that it can provide health benefits. Such intentional exposure can lead to life-threatening toxicity, including death. A recent publication highlights the dangers of misuse of H_2O_2 . A 39-year old man was admitted to a hospital emergency room in haemolytic crisis 24h after alleged intravenous injection of H_2O_2 (amount and concentration not specified), provided by a local physician, for treatment of cancer. Clinically, the patient was experiencing an acute haemolytic crisis (haematocrit 13.5%). He died following cardiopulmonary arrest on his twelfth day in the hospital (Jordan *et al*, 1991).

9.4 ACUTE EFFECTS FROM IRRIGATING AND CLEANSING SOLUTIONS

9.4.1 Oral Rinses

In a study designed to evaluate the antiseptic activity of a mouth rinse containing 0.5% or 0.75% H_2O_2 in 33% or 50% glycerine, respectively, no irritation of the oral mucosa was noted at the lower concentration in six subjects over a five-day use period with two one-minute rinses daily. Some subjects noted irritation of the mouth and gums at the higher concentration of H_2O_2 , but these effects may have been due to the humectant effect of the 50% glycerine (Slanetz and Brown, 1949).

Rees and Orth (1986) report that the use of 3% H_2O_2 three to five times/day as a mouthrinse resulted in mucosal irritations in two individuals with prior tissue injury. The pre-existing lesions worsened after exposure to 3% solution. Herrin *et al*. (1987) has shown that use of 3% H_2O_2 with sodium bicarbonate did not cause any lesions in healthy individuals. Gingival lesions were seen in patients who used home care solutions employing 5M sodium chloride in addition to 3% H_2O_2 and sodium bicarbonate.

A group of 88 dental students self-administered 6-12.5% H_2O_2 solutions. They used it as a mouthwash and dipped their toothbrushes into the solution before brushing their teeth. Application of the H_2O_2 was 2-3 times/day for 1-2.5 months. Some gingival changes were noted: 6.4% of the subjects showed "redder" gums, 3.4% showed "paler" gums and 6.6% developed hyperkeratinised filiform papillae of the tongue (Miller *et al*, 1938).

The US FDA concluded that concentrations of up to 3% H_2O_2 are safe for over-the-counter use on the mucous membranes of the mouth and throat (US FDA, 1983).

9.4.2 Wound Irrigating and Disinfecting Solutions

A 3% solution of H_2O_2 is widely used as a topical antiseptic agent for suppurative wounds and inflammations of the skin and mucous membranes; as an irritating solution during root-canal therapy and as a mouth rinse for acute necrotising gingivitis (US FDA, 1983).

Cases of rupture of the colon, inflammation of the anus or rectum and ulcerative colitis have been reported following H_2O_2 use of a 1% or 2% solution as an enema (Pumphrey, 1951; Ludington *et al*, 1958; Sheehan and Brynjolfsson, 1960). This is no longer used.

Shock and coma developed suddenly in a 54-year old patient in which 3% H_2O_2 was being used for wound irrigation of the inguinal area during surgery. The patient recovered within one hour. The cause of the near-fatality was considered to be microembolisms formed by H_2O_2 degradation in the closed spaces of the wound irrigation (Bassan *et al*, 1982).

Oxygen embolism has been reported in several infants following intestinal irrigation with H_2O_2 to remove meconium (Danis *et al*, 1967; Shaw *et al*, 1967). In one case, a 36-h old infant died following use of 1% H_2O_2 to remove inspissated meconium from the bowel due to meconium ileus (Shaw *et al*, 1967).

There are several reports of chemically-induced colitis in patients undergoing endoscope examination with instruments which had been cleaned with 3% H_2O_2 (Jonas *et al*, 1988; Bilotta and Wayne, 1989). Discrete white plaques adherent to the colonic mucosa and mild to severe erythema of the surrounding mucosa were observed in one report of twenty patients (Jonas *et al*, 1988). In the other reports, areas of white mucosa extending over areas of the colon were seen following endoscopic procedures (Bilotta and Wayne, 1989). Pathological examination of biopsy specimens revealed nonspecific inflammation. Colitis has also been reported in patients following the use of 3% H_2O_2 diluted with water as an enema (Meyer *et al*, 1981).

Use of H_2O_2 to sanitise haemodialysis equipment has resulted in haemolysis in three pediatric patients, as shown by a significant decline in blood haemoglobin levels (Gordon *et al*, 1990). Apparently, the H_2O_2 sanitising solution was not adequately rinsed out prior to use.

A United States Food and Drug Administration panel indicated that 3% H_2O_2 is safe for over-the-counter use as a topical antiseptic and cleansing solution and sanitising lotions (US FDA, 1983).

9.5 OCULAR EXPOSURE

Several early reports indicate that H_2O_2 was used directly in human eyes to treat corneal ulcers, particularly in herpetic dendritic keratitis (Grant, 1986). In one case, a 20% solution was applied after a local anaesthetic every two hours, as a localised cautery to the ulcer. No apparent corneal damage resulted (Vala, 1965). A 10% solution of H_2O_2 was dropped on one eye of a patient after cocaine with no adverse consequences (Lewin and Guillery, 1913). It is not known how much reliance can be placed on these reports.

Historically, 1-3% solutions have been used as topical ocular antibacterial agents three to five times per day without significant injury (Grant, 1986). Dropping 3% H_2O_2 solution into human eyes causes severe pain, which soon subsides. Even a 0.5% solution caused pain and conjunctival hyperemia (Grant, 1986).

An accident report describes ocular inflammation, hyperaemia, tears and eyelid spasm in a woman who inadvertently placed in her eye a contact lens which had been stored in 3% H_2O_2 . Vision was reduced insignificantly and recovery was complete in a few days following treatment with anti-inflammatory drops (Knopf, 1984).

Contact lens solutions of H_2O_2 generally contain buffers and preservatives which may reduce the ocular toxicity. The threshold irritant concentration was determined by introducing hydrogel contact lens treated with various concentrations of H_2O_2 into human eyes. Concentrations of 0, 25, 50, 100, 200, 400 and 800ppm were tested; none affected the physiological integrity of the eye. An initial conjunctival hyperemia occurred with the 800 and 400ppm levels but not with 200ppm and below. Levels of H_2O_2 in excess of 100ppm were associated with an initial subjective stinging response. Subjective effects, such as stinging were first noticed between 50 and 100ppm; the NOEL was 50ppm. There were no disruptive changes to the corneal or conjunctival epithelium at any concentration (Paugh *et al*, 1988).

The mean ocular threshold for discomfort in human subjects was 247ppm when solutions were instilled directly into the eyes (McNally, 1990).

H_2O_2 has gained wide-spread use as a disinfectant of contact lenses, resulting in maximum residual concentrations of 50-60ppm after neutralisation; this is close to the natural background level of H_2O_2 in the eye lens (section 7).

9.6 DERMAL EXPOSURE

A characteristic temporary whitening of the skin occurs after dermal application of 20% to 30% H_2O_2 . The whitening is due to oxygen bubbles acting as microemboli in the capillaries and blocking circulation (Hauschild *et al.*, 1958). H_2O_2 is used topically to bleach hair at concentrations of 3% to 6%. No clear data are available on the skin irritation effects of various H_2O_2 solutions in human beings (see also next section).

9.7 VAPOUR EXPOSURE

There are few reports of adverse effects resulting from inhalation exposure to H_2O_2 , although bleaching of the hair is a common occurrence. Bleaching has been known to occur with levels of 0.5-1ppm vapour (FMC, 1990e), but it is uncertain whether bleaching was due to vapour exposure or to transfer of H_2O_2 from the hands to hair.

Eye and throat irritation and gradual bleaching of hair have been reported among factory workers exposed to H_2O_2 aerosol concentrations of 12 to 41mg/m³ (8.6 to 29.5ppm) (Kaelin *et al.*, 1988). The workers were operators of a semi-automatic milk packing machine which used a H_2O_2 bath to sanitise cardboard packaging for milk. One worker developed interstitial lung disease and impaired gas exchange, but since he was a heavy smoker, the cause of the pulmonary disease could not be ascertained.

In a USSR study, the irritant threshold to the lungs was determined in human volunteers exposed from 5 minutes to 1-4h. The threshold level was considered to be 10mg/m³ for the full period (up to 4h) with a NOEL of 5mg/m³ (Kondrashov, 1977). Increasing the time over which skin was exposed to H_2O_2 aerosols from 5min to 4h decreased the irritation threshold from 180 to 20mg/m³ (Kondrashov, 1977).

9.8 INDIVIDUAL SUSCEPTIBILITY

Some individuals are more susceptible to H_2O_2 exposure because of hereditary disorders in H_2O_2 metabolising enzymes (section 7).

9.9 SKIN SENSITISATION

No data are available.

9.10 CHRONIC TOXICITY AND CARCINOGENICITY

No data are available.

9.11 REPRODUCTIVE TOXICITY

Human sperm was exposed to solutions containing 0.3, 3, 30, 300 and 3,000ppm H_2O_2 *in vitro*. In unwashed spermatozoa, the ED_{50} (the concentration that results in a 50% reduction of the mobility index of spermatozoa) of H_2O_2 was between 30 and 300ppm (Wales *et al*, 1959). Removal of endogenous catalase by washing increased the toxicity of H_2O_2 to spermatozoa.

9.12 NEUROTOXICITY

No information is available.

9.13 EVALUATION

Accidental exposure to H_2O_2 has occurred from its use in household topical disinfectants and cosmetic/personal care products. Young children are particularly vulnerable to accidental oral exposure. Deaths have been reported from ingestion of unknown quantities of 30-40% solutions. The effects seen were related to a corrosive action on the gastro-intestinal tract and to generation of large volumes of oxygen. Complete recovery has occurred in near fatal cases within 2-3 weeks with medical care.

H_2O_2 solutions of 1-3% have been used as anti-bacterial agent in the eye and mouth washes without causing significant injury. In the eye, H_2O_2 solutions of >200ppm caused hyperemia; pain and stinging occurred with concentrations of ≥ 100 ppm.

In volunteers exposed to aerosol for 4h, the irritation threshold for the respiratory tract was $10\text{mg}/\text{m}^3$ H_2O_2 and for the skin $20\text{mg}/\text{m}^3$. At these concentrations, eye and throat irritation and gradual bleaching of hair has been reported. At concentrations at or below the present occupational exposure limit of $1.4\text{mg}/\text{m}^3$ only hair bleaching was observed; this may, however, also have been caused by H_2O_2 transferred from the hands to the hair.

SECTION 10. FIRST AID AND SAFE HANDLING ADVICE

This section has been based on Material Safety Datasheets from L'Air Liquide (1990b-j), Amylum (1990), Atochem (1989a-e, 1990), Degussa (1986, 1988, 1989), Du Pont (1990a,b), Eka Nobel (1986a-c, 1988a,b, 1989), FMC (1990a-d), Mallinckrodt (1989), Montefluos (1990a-c) and Solvay (1984a,b), and on information from CEDRE (1990) US DOT (1987) and Weiberg and Leuchtenberger (1978).

10.1 FIRST AID AND MEDICAL TREATMENT

This advice applies to all commercially available grades of H_2O_2 solutions (3-70%), unless stated otherwise.

10.1.1 First Aid

Eye contact. Immediately flush eyes with plenty of water for at least 15 minutes (keep eyelids apart). Seek medical attention; with $\leq 5\%$ solutions, medical attention is required only if irritation or burning sensation persists.

Skin contact. Remove contaminated clothing and shoes. Soak contaminated clothes in water to prevent risk of fire. Wash contaminated skin with water for at least 15 minutes. Keep patient warm and quiet. Seek medical attention. With a $\leq 5\%$ solution, medical attention is required only if irritation or ulcers develop.

Inhalation. Move subject to fresh air and keep warm and quiet. If not breathing, give artificial respiration. Obtain medical attention immediately.

Ingestion. Do not induce vomiting. Flush mouth with water. Give several glasses of water to drink if subject is conscious. Keep subject quiet and warm. Seek medical attention.

10.1.2 Notes to Physicians

H_2O_2 solutions and their vapours and airborne mists are of low toxicity, but, depending on the concentration and duration of exposure, may cause mild ($\leq 5\%$ solutions) to severe damage to the skin and mucous membranes. Symptoms will subside after exposure ceases except for highly concentrated solutions ($\geq 70\%$) when symptoms are likely to persist. It is often difficult to estimate the amount of H_2O_2 ingested or the concentration of H_2O_2 solutions.

Eye contact. Stinging and burning sensation of the eyelids and conjunctivitis occur with 5% solutions and with vapours. Direct contact with 8% solutions may cause burning of the eyelids, keratitis; necrosis of mucous membranes, ulceration of the cornea. Appearance of

damage, even corneal ulceration, may be delayed for hours or days after an initial (false) healing period.

Following first aid, instil 1-2 drops of an anaesthetic or apply local corticosteroids. Keep subject under surveillance for 1-2 weeks.

Skin contact. Contact on a burn or open skin may cause a stinging or burning pain. A short period (2-3h) of whitening may occur at the point of contact, followed by redness, blisters, deep cutaneous necrosis. If large areas of skin are affected, the subject may suffer from shock. Treat as for thermal burns. No adverse effects are expected from dilute ($\leq 5\%$) solutions on intact skin.

Inhalation. Effects depend on vapour or aerosol concentration and may include irritation of the nose, eyes and throat. Coughing, dyspnoea, shortness of breath may occur and, in the worst cases, there is a risk of lung oedema.

Ingestion. Ingestion may produce a burning sensation in the throat, oesophagus and stomach, abdominal pain, vomiting and diarrhoea. High concentrations may produce buccal foaming, micro-haemorrhage of the gastric mucosa with risk of micro-embolism. Blood-stained vomit, glottis oedema and shock may occur in severe cases.

Do not induce vomiting because of the danger of aspiration of foam. Prevent and treat shock, treat digestive tract burns. Large amounts of oxygen may rapidly be released from H_2O_2 . Severe distension may be treated by flexible nasogastric or orogastric tube. Evacuation of the stomach by emesis induction or gastric lavage should be avoided.

10.2 SAFE HANDLING

General precautions.

Protect against heat.

Work in strictly clean areas and use only the minimal quantity necessary in the workplace.

All installations, containers and equipment must be made of completely pure materials compatible with H_2O_2 (section 3.2) and provided with the necessary devices to avoid pressure build-up, such as pressure relief valves and/or bursting discs, these should release to a safe area.

The design and operation of the system should ensure that no contamination can enter it. If a breather vent or similar device is used it should be provided with a filter to avoid any airborne contaminants entering the H_2O_2 system.

Depending upon the quantity of product being stored/handled, additional venting may be required by means of a ventable lid.

All materials must be cleaned before coming into contact with H_2O_2 according to the suppliers recommendations. Vessels and other ancillary items used for H_2O_2 handling should be exclusively reserved for this purpose. Never return unused solution to containers of H_2O_2 .

When emptying plastic containers the preferred methods are syphoning or pumping, pressure should not be used. For stronger containers, pressure may be used, but only good quality air (or nitrogen), free from oil and other airborne contaminants, should be used.

Never confine H_2O_2 between two valves.

Personnel protection. Personnel must be educated in the hazards of H_2O_2 and that it slowly decomposes. Contamination causes the decomposition to take place at a much enhanced rate, and can be initiated even with trace amounts (a few ppm) of transition metal ions, and pH changes. Other factors such as direct sunlight and high temperatures can also lead to decomposition.

Decomposition results in a rapid increase in the rate of gas (plus steam) production which can lead to a pressure build up and bursting of the container if adequate relief is not provided.

H_2O_2 at strength >8% is an oxidising substance, having the potential to ignite combustible materials. Evaporation of water, e.g. by heat, may increase the concentration of a H_2O_2 solution, the higher the strength of the solution the greater is the potential for ignition.

When handling solutions of <8%, normal work clothing can be worn with impervious gloves, apron and shoes as well as eye protection. For stronger solutions, wear a non-flammable worksuit made of natural rubber, neoprene, nitrile rubber or PVC, including gloves, boots, trousers, jacket, and hood.

Leather shoes should not be worn as these can ignite within a few minutes of contact with H_2O_2 . Cotton clothing can also ignite quickly. Protective skin creams offer no protection from H_2O_2 .

Eye protection using chemical safety goggles must be worn when handling H_2O_2 . In addition, a face-shield is desirable to protect against accidental splashing.

Safety showers and eye fountains must be provided at the work place, in addition sufficient clean water should be provided to deal with spillages.

Respiratory protection should be provided by natural or general ventilation of the workplace, with local exhaust points, where necessary, to control mists or vapours. Where there is a potential for airborne exposure in excess of applicable limits (section 5.2), a face-mask with an approved cartridge must be provided, used as an escape mask. In cases of severe exposure, e.g. in dealing with an emergency, self-contained breathing apparatus should be used by qualified personnel.

Storage. Store H_2O_2 solutions outside in a cool, clean place or in clean, well ventilated rooms, built with non-combustible materials. Do not stack containers next to the air vents. The floors should be impermeable so that when accidental leakage occurs, the product will flow away to a safe area or be retained.

Do not store on wooden pallets. Store away from reactive substances (e.g. finely divided metals and their salts, strong acids and alkalis, organic substances, oils and greases) and combustible materials (e.g. straw, paper, wood, textile), and heat sources.

Containers should be regularly checked for signs of abnormality, e.g. bulging or increased temperature. Regularly verify the availability of ample supply of water for emergencies. For bulk storage, the supplier should be consulted regarding the design and installation.

10.3 MANAGEMENT OF SPILLAGE AND WASTE

Spillage. In the event of a container spillage, evacuate and isolate the hazard area and keep people away, stay upwind and keep out of low-lying areas. Contain the spill, if possible, and dilute with large amounts of water before disposal. If containment is not possible wash down with copious amounts of water. If water pollution occurs, notify the appropriate authorities.

If containers are involved in a fire situation, drench with water, keep adjacent containers cool by spraying with water. These operations should only be undertaken by personnel equipped with fire fighting protection. For bulk storage spillages, a pre-arranged plan must be worked out in conjunction with the supplier and the competent authority, if applicable.

Waste disposal. In most countries, there will be statutory regulations governing the disposal of H_2O_2 . Solutions containing <1% can normally be flushed down drains (no explosion hazard); in certain cases it may need to be more dilute.

For further and more detailed safety instructions, contact your H_2O_2 supplier.

BIBLIOGRAPHY

- Abu-Shakra, A. and Zeiger, E. (1990). Effects of *Salmonella* genotypes and testing protocols on hydrogen peroxide-induced mutation. *Mutagenesis* 5, 469-473.
- ACGIH (American Conference of Governmental Industrial Hygienists) (1986). Documentation of the threshold limit values and biological exposure indices (5th ed). ACGIH, Cincinnati, OH, 316.
- ACGIH (American Conference of Governmental Industrial Hygienists) (1991). Threshold limit values and biological exposure indices for 1990-1991. ACGIH, Cincinnati, OH, 24.
- Aebi, H. and Suter, H. (1972). Acatalasemia. In: Stanbury, J. B., Wyngaarden, J. B. and Fredrickson, D. S. (eds), *The metabolic basis of inherited diseases*. McGraw-Hill, New York, 1710-1750.
- AFS (Arbetarskyddsstyrelsens Foerfattningssamling) (1990). Hygieniska graensvaerden, AFS 1990:13. AFS, Stockholm, 40.
- Aguinaldo, E. R., Weiner, M. L., Freeman, C., McCarty, J. D., Fletcher, M. J. and McConnell, F. (1992). Skin irritation studies with hydrogen peroxide, abstract 354. *The Toxicologist* 12, 110.
- Air Liquide, L'. (1990a). Catalogue produits, fiche de caractéristiques n° 56, peroxyde d'hydrogène type 30% électronique. L'Air Liquide, dép. chimique, Paris, 1.
- Air Liquide, L'. (1990b). Peroxyde d'hydrogène à 10%, fiche de données de sécurité 540. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990c). Peroxyde d'hydrogène à 18%, fiches de données de sécurité 535. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990d). Peroxyde d'hydrogène à 30%, fiche de données de sécurité 530. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990e). Peroxyde d'hydrogène à 35%, fiches de données de sécurité 525. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990f). Peroxyde d'hydrogène à 43%, fiche de données de sécurité 520. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990g). Peroxyde d'hydrogène à 50%, fiches de données de sécurité 515. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990h). Peroxyde d'hydrogène à 60%, fiche de données de sécurité 510. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990i). Peroxyde d'hydrogène à 70%, fiches de données de sécurité 505. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1990j). Peroxyde d'hydrogène à 85%, fiche de données de sécurité 501. L'Air Liquide, Paris, 1-4.
- Air Liquide, L'. (1991). Evolution de la concentration en peroxyde d'hydrogène dans des échantillons tirés de la Saône. Unpublished data.
- Amylum (1990). Waterstofperoxide - 27,5% H₂O₂. Amycor, Aalst, 1-2.
- Andrae, U., Singh, J. and Ziegler-Skylakakis, K. (1985). Pyruvate and related alpha-ketoacids protect mammalian cells in culture against hydrogen peroxide-induced cytotoxicity. *Toxicol. Let.* 28, 93-98.
- Antonova, A. (1974). Argumentation hygiénique de la concentration limitée admissible en peroxyde d'hydrogène dans les réservoirs d'eau. *O. Gigiena i sanit.* 10, 20-22 [Russian].
- Aoki, M. and Tani, Y. (1972). Growth and histopathological changes in mice fed with hydrogen peroxide solution in stead of water. *Icaku to Seibutsugaku (Medicine and Biology)* 84, 3, 159-162 [Japanese].
- Arbeidsinspectie (1989). De nationale MAC-lijst 1989. Directoraat-Generaal van de Arbeid van het Ministerie van Sociale Zaken en Werkgelegenheid, Voorburg, 24-25.
- Arbeidstilsynet, Direktoratet for (1989). Administrative normer for forurensning i arbeidsatmosfaere. Veiledning til arbeidsmiljøloven. Arbeidstilsynet, Oslo, 12.
- Arbejdstilsynet (1988). Graensevaerdier for stoffer og materialer. At-anvisning Nr. 3. 1. 0. 2, April 1988. Arbejdstilsynet, København, 21.
- Atochem, (1989a). Fiche de données de sécurité. Peroxyde d'Hydrogene 30% Medicinal. Atochem, Paris, 1-4.
- Atochem, (1989b). Fiche de données de sécurité. Peroxyde d'Hydrogene 35% Industriel. Atochem, Paris, 1-4.

- Atochem. (1989c). Fiche de données de sécurité. Peroxyde d'Hydrogene 35% Parfumerie. Atochem. Paris. 1-4.
- Atochem. (1989d). Fiche de données de sécurité. Peroxyde d'Hydrogene 50% Industriel. Atochem. Paris. 1-4.
- Atochem. (1989e). Fiche de données de sécurité. Peroxyde d'Hydrogene 70%. Atochem. Paris. 1-4.
- Atochem. (1990). Fiche de données de sécurité. Peroxyde d'Hydrogene 35% A1. Atochem. Paris. 1-4.
- Aust, S. D. and White, B. C. (1985). Iron chelation prevents tissue injury following ischemia. *Adv. Free Radical Biol. Med.* 1, 1-17.
- Baehner, R. L., Boxer, L. A. and Ingraham, L. M. (1982). Reduced oxygen by-products and white blood cells. Pryor, W. (ed.), *Free Radicals in Biology* 5. Academic Press, New York, 91-113.
- Báldry, M. G. C. (1983). The bactericidal, fungicidal and sporicidal properties of hydrogen peroxide and peracetic acid. *J. Appl. Bacteriol.* 54, 417-423.
- Balvay, G. (1981). Biological consequences of treating a lake with hydrogen peroxide on the plankton biocenosis. *Water Res.* 15, 691-696.
- Barenschee, E. R. (1990). Halbwertszeit H₂O₂ in Boeden, Wasser, Abwasser. Personal communication 29. 10. 90. Degussa. Hanau, 1-3
- Barth, M. C., Hegg, D. A., Hobbs, P. V. and Walega, J. G., Kok, G. L., Heikes, B. G., Lazrus, A. L. (1989). Measurements of atmospheric gas-phase and aqueous-phase hydrogen peroxide concentrations in winter on the east coast of the United States. *Tellus* 41B, 61-69.
- Bassan, M. M., Dudai, M. and Shalev, O. (1982). Near-fatal systemic oxygen embolism due to wound irritation with hydrogen peroxide. *Postgrad. Med. J.* 58, 448-450.
- Beales, S. and Suter, W. (1989). DNA repair induction by UV irradiation and various mutagens requiring metabolic activation in rat hepatocytes maintained in culture for several days. *Mutagenesis* 4, 456-461.
- Becewa (1984). Wasserstoffperoxide. *Tijdschr. Becewa* 77, 111-118.
- Bhuyan, K. C. and Bhuyan, D. K. (1977). Regulation of hydrogen peroxide in eye humors; effect of 3-amino-1H-1,2,4-triazole on catalase and glutathione peroxidase of rabbit eye. *Biochem. Biophys. Acta* 497, 641-651.
- Bilotta, J. J. and Wayne, J. D. (1989). Hydrogen peroxide enteritis: the "snow white" sign. *Gastro-intestinal Endoscopy* 35, 428.
- Birnboim, H. C. (1982). Factors which affect DNA strand breakage in human leukocytes exposed to a tumor promoter, phorbol myristate acetate. *Can. J. Physiol. Pharmacol.* 60, 1359-1366.
- Boatman, J. F., Wellman, D. L., Van Valin, C. C., Gunter, R. L., Ray, J. D., Sievering, H., Kim, Y., Wilkison, S. W. and Luria, M. (1989). Airborne sampling of selected trace chemicals above the central United States. *J. Geophys. Res.* 94, D4, 5081-5093.
- Bock, F. G., Myers, H. K. and Fox, H. W. (1975). Cocarcinogenic activity of peroxy compounds. *J. Nat. Cancer Inst.* 55, 1359-1361.
- Bors, W., Saran, M., Lengfelder, E., Spoettl, R. and Michel, C. (1974). The relevance of the superoxide anion radical in biological systems. *Curr. Top. Radiat. Res. Quart.* 9, 247-309.
- Bosworth, D., Crofton-Sleigh, C. and Venitt, S. (1987). A forward mutation assay using ampicillin-resistance in *Escherichia coli* designed for investigating the mutagenicity of biological samples. *Mutagenesis* 2, 455-467.
- Boveris, A., Oshino, N. and Chance, B. (1972). The cellular production of hydrogen peroxide. *Biochem. J.* 128, 617-630.
- Boveris, A. and Chance, B. (1973). The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem. J.* 134, 707-716.
- Boveris, A. (1977). Mitochondrial production of superoxide radical and hydrogen peroxide. In: Reivich, M., Coburn, R., Lahiri, S. and Chance, B. (eds), *Tissue hyperoxia and ischemia*. Plenum, New York, 67-82.
- Bradley, M. O., Hsu, I. C. and Harris, C. C. (1979). Relationships between sister chromatid exchange and mutagenicity, toxicity and DNA damage. *Nature* 282, 318-320.

- Bradley, M. O. and Erickson L. C. (1981). A comparison of the effects of hydrogen peroxide and X-ray irradiation on toxicity, mutation, and DNA damage/repair in mammalian cells (V, 79). *Biochimica et Biophysica Acta* 654, 135-141.
- Bringmann, G. and Kuehn, R. (1982). Ergebnisse der Schädlichkeitsbewertung von wasser- und luftgefährdenden Stoffen gegen *Daphnia magna* in einem weiterentwickelten standardisierten Testverfahren. *Z. Wasser-Abwasser-Forsch.* 15, 1, 1-6.
- Bryan, C. L. and Jenkinson, S. G. (1981). Species variation in lung antioxidant enzyme activities. *J. Appl. Physiol.* 63, 597-602.
- Budagovskija, M. T., Vadachkoniya, M. K. and Danyalov, A. I. (1971). Poisoning by perhydrol. *Voen. Med. Zh.* 9, 79-81 [Russian].
- Bufalini, J. J., Gay, B. W. and Brubaker, K. L. (1972). Hydrogen peroxide formation from formaldehyde photooxidation and its presence in urban atmospheres. *Environ. Sci. Technol.* 6, 818-821.
- Bundesminister fuer Jugend, Familie, Frauen und Gesundheit (1990). Verordnung zur Aenderung der Trinkwasserverordnung und der Mineral- und Tafelwasser-Verordnung. *Bundesgesetzblatt I*, 2600, 2604.
- Calvert, J. G. and Stockwell, W. R. (1983). Acid generation in the troposphere by gas-phase chemistry. *Envir. Sci. Technol.* 17, 428A-443A.
- Cantoni, O., Brandi, G., Cerutti, P., Meyn, E., Murray D. (1989). Mechanisms of hydrogen peroxide cytotoxicity in mammalian and bacterial cells. In: *Proc. Int. Conf. DNA Damage Repair*, 281-289.
- Cardinaels, J. (1990). Valeurs limites tolérables pour substances dangereuses, T. L. V. 1989-1990. Ced. Samson, Brussels, 38.
- Carlsson, J., Berglin, E. H., Claesson, R., Edlund, M. B. K. and Persson, S. (1988). Catalase inhibition by sulfide and hydrogen peroxide-induced mutagenicity in *salmonella typhimurium* strain TA 102. *Mutation res.*, 202, 59-64.
- Cattley, R. C., Smith-Oliver, T., Butterworth, B. E. and Popp, J. A. (1986). Failure of the peroxisome proliferator WY-14,643 to induce unscheduled DNA synthesis in rat hepatocytes following *in vivo* treatment. *Carcinogenesis* 9, 7, 1179-1183.
- Cavarocchi, N. C., England, M. D., Schaff, H. V., Russo, P., Orszulak, T. A., Schnell, W. A., O'Brian, J. F. and Pluth, J. R. (1986). Oxygen free radical generation during cardiopulmonary bypass: correlation with complement activation. *Circulation* 74, suppl. III, 130-133.
- CEDRE (Centre de Documentation de Recherche et d'Experimentations sur les Pollutions Accidentelles des Eaux) (1990). Mini-guide d'intervention et de lutte face au risque chimique, peroxyde d'hydrogene, eau oxygénée, bioxyde d'hydrogene. CEDRE, Plouzané, F. 1-2.
- Chalmers, R. L. (1989). Hydrogen peroxide in anterior segment physiology: a literature review. *Optometry and Vision Science* 66, 11, 796-803.
- Chance, B., Sies, H. and Boveris, A. (1979). Hydroperoxide metabolism in mammalian organs. *Physiol. Rev.* 59, 527-605.
- Chance, K. V. and Traub, W. A. (1987). Evidence for stratospheric hydrogen peroxide. *J. Geophys. Res.* 92, D3, 3061-3066.
- ✓ Chemoxal (1992). Chemoxal data on industrial sources of hydrogen peroxide; Annual input of hydrogen peroxide in surface waters through rain in France. Personal communications by Dr. A. Balay, 07/01/1992 and 29/04/1992. Chemoxal, Chalon-sur-Saone, 1-3, 1-2.
- Chiu, D., Lubin, B. and Shohet, S. B. (1982). Peroxidative reactions in red cell biology. In: Pryor, W. A. (ed.), *Free Radicals in Biology* 5. Academic Press, New York, 115-160.
- Chow, C. K. and Tappel, A. L. (1972). An enzymatic protective mechanism against lipid peroxidation damage to lungs of ozone-exposed rats. *Lipids* 7, 518-524.
- Christensen, G. M. (1971/72). Effects of metal cations and other chemicals upon the *in vitro* activity of two enzymes in the blood plasma of the white sucker, *Catostomus commersoni* (Lacépède). *Chem. - Biol. Interactions* 4, 351-361.
- CNA (Caisse Nationale Suisse d'Assurance en Cas d'Accidents) (1987). Valeurs limites tolérables aux postes de travail: substances toxiques (valeurs MAC), agents physiques. CNA, 57.
- COLIPA (Comité de Liaison des Associations Européennes de l'Industrie de la Parfumerie, des Produits Cosmétiques et de Toilette) (1991). Hydrogen peroxide used in cosmetics. Personal communication by J. Dupuis, COLIPA, Brussels, 1.

- Comstock, C. C., Hackley, E. B., Oberst, F. W., Lawson, L. and Greene, E. (1954). The inhalation toxicity of 90% hydrogen peroxide vapour for acute, subacute and chronic exposures to laboratory animals. Chemical Corps Medical Laboratories Res. Rep. 243. Army Chemical Center, Edgewood, MA. 1-17.
- Cooper, W. J. and Lean, D. R. S. (1989). Hydrogen peroxide concentration in a northern lake: photochemical formation and diel variability. *Environ. Sci. Technol.* 23, 11, 1425-1428.
- Cooper, W. J., Lean, D. R. S. and Carey, J. H. (1989). Spatial and temporal patterns of hydrogen peroxide in lake waters. *Can. J. Fish. Aquat. Sci.* 46, 1227-1231.
- Cooper, W. J., Saltzman, E. S. and Zika, R. G. (1987). The contribution of rainwater to variability in surface ocean hydrogen peroxide. *J. Geophys. Res.* 92, C3, 2970-2980.
- Cooper, W. J. and Zika, R. G. (1983). Photochemical formation of hydrogen peroxide in surface and ground waters exposed to sunlight. *Science* 220, 711-712.
- Cooper, W. J., Zika, R. G., Petasne, R. G. and Plane, J. M. C. (1988). Photochemical formation of hydrogen peroxide in natural waters exposed to sunlight. *Environ. Sci. Technol.* 22, 1156-1160.
- Coppinger, W. J., Wong, T. K. and Thompson, E. D. (1983). Unscheduled DNA Synthesis and DNA repair studies of peroxyacetic and monoperoxydecanoic acids. *Environmental Mutagenesis* 5, 177-192.
- Cristov  o, L., Teixeira, M., Br  s, A. and Rueff, J. (1991). DNA and chromosomal damage by hydrogen peroxide in human cells. EEMS meeting, Prague, September 1991, 1-2.
- Damschen, D. E. and Martin, L. R. (1983). Aqueous aerosol oxidation of nitrous acid by O₂, O₃ and H₂O₂. *Atmosph. Envir.* 17, 2005-2011.
- Danis, R. K., Brodeur, A. E. and Shields, J. (1967). The danger of hydrogen peroxide as a colonic irrigating solution. *J. Pediatric Surg.* 2, 131.
- Das, T. N., Moorthy, P. N. and Rao, K. N. (1983). Chemiluminescent measurement of atmospheric hydrogen in the bombay area. *Atmosph. Environ.* 17, 79-82.
- Daum, P. H., Kelly, T. J., Schwartz, S. E. and Newman, L. (1984). Measurements of the chemical composition of stratiform clouds. *Atmos. Environ.* 18, 2671-2684.
- Daum, P. H., Kleinman, L. I., Hills, A. J., Lazrus, A. L., Leslie, A. C. D., Busness, K. and Boatman, J. (1990). Measurement and interpretation of concentrations of hydrogen peroxide and related species in the upper Midwest during summer. *J. of Geophysical Res.* 95, 9857-9871.
- Daum, P. H. (1990). Observations of H₂O₂ and S⁴⁺ in air, cloudwater and precipitation and their implications for the reactive scavenging of SO₂. *Atmospheric Research* 25, 89-102.
- Dawson, J., Bryant, M. G., Bloom, S. R. and Peters, T. J. (1981). Characterisation of gut hormone storage granules from normal human jejunum using metrizamide density gradients. *Regulatory Peptides* 2, 305-315.
- De Duve, C. (1965). The separation and characterisation of subcellular particles. *Harvey Lect. Ser.* 59, 48-87. [Quoted in Chance *et al.* 1979]
- De Duve, C. and Bauduin, P. (1966). Peroxisomes (microbodies and related particles). *Physiol. Rev.* 46, 323-353.
- De Flora, S., Camoirano, A., Zancchi, P. and Bennicelli, C. (1984). Mutagenicity testing with TA97 and TA102 of 30 DNA-damaging compounds, negative with other *Salmonella* strains. *Mutation Res.* 134, 159-165.
- Degussa (1977). Vorversuche zum Fishtest, Ermittlung der kritischen Konzentration an H₂O₂ in Wasser. Personal communication by Dr. Scharzkoff and Mr. Muehlig, ISEGA, US-IT 77-0013-DKO. Degussa, Hanau (D).
- Degussa (1988). DIN-Sicherheitsdatenblatt, Wasserstoffperoxid 70 G%. Degussa, Hanau, 1-3.
- Degussa (1989). DIN-Sicherheitsdatenblatt, Wasserstoffperoxid 30 G%. Degussa, Hanau, 1-3.
- DFG (Deutsche Forschungsgemeinschaft) (1991). Maximale Arbeitsplatzkonzentrationen und Biologische Arbeitsstofftoleranzwerte 1991. VCH, Weinheim, 71.
- Dick, D. A. T. (1964). The permeability coefficient of water in the cell membrane and the diffusion coefficient in the cell interior. *J. Theoret. Biol.* 7, 504-531.

- Dieter, H. H. (1988). Probleme der humantoxikologischen Verträglichkeit von Wasserstoffperoxid in Bade- und Trinkwasser aus biochemischer und pharmakokinetischer Sicht. Z. Wasser Abwasser Forsch. 21, 133-140.
- Di Paolo, J. A. (1952). Studies on chemical mutagenesis utilizing nucleic acid components, urethane, and hydrogen peroxide. The American Naturalist 86, 49-56.
- Dixon, M. (1971). The acceptor specificity of flavins and flavoproteins III, flavoproteins. Biochem. Biophys. Acta 226, 269-284.
- Dollard, G. J., Jones, B. M. R., Chandler, A. S., Gay, M. J., (1988). Measurements of ambient SO₂ and H₂O₂ at Great Dun Fell and evidence of their reaction in cloud. In: M. H. Unsworth and D. Fowler (eds), Acid deposition at high elevation sites. NATO ASI series C 252. Kluwer Academic, Dordrecht, 215-229.
- Dominguez, L., Garayazabal, J. F. F., Ferri, E. R., Vasquez, J. A. Gomez-Lucia, E., Ambrosio, C. and Suarez, G. (1987). Viability of *Listeria monocytogenes* in milk treated with hydrogen peroxide. J. Food Protection 50, 636-639.
- Dorman, H. L. and Bishop, J. G. (1970). Production of experimental edema in dog tongue with dilute hydrogen peroxide. Oral Surgery, Oral Medicine, and Oral Pathology 29, 38-43.
- Draper, W. M. and Crosby, D. G. (1983). The photochemical generation of hydrogen peroxide in natural waters. Arch. Environ. Contam. Toxicol. 12, 121-126.
- Dreosti, I. E., Baghurst, P. A., Partick, E. J. and Turner, J. (1990). Induction of micronuclei in cultured murine splenocytes exposed to elevated levels of ferrous ions, hydrogen peroxide and ultraviolet irradiation, Mutation Res. 244, 337-343.
- Du Pont (1953). Primary irritancy and skin sensitization tests, Memphis hydrogen peroxide 3%, Medical research project. Du Pont, Wilmington, DE, 1-5 + 1-2.
- Du Pont (1972a). Federal Hazardous Substances Act - Skin irritation test on rabbits, hydrogen peroxide 3% (aqueous solution), Haskell Laboratory Report. Du Pont, Wilmington, DE, 1-3.
- Du Pont (1972b). Federal Hazardous Substances Act test - Rabbit eye irritation, hydrogen peroxide 3% water, Haskell Laboratory Report. Du Pont, Wilmington, DE, 1.
- Du Pont (1972c). Eye irritation test in rabbits, hydrogen peroxide 8%, 10%, 12%, Haskell Laboratory Report. Du Pont, Wilmington, DE, 1-3.
- Du Pont (1973a). Federal Hazardous Substances Act - Skin irritation test on rabbits, 6% hydrogen peroxide (by weight), Haskell Laboratory Report. Du Pont, Wilmington, DE, 1-3.
- Du Pont (1973b). Federal Hazardous Substances Act test - Rabbit eye irritation, 6% hydrogen peroxide (by weight), Haskell Laboratory Report. Du Pont, Wilmington, DE, 2.
- Du Pont (1974a). Federal Hazardous Substances Act - Skin irritation test on rabbits, hydrogen peroxide 8%, Haskell Laboratory Report. Du Pont, Wilmington, DE, 1-2.
- Du Pont (1974b). Skin irritation test on rabbits, hydrogen peroxide 35%, Haskell Laboratory Report. Du Pont, Wilmington, DE, 1.
- Du Pont (1990a). Material Safety data sheet. Albione 35, 50, 35 CG, 50 CG, 50 M hydrogen peroxide. Du Pont, Wilmington, DE, 1-7.
- Du Pont (1990b). Material Safety data sheet. Albione 70, 70 DG, 70 CG hydrogen peroxide. Du Pont, Wilmington, DE, 1-7.
- ECETOC (European Chemical Industry Ecology and Toxicology Centre) (1991). Measurements of H₂O₂ at the workplace, personal communications from L'Air Liquide, Du Pont, Eka Nobel and Interlox. ECETOC, Brussels, 1-5.
- ECETOC (European Chemical Industry Ecology and Toxicology Centre) and IPCS (International Programme on Chemical Safety, UNEP, ILO, WHO) (1991). Early indicators of non-genotoxic carcinogenesis. Proc. joint workshop, Brussels, June 1990. Mutation Res. 248, 213-216.
- ECN (European Chemical News) (1988). H₂O₂ in oversupply but prospects are bright, ECN special report. Eur. Chem. News, May 30, 9.
- Eka Nobel (1986a). Varuinformation, vaeteperoxid 35%. Eka Nobel, Surte, S, 1-2.
- Eka Nobel (1986b). Varuinformation, vaeteperoxid 50%. Eka Nobel, Surte, S, 1-2.

- Eka Nobel (1986c). Product information, hydrogen peroxide 50%. Eka Nobel, Surte. S. 1-2.
- Eka Nobel (1988a). Varuininformation, vaeteperoxid 70%. Eka Nobel, Surte. S. 1-2.
- Eka Nobel (1988b). Product information, hydrogen peroxide 70%. Eka Nobel, Surte. S. 1-2.
- Eka Nobel (1989). Product information, Ekarox T, hydrogen peroxide 25-35%. Eka Nobel, Surte. S. 1-2.
- Eka Nobel (1990). Some estimations of H_2O_2 amounts released into waste water from different types of usage, personal communication by I. Dalin. Eka Nobel, Surte. S. 1-2.
- Emerit, I., Keck, M., Levy, A., Feingold, J. and Michelson, A. M. (1982). Activated oxygen species at the origin of chromosome breakage and sister-chromatide exchanges. *Mutation Res.*, 103, 165-172.
- England, M. D., Cavarocchi, N. C., O'Brian, J. F., Solis, E., Pluth, J. R., Orszulak, T. A., Kaye, M. P. and Schaff, H. V. (1986). Influence of antioxidants (mannitol and allopurinol) on oxygen free radical generation during and after cardiopulmonary bypass. *Circulation* 74, 134-137.
- Estervig, D. and Wang, R. J. (1984). Sister chromatid exchanges and chromosome aberrations in human cells induced by H_2O_2 and other photoproducts generated in fluorescent light-exposed medium. *Photochemistry and Photobiology* 40, 333-336.
- FCC (US Food Chemical Codex) (1981). Hydrogen peroxide, FCC 111 Monographs (3rd ed.). National Academy, New York, 146-147.
- Ferm, M. (1988). A simple volumetric technique for determination of atmospheric H_2O_2 . EMEP expert meeting on sampling, chemical analysis and quality assurance, Arona, Italy, EMEP-CCC-Rep. 4/88, 139-150.
- Flohé, L. and Schlegel, W. (1971). Glutathion-peroxidase IV. *Hoppe-Seyler's Z. Physiol. Chem.* 352, 1401-1410.
- Florence, T. M. and Stauber, J. L. (1986). Toxicity of copper complexes to the marine diatom *Nitzschia closterium*. *Aquatic Toxicology* 8, 11-26.
- FMC (1979a). Acute oral toxicity of 70% hydrogen peroxide in rats, study ICG/T79027-01. FMC, Princeton, NJ, 1.
- FMC (1979b). Acute dermal toxicity of 70% hydrogen peroxide in rabbits, study ICG/T79027-02. FMC, Princeton, NJ, 1.
- FMC (1979c). Acute eye irritation of 70% hydrogen peroxide in rabbits, study ICG/T79027-04. FMC, Princeton, NJ, 1.
- FMC (1983a). Acute oral toxicity of 35% hydrogen peroxide in rats, study I83-745. FMC, Princeton, NJ, 1.
- FMC (1983b). Acute dermal toxicity of 35% hydrogen peroxide in rabbits, study I83-746. FMC, Princeton, NJ, 1.
- FMC (1983c). Primary skin irritation and corrosion of 35% hydrogen peroxide in rabbits, study I83-747. FMC, Princeton, NJ, 1.
- FMC (1986). Acute oral toxicity of 50% hydrogen peroxide in rats, study I86-0914. FMC, Princeton, NJ, 1.
- FMC (1987a). Preliminary eye irritation of 8% hydrogen peroxide in rabbits, study I87-0950. FMC, Princeton, NJ, 1.
- FMC (1987b). Primary skin irritation/corrosion study (modified) of 70% hydrogen peroxide in one anesthetized rabbit, study I87-972. FMC, Princeton, NJ, 1-2.
- FMC (1989a). Acute oral toxicity of 10% hydrogen peroxide in rats, study I89-1077. FMC, Princeton, NJ, 1-2.
- FMC (1989b). Primary skin irritation of 10% hydrogen peroxide in rabbits, study I89-1078. FMC, Princeton, NJ, 1.
- FMC (1989c). Acute inhalation screen of 50% hydrogen peroxide in rats, study I89-1080. FMC, Princeton, NJ, 1-2.
- FMC (1990a). Material safety data, hydrogen peroxide 5% technical grade. FMC, Princeton, NJ, 1-10.
- FMC (1990b). Material safety data, hydrogen peroxide 35% standard grade. FMC, Princeton, NJ, 1-10.
- FMC (1990c). Material safety data, hydrogen peroxide 50% standard grade. FMC, Princeton, NJ, 1-10.
- FMC (1990d). Material safety data, hydrogen peroxide 70% standard grade. FMC, Princeton, NJ, 1-10.
- FMC (1990e). Occupational exposure data hydrogen peroxide. Pers. comm. by S. T. Wagner, 1 June 1990. FMC, Philadelphia, NJ, 1.

- Frenkel, K. and Chrzan, K. (1987). Hydrogen peroxide formation and DNA base modification by tumor promoter-activated polymorphonuclear leukocytes. *Carcinogenesis* 8, 455-460 [Abstract].
- Fridovich, J. (1978). The biology of oxygen radicals. *Science* 201, 875-879.
- Fridovich, I. (1983). Superoxide radical: an endogenous toxicant. *Ann. Rev. Pharmacol. Toxicol.* 23, 239-257.
- Gannon, J. E. and Gannon, S. A. (1975). Observations on the narcotization of crustacean zooplankton. *Crustaceana* 28, 2, 220-225.
- Garberg, P., Akerblom, E. L. and Bolcsfoldi, G. (1988). Evaluation of a genotoxicity test measuring DNA-strand breaks in mouse lymphoma cells by alkaline unwinding and hydroxyapatite elution. *Mutation Res.* 203, 155-176.
- Gensler, H. L. and Bowden, G. T. (1983). Evidence suggesting a dissociation of DNA strand scissions and late stage promotion of tumor cell phenotype. *Carcinogenesis* 4, 12, 1507-1511.
- Ghadially, F. N. and Wiseman, G. (1958). Inability of hydrogen peroxide to influence the survival-time of rats bearing RD3 sarcoma and the Walker carcinoma. *Nature* 4615, 1067-1069.
- Giberson, T. P., Kern, J. D., Pettigrew, D. W., Eaves, C. C. and Haynes, J. F. (1989). Near-fatal hydrogen peroxide ingestion. *Ann. Emerg. Med.* 18, 778-779.
- Giguère, P. A. (1975). Peroxyde d'hydrogène. In: Pascal, P., Compléments au nouveau traité de chimie minérale. Masson, Paris, 105-106.
- Giusti, G. V. (1973). Fatal poisoning with hydrogen peroxide. *Forensic Science* 2, 99-100.
- Glatt, H. (1989). Mutagenicity spectra in *Salmonella typhimurium* strains of glutathione, L-cysteine and active oxygen species. *Mutagenesis* 4, 221-227.
- Goor, G., Kunkel, W., Weiberg, O. (1989). Hydrogen peroxide. In: Elvers, B., Hawkins, S., Ravenscroft, M. and Schulz, G., Ullmann's encyclopedia of industrial chemistry. VCH, Weinheim, A13, 443-466.
- Gordon, S. M., Bland, L. A., Alexander, S. R., Newman, H. F., Arsuino, M. J., Jarvis, W. R. (1990). Hemolysis associated with hydrogen peroxide at a pediatric dialysis center. *Am. J. Nephrol.* 10, 123-127.
- Goth, L., Nemeth, H. and Meszaros, U. (1983). Serum catalase activity for detection of hemolytic diseases. *Clin. Chem.* 29, 741-743.
- Grant, W. M. (1974). Toxicology of the eye (2nd ed). Thomas, Springfield, IL, 559-560.
- Grant, W. M. (1986). Toxicology of the eye (3rd ed). Thomas, Springfield, IL, 492-494.
- Green, H. N. and Westrop, J. W. (1958). Hydrogen peroxide and tumour therapy. *Nature* 4602, 128-129.
- Grisham, M. B. and Kvietys, P. R. (1989). Sublethal amounts of hydrogen peroxide (H_2O_2) produce breaks in double stranded deoxyribonucleic acid (DNA) of intestinal epithelial cells and lymphocytes. *Gastroenterology* 96/5-2, A185 [Abstract].
- Guenzler, W. A., Kremers, H. and Flohé, L. (1974). An improved coupled test procedure for glutathione peroxidase (EC 1. 11. 1. 9) in blood. *Z. Klin. Chem. Klin. Biochem.* 12, 444-448.
- Gupta, C. B., Eskin, N. A. M., Frenkel, C. and Sadovski, A. Y. (1977). An improved method for the determination of residual hydrogen peroxide in milk. *J. Food Sci.* 42, 2, 537-538.
- Halliwell, B. (1974). Superoxid dismutase, catalase and glutathione peroxidase: solutions to the problems of living with oxygen. *New Phytol.* 73, 1075-1086.
- Halliwell, B. and Gutteridge, J. M. C. (1984). Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem. J.* 219, 1-14.
- Hankin, L. (1958). Hydrogen peroxide ingestion and the growth of rats. *Nature* 4647, 1453.
- Hasle, P. (1987). Protection from dangerous substances in the workplace, draft document for discussion. NICE/UNDP/ILO Project THA/82/005. ILO, Geneva, 40.
- Hauschild, F., Ludwig, R. and Muelberg, H. (1958). Ueber die "aetzende" Wirkung von Wasserstoffperoxyd. *Naunyn-Schmiedeberg's Arch. Exp. Pathol. Pharmac.* 235, 51-62.

- Heikes, B. G. (1987). Measurements of H_2O_2 and their relationship to photochemical oxidants. Presented at: Graduate School of Oceanography, Univ. Rhode Island, Narragansett, abstract 6. PL.
- Heikes, B. G., Kok, G. L., Walega, J. G. and Lazrus, A. L. (1987). H_2O_2 , O_3 and SO_2 measurements in the lower troposphere over the eastern United States during fall. *J. Geophys. Res.* 92, D1, 915-931.
- Heilpointner, E. and Gaeb, S. (1989). Detection of methyl, hydroxymethyl and hydroxyethyl hydroperoxides in air and precipitation. *Nature* 337, 631-634.
- Helz, G. R. and Kieber, R. J. (1985). Hydrogen peroxide in estuarine waters: a minor but significant contributor to chlorine demand. In: *Water chlorination, photochemistry of oxidants* 81, 1033-1040.
- Hemmerich, P., Nagelschneider, G. and Veeger, C. (1970). Chemistry and molecular biology of flavins and flavoproteins. *FEBS Lett.* 8, 69-83.
- Herrin, J. R., Squier, C. A. and Rubright, W. C. (1987). Development of erosive gingival lesions after use of a home care technique. *J. Periodontol.* 58, 11, 785-788.
- Hiatt, R. W., Naughton, J. J. and Matthews, D. C. (1953). Effects on chemicals on a schooling fish, *Kuhlia sandvicensis*. *Biol. Bull.* 104, 28-44.
- Hinchee and Downey, 1988. The role of hydrogen peroxide in enhanced bioreclamation. In: *Proc. Conf. Petroleum hydrocarbons and organic chemicals in ground water: prevention, detection and restoration*. Ass. Ground Water Scientists and Engineers (division of NWWA) and the American Petroleum Institute, Houston, TX, 715-722.
- Hincks, J. R. and Gibson, N. W. (1988/89). A comparison of the alkaline elution of DNA through filters by pumping versus gravity-flow. *In Vitro Toxicology* 2, 259-266.
- Hirota, N. and Yokoyama, T. (1981). Enhancing effect of hydrogen peroxide upon duodenal and upper jejunal carcinogenesis in rats. *Gann* 72, 811-812.
- Hochstein, P. (1988). Perspectives on hydrogen peroxide and drug-induced hemolytic anemia in glucose-6-phosphate dehydrogenase deficiency. *Free Radical Biol. Med.* 5, 387-392.
- Hochstein, P. and Atallah, A. S. (1988). The nature of oxidants and antioxidant systems in the inhibition of mutation and cancer. *Mutation Res.* 202, 363-375.
- Holm, T. R., George, G. K. and Barcelona, M. J. (1987). Fluorometric determination of hydrogen peroxide in groundwater. *Anal. Chem.* 59, 582-586.
- Holman, R. A. (1957). A Method of destroying a malignant rat tumour *in vivo*. *Nature* 4568, 1033.
- Holt, B. D. and Kumar, R. (1986). Method for determination of $\delta^{18}O$ and concentration of H_2O_2 in rainwater, preprint extended abstract, Division of Environmental Chemistry, American Chemical Society, New York, 186-187.
- Hrubetz, M. C., Conn, L. W., Gittes, H. R., MacNamee, J. K. and Williams, J. H. (1951). The cause of the increasing intravenous toxicity of 90% hydrogen peroxide with progressive dilutions. *Chemical Corps Medical Laboratories Res. Rep.* 75. Army Chemical Center, Edgewood, MA, 1-13.
- Hsie, A. W., Recio, L., Katz, D. S., Lee, C. Q., Wagner, M. and Schenley, R. L. (1986). Evidence for reactive oxygen species inducing mutations in mammalian cells. *Proc. Natl. Acad. Sci. USA* 83, 9616-9620.
- Humberston, C. L., Dean, S. D., Krenzelok, E. D. (1990). Ingestion of 35% hydrogen peroxide. *Clinical Toxicol.* 28, 95-100.
- Hwang, H., and Dasgupta, P. K. (1985). Thermodynamics of the hydrogen peroxide - water system. *Environ. Sci. Technol.* 19, 3, 255-258.
- IARC (International Agency for Research on Cancer) (1985). Hydrogen peroxide. In: *Allyl compounds, aldehydes, epoxides and peroxides*. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans 36. WHO, Geneva, 285-314.
- ILO (International Labour Office) (1991). Occupational exposure limits for airborne toxic substances (3rd ed.). *Occupational Safety and Health Series N°37*. ILO, Geneva, 224.
- INRS (Institut National de Recherches Scientifiques) (1987). Valeurs limites pour les concentrations des substances dangereuses dans l'air des lieux de travail, France. *Cahiers de notes documentaires* 129. INRS, Paris, 619.

- INRS (Institut National de Recherches Scientifiques) (1990). Valeurs limites pour les concentrations des substances dangereuses dans l'air des lieux de travail. Union Soviétique. Cahiers de notes documentaires 138, 138-139, 170.
- Interox (1990). Effluent and its treatment at NPO Fonyama works. report 89/10/30, personal communication at environmental effluent review meeting, Muenich, 12 June 1990. Interox Chemicals, Warrington, UK, 1-5.
- Interox (1991). Personal communication by C. de Rooij, 4 April 1991 [concerning workplace monitoring of H_2O_2]. Solvay Brussels, 2.
- Ishida, A. and Sasaki, T. (1981). Hydrogen peroxide induced synthesis of catalase in *Bacillus subtilis*. Kumamoto J. Sci. Biol. 15, 39-48.
- Ishidate, M., Sofuni, T., Yoshikawa, K., Hayashi, M., Nohmi, T., Sawada, M. and Matsuoka, A. (1984). Primary mutagenicity screening of food additives currently used in Japan. *Fd Chem. Toxic.* 22, 623-636.
- Ishikawa, T. and Takayama, S. (1984). Cited in Information Bulletin on the Survey of Chemicals Being Tested for Carcinogenicity. No. 11, 86.
- ISO (International Organization for Standardization) (1984). Draft international standard ISO/DIS 7157, hydrogen peroxide for industrial use. determination of hydrogen peroxide content, titrimetric method. ISO, Geneva, 1-5.
- Ito, K. A., Denny, C. B., Brown, C. K., Yao, M. and Seeger, M. L. (1973). Resistance of bacterial spores to hydrogen peroxide. *Food Technol.* Nov. 1973, 58-66.
- Ito, R., Kawamura, H., Chang, H. S., Toida, S., Matsuura, S., Hidano, T., Nakai, S., Inayoshi, Y., Matsuura, M. and Akuzawa, K. (1976). Oral safety of hydrogen peroxide acute and subacute toxicities. *Toho Igakki Zasshi (J. Medical Soc. Toho)*, 23, 531-537 [Japanese].
- Ito, A., Naito, M. and Watanabe, H. (1981a). Implication of chemical carcinogenesis in the experimental animal. Tumorigenic effect of hydrogen peroxide in mice. *Ann. Rep. of Hiroshima Univ. Res. Inst. Nuclear Medicine and Biology*, 22, 147-158 [Japanese].
- Ito, A., Watanabe, H., Naito, M. and Naito, Y. (1981b). Induction of duodenal tumors in mice by oral administration of hydrogen peroxide. *Gann* 72, 174-175.
- Ito, A., Naito, M., Naito, Y. and Watanabe, H. (1982). Induction and characterization of gastro-duodenal lesions in mice given continuous oral administration of hydrogen peroxide. *Gann* 73, 315-322.
- Ito, A., Watanabe, H., Naito, M., Naito, Y. and Kawashima, K. (1984). Correlation between induction of duodenal tumor by hydrogen peroxide and catalase activity in mice. *Gann* 75, 17-21.
- Jacob, P., Bambauer, A., Tavares, T. M. and Klockow, D. (1987). Field investigations on the variability of hydrogen peroxide concentration in atmospheric gas-phase. In: CEC, Phys.-Chem. Behav. Atmos. Pollut., rep. EUR 10832. Off. Off. Publ. EC, Luxembourg, 258-270.
- Jacob, P., Tavares, T. M., Rocha, V. C. and Klockow, D. (1990). Atmospheric hydrogen peroxide field measurements in a tropical environment: Bahia, Brazil. *Atmospheric Env.* 24A, 377-382.
- Jacob, P. and Klockow, D. (1987). Hydrogen peroxide - its role with the environmental chemistry. Presented at: Becker, H. H. and Carlier, P., 3rd French-German workshop tropospheric chemistry by laboratory studies, Lensweiler, 6-8 October 1987, Landau, D. Bergische Universitaet, Gesamthochschule Wuppertal, Fachbereich 9, Physikalische Chemie, Bericht 15.
- Joenje, H. (1989). Genetic toxicology of oxygen. *Mutation Res.* 219, 193-208.
- Johnson, K. S., Sakamoto-Arnold, A., Willason, S. T. and Beehler, C. L. (1987). Reagent-injection flow analysis: application to the determination of nanomolar levels of hydrogen peroxide in seawater. *Anal. Chim. Acta* 201, 83-94.
- Johnson, K. S., Willason, S. W., Wiesenburg, D. A., Lohrenz, S. E. and Arnone, R. A. (1989). Hydrogen peroxide in the western Mediterranean Sea: a tracer for vertical advection. *Deep-Sea Res.* 36, 2, 241-254.
- Jonas, G., Mahoney, A., Murray, J. and Gentler, S. (1988). Chemical colitis due to endoscope cleaning solutions: a mimic of pseudomembranous colitis. *Gastroenterology* 95, 1403-1408.
- Jordan, K. S., Mackey, D. and Garvey, E. (1991). Case review, a 39-year-old man with acute hemolytic crisis secondary to intravenous injection of hydrogen peroxide. *J. Emergency Nursing* 17, 8-10.

- Kaelin, R. M., Kapanci, Y. and Tschopp, J. M. (1988). Diffuse interstitial lung disease associated with hydrogen peroxide inhalation in a dairy worker. *Ann. Rev. Respiratory Disease* 137, 1233-1235.
- Kappus, H. (1987). Oxidative stress in chemical toxicity. *Arch. Toxicol.* 60, 144-149.
- Kasai, H., Crain, P. F., Kuchino, Y., Nishimura, A., Ootsuyama, A. and Tanooka, H. (1986). Formation of 8-hydroxyguanine moiety in cellular DNA by agents producing oxygen radicals and evidence for its repair. *Carcinogenesis* 7, 1849-1851.
- Kawachi, T., Yahagi, T., Kada, T., Tazima, Y., Ishidate, M., Sasaki, M. and Sugiyama, T. (1980). Cooperative programme on short-term assays for carcinogenicity in Japan. In: Montesano, R., Bartsch, H. and Tomatis, L. (eds), *Molecular and cellular aspects of carcinogen screening tests*, IARC Scient. Publ. 27. IARC, Lyon, F. 323-330.
- Kawasaki, C., Kondo, M., Nagayama, T., Takeuchi, Y. and Nagano, H. (1969). Effect of hydrogen peroxide on the growth of rats. *Shokuhin Eisegaku Zasshi (J. Jap. Food Hygiene Soc.)* 10, 68-72 [Japanese].
- Kay, S. H., Quimby P. C. Jr. and Ouzts, J. D. (1982). Hydrogen peroxide: a potential algicide for aquaculture. In: *Proc. South. Weed Sci. Soc. 35th (New Perspect. Weed Sci.)*. Delta State University, Cleveland, MS, 275-289.
- Keck, M., Stehlik, G. and Binder, W. (1980). *Mutagenitätsuntersuchungen von Wasserstoffperoxid- bzw. Wasserstoffperoxid-Katalase behandelter Milch*. *Oesterreichische Milchwirtschaft* 2, 7-14.
- Kelly, T. J., Daum, P. H. and Schwartz, S. E. (1985). Measurements of peroxides in cloudwater and rain. *J. Geophys. Res.* 90, D5, 7861-7871.
- Kensese, S. M. and Smith, L. L. (1989). Hydrogen peroxide mutagenicity towards *Salmonella typhimurium*. *Teratogenesis, Carcinogenesis and Mutagenesis* 9, 211-218.
- Keuken, M. P., Bakker, F. P., Lingerak, W. A. and Slanina, J. (1987). Flow injection analysis of hydrogen peroxide, sulfite, formaldehyde and hydroxymethanesulfonic acid in precipitation samples. *Intern. J. Environ. Anal. Chem.* 31, 263-279.
- Kieber, R. J. and Heinz, G. R. (1986). Two-method verification of hydrogen peroxide determinations in natural waters. *Anal. Chem.* 58, 2312-2315.
- Kihlstrom, M., Salminen, A. and Vihko (1986). Chronic hydrogen peroxide intake and peroxide metabolizing enzyme activities in some tissues of mice and rats. *Experientia* 42, 1018-1020.
- Kleiman, N. J., Wang, R. R. and Spector, A. (1990). Hydrogen peroxide-induced DNA damage in bovine lens epithelial cells. *Mutation Res.* 240, 35-44.
- Kleinman, L. (1986). Photochemical formation of peroxides in the boundary layer. *J. Geophys. Res.* 91, 10,889-10,904.
- Klein-Szanto, A. J. P. and Slaga, T. J. (1982). Effects of peroxides on rodent skin: epidermal hyperplasia and tumor promotion. *J. Invest. Dermatol.* 79, 30-34.
- Klockow, D. and Jacob, P. (1986). The peroxyoxalate chemiluminescence and its application to the determination of hydrogen peroxide in precipitation. In: *NATO ASI Series G6, Chemistry of multiphase atmospheric systems*. Springer Verlag, Heidelberg, 117-130.
- Knie, J., Haeke, A., Juhnke, I., Schiller, W. (1983). *Ergebnisse der Untersuchungen von chemischen Stoffen mit vier Biotests*. *Deutsche Gewässerkundliche Mitteilungen* 27, 77-79.
- Knopf, H. L. S. (1984). Reaction to hydrogen peroxide in a contact-lens wearer. *Amer. J. Ophthalmol.* June 1984, 796.
- Kok, G. L. (1980). Measurements of hydrogen peroxide in rainwater. *Atmospheric Environment* 14, 653-656.
- Kok, G. L., Damall, K. R., Winer, A. M., Pitts Jr, J. N. and Gay, B. W. (1978a). Ambient air measurements of hydrogen peroxide in the California south coast air basin. *Environ. Sci. Technol.* 12, 9, 1077-1080.
- Kok, G. L., Holler, T. P., Lopez, M. B., Nachtrieb, H. A. and Yuan, M. (1978b). Chemiluminescent method for determination of hydrogen peroxide in the ambient atmosphere. *Environ. Sci. Technol.* 12, 1072-1076.
- Kolthoff, I. M., Sandell, E. B., Meehan, E. J., Bruckenstein, S., (eds) (1969). Volumetric determinations, iodometry, hydrogen peroxide. In: *Quantitative chemical analysis* (4th ed.). MacMillan, London, 854.
- Kondrashov, V. A. (1977). On comparative toxicity of hydrogen peroxide vapours with their inhalation and dermal modes of action. *Gig. Tr. Prof. Zabol* 10, 22-25 [Russian].

- Korhonen, A., Hemminki, K. and Vainio, H. (1984). Embryotoxic effects of eight organic peroxides and hydrogen peroxide on three-day chicken embryos. *Environ. Res.* 33, 54-62.
- Koseishyo (Japanese Ministry of Health and Welfare) (1977-1981). Annual report of the cancer research (excerpts). Kosesnyo, Tokyo [Japanese]. Degussa, Hanau.
- Koseishyo (Japanese Ministry of Health and Welfare) (1980a). On the use of hydrogen peroxide in relation to food. Koseishyo, Tokyo. [Japanese]
- Koseishyo (Japanese Ministry of Health and Welfare) (1980b). Standard of usage of chemicals containing H_2O_2 , final decision, notice 24, Feb. 20, 1980. Koseishyo, Tokyo. [Japanese]
- Kurokawa, Y., Takamura, N., Matsushima, Y., Imazawa, T. and Hayashi, Y. (1984). Studies on the promoting and complete carcinogenic activities of some oxidizing chemicals in skin carcinogenesis. *Cancer Lett.* 24, 299-304.
- Lagrange, J. and Lagrange P. (1990). Evaluation de trois méthodes de détermination de H_2O_2 dans les pluies. *Analisis* 18, 37-44.
- Lal, J., Khan, B. L. and Kuhlshetha, D. C. (1985). Effect of levels of hydrogen peroxide on survival of bacteria in milk. *Asian J. Dairy Res.* 4, 1-6.
- Lazrus, A. L., Kok, G. L., Gitlin, S. N., Lind, J. A. and McLaren, S. E. (1985). Automated fluorometric method for hydrogen peroxide in atmospheric precipitation. *Anal. Chem.* 57, 917-922.
- Lazrus, A. L., Kok, G. L., Lind, J. A., Gitlin, S. N., Heikes, B. G. and Sheffer, R. E. (1986). Automated fluorometric method for hydrogen peroxide in air. *Anal. Chem.* 58, 594-597.
- Lee, J. H., Chen, Y. and Tang, I. N. (1991). Heterogeneous loss of gaseous H_2O_2 in an atmospheric air sampling system. *Environ. Sci. Technol.* 25, 339-342.
- Levin, D. E., Hollstein, M., Christman, M. F., Schwieters, E. A. and Ames, B. N. (1982). A new *Salmonella* tester strain (TA102) with AT base pairs at the site of mutation detects oxidative mutagens. *Proc. Natl. Acad. Sci. USA* 79, 7445-7449.
- Lewin, L. and Guillery, H. (1913). Die Wirkungen von Arzneimitteln und Giften auf das Auge. 2nd ed. August Hirschwald, Berlin, 721.
- Ljarskii, P. P., Gleiberman, S. E., Pankratova, G. P., Laroslavskaja, L. A. and Iurchenko, V. V. (1983). Toxicological and hygienic characterization of disinfectants based on hydrogen peroxide and its derivatives. *Gig. Sanit.* 6, 28-31 [Russian].
- Litovitz, T. L., Schmitz, B. F., Matyunas, N. and Martin, T. G (1988). 1987 Annual Report on the American Association of Poison Control Centers National Data Collection System. *Am. J. Emerg. Med.* 6, 479-515.
- Litovitz, T. L., Schmitz, B. F., Bailey, K. M. (1990). 1989 Annual Report of the American Association of Poison Control Centers National Data Collection System. *American J. Emergency Medicine* 8, 394-429.
- Liu, S. C., Trainer, M., Fehsenfeld, F. C., Parrish, D. D., Williams, E.J., Fahey, D. W., Huebler, G. and Murphy, P. C. (1987). Ozone production in the rural troposphere and the implications for regional and global ozone distribution. *J. Geophys. Res.* 92, 4191-4207.
- Little, C. (1972). Steroid hydroperoxidases as substrates for glutathione peroxidase. *Biochem. Biophys. Acta* 284, 375-381.
- Loschen, G., Azzi, A., Richter, C. and Flohé, L. (1974). Superoxide radicals as precursors of mitochondrial hydrogen peroxide. *FEBS Lett.* 42, 68-72.
- Ludewig, R. (1959). Zur intraoralen Anwendung von Wasserstoffperoxyd. *Z. Gesamte Exp. Med.* 131, 452-465.
- Ludewig, R. (1964). Hydroperoxidase-Verteilung in der Haut und transepidermale Penetration von Wasserstoffperoxid nach epikutaner Applikation. *Acta Histochem.* 19, 303-315.
- Ludewig, R. (1965). Nachweis von 18_2 in Expirationsluft und Blut waehrend sublingualer Einwirkung 18_2 -markierten Wasserstoffperoxids. *Abhandl. Deutsch. Akad. Wiss. Berlin, Kl. Chem. Geol. Biol.* 7, 549-552.
- Ludington, L. G., Hartman, S. W., Keplinger, J. E. and Williams, F. S. (1958). Incomplete rupture of the colon following hydrogen peroxide enema. *Arch. Surg.* 76, 658-660.
- MacPhee, C. and Ruelle, R. (1969). Lethal effects of 1888 chemicals upon four species of fish from western North America. *Univ. of Idaho Forest. Wildl. Range Exp. Station Bull. N°3, Moscow, ID.* 1-112.

- MacRae, W. D. and Stich, H. F. (1979). Induction of sister-chromatid exchanges in Chinese hamster ovary cells by thiol and hydrazine compounds. *Mutation Res.* 68, 351-365.
- Mahmoud, S. Z., Naguib, K., El-Nockeashy, S. and Tawfeek, N. (1984). Use of hydrogen peroxide as a dairy preservative. *Egyptian J. Dairy Sci.* 12, 163-172.
- Mallant, R. K. A. M., Slanina, J., Masuch, G. and Kettrup, A. (1988). Effects of H₂O₂-containing acidic fog on young trees. In: Symposium. Air pollution and ecosystems, Grendel, 18 May 1988. Heidelberg, Dordrecht, 306-311.
- Mallinckrodt (1989). Hydrogen peroxide solution 3%. material safety data sheet. Mallinckrodt, Du Pont, Wilmington, DE, 1-6.
- Mannervik, B. (1980). Thioltransferases. In: Jakoby, W. B. (ed.), *Enzymatic basis of detoxication*, Vol. 2. Academic Press, New York, 229-244.
- Manohar, M. and Balasubramanian, K. A. (1986). Antioxidant enzymes in rat gastrointestinal tract. *Indian J. Biochem. Biophys.* 23, 274-278.
- Marklund, S. L., Holme, E. and Hellner, L. (1982). Superoxide dismutase in extracellular fluids. *Clin. Chim. Acta* 126, 41-51.
- Martin, J. H., Bishop, J. G., Guentherman, H. and Dorman, H. L. (1968). Cellular responses of gingiva to prolonged application of dilute hydrogen peroxide. *Periodontics* 39, 208-210.
- Masuch, G. and Kettrup, A. (1986). Effects of H₂O₂-containing acidic fog on young trees. *Intern. J. Environ. Anal. Chem.* 27, 183-213.
- Masuch, G., Kicinski, H. G., Duellme, W. and Kettrup, A. (1989). Hydrogen peroxide dissolved in acidic fog as air pollutant-effects on spruce needles. *Intern. J. Environ. Anal. Chem.* 37, 161-185.
- Matkovic, B. and Novák, R. (1977). The effects of chronic peroxide intake on the peroxide metabolism enzyme activities of rat organs. *Experientia* 33, 1574-1575.
- Matz, J. and Dietze, P. (1971). Zur Bestimmung von H₂O₂ in H-Milch. *Z. Gesamte Hyg. Grenzget.* 17, 275-277.
- MCA (Manufacturing Chemists Association) (1969). Chemical safety data sheet SD-53, hydrogen peroxide. CMA, Washington, DC, 1-31.
- McAllister, W. and Cohle, P. (1984a). Acute toxicity of 35% hydrogen peroxide to bluegill sunfish (*Lepomis macrochirus*): static acute toxicity report #31979, prepared by Analytical Bio-Chemistry Laboratories. US EPA, Pesticide document management system (PDMS), report PDR20804. [confidential]
- McAllister, W. and Cohle, P. (1984b). Acute toxicity of 35% hydrogen peroxide to rainbow trout (*Salmo gairdneri*): static acute toxicity report #31979, prepared by Analytical Bio-Chemistry Laboratories. US EPA, Pesticide document management system (PDMS), report PDR20804. [confidential]
- McKee, J. E. and Wolf, H. W. (1963). *Water quality criteria*, 2nd ed. State Water Quality Control Board, Pasadena, CA.
- McNally, O. D. (1990). Clinical aspects of topical application of dilute hydrogen peroxide solutions. *CLAO J.* 16, S46.
- Mehnert, K., Duering, R., Vogel, W. and Speit, G. (1984a). Differences in the induction of SCEs between human whole blood cultures and purified lymphocyte cultures and the effect of an S9 mix. *Mutation Res.* 130, 403-410.
- Mehnert, K., Vogel, W., Benz, R. and Speit, G. (1984b). Different effects of mutagens on sister chromatid exchange induction in three chinese hamster cell lines. *Environmental Mutagenesis* 6, 573-583.
- Mello Filho, A. C. and Meneghini, R. (1984). *In vivo* formation of single-strand breaks in DNA by hydrogen peroxide is mediated by the Haber-Weiss reaction. *Biochem. Biophys. Acta* 781, 56-63.
- Meneghini, R. (1988). Genotoxicity of active oxygen species in mammalian cells. *Mutation Res.* 195, 215-230.
- Merrifield, G. (1988). Fire and explosion hazards associated with the storage and handling of hydrogen peroxide, rep. 19. Health and Safety Executive, Technology Division, Merseyside, 1-7.
- Meyer, C. T., Brand, M., DeLuca, V. A. and Spiro, H. M. (1981). Hydrogen peroxide colitis: a report of three patients. *J. Clin. Gastroenterol.* 3, 31-35.
- Miller, S. C., Sorrin, S. S., Greenhut, W. H. and Pelzer, R. H. (1938). Hydrogen peroxide and sodium perborate: their comparative oral irritant action. *J. Am. Dental Assoc.* 25, 1957-1973.

- Miller, H. A. (1958). Les brulures du globe oculaire. Bull. Soc. Ophthalmol. France 4, 86.
- Mills, G. C. (1960). Glutathione peroxidase and the destruction of hydrogen peroxide in animal tissues. Arch. Biochem. Biophys. 86, 1-5.
- Ministere de la Solidarité, de la Santé et de la Protection Sociale (1990). Utilisation du peroxyde d'hydrogene pour le traitement des eaux destinées à la consommation humaine, autorisation DGS/PGE/1. D., n°242 du 20 Fev. 1990. DG Santé, Paris, 1-2.
- Misra, H. P. and Fridovich, J. (1972). The generation of superoxide radical during the autoxidation of hemoglobin. J. Biol. Chem. 247, 6960-6962.
- Mitchell, I. De G. (1974). A comparison of the sensitivity and specificity of microbial systems for assessing genetic damage. Agents and Actions 4, 286-294.
- Mitsubishi (1981). Acute toxicity test on 60 per cent aqueous hydrogen peroxide solution. In: IMCO Sub-Committee on Bulk Chemicals. BHC/66 Appendix 1. Mitsubishi Gas Chemical Company, Environmental Section, 2-7.
- Miyasaki, K. T., Genco, R. J. and Wilson, M. E. (1987). Influence of growth phase on the sensitivity of *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* to the bactericidal effects of hydrogen peroxide. Microbios Letters 34, 117-122.
- Miyasaki, K. T., Wilson, M. E., Zambon, J. J. and Genco, R. J. (1985). Influence of endogenous catalase activity on the sensibility of the oral bacterium *Actinobacillus actinomycetemcomitans* and the oral haemophili to the bactericidal properties of hydrogen peroxide. Archs oral Biol. 30, 843-848.
- Moffett, J. M. and Zika, R. G. (1987). Reaction kinetics of hydrogen peroxide with copper and iron in seawater. Environ. Sci. Technol. 21, 804-810.
- Mohnen, V. A. and Kadlecak, J. A. (1989). Cloud chemistry research at Whiteface Mountain. Tellus 41B, 79-91.
- Montefluos (1990a). Hydrogen peroxide 35%, material safety data sheet. Montefluos, Italy, 1-2.
- Montefluos (1990b). Hydrogen peroxide 50%, material safety data sheet. Montefluos, Italy, 1-2.
- Montefluos (1990c). Hydrogen peroxide 60%, material safety data sheet. Montefluos, Italy, 1-2.
- Mopper, K. and Zika, R. G. (1987). Natural photosensitizers in sea water: riboflavin and its breakdown products. In: Photochemistry of environmental aquatic systems. Zika, R. G. and Cooper, W. J. (ed). American Chemical Society, Washington, DC, 174-190.
- Moraes, E. C., Keyse, S. M. and Tyrrell, R. M. (1990). Mutagenesis by hydrogen peroxide treatment of mammalian cells: a molecular analysis. Carcinogenesis 11, 283-293.
- Moriyama, I., Hiroaka, K., Fujita, M., Ichijo, M. and Ioka, H. (1982). Effects of food additive hydrogen peroxide studied in fetal development. Acta Obst. Gynaec. Jpn 34, 2149-2154 [Japanese].
- Morse, D. E., Duncan, H., Hooker, N. and Morse, A. (1976). Hydrogen peroxide induces spawning in mollusks, with activation of prostaglandin endoperoxide synthetase. Science 196, 298-300.
- Mouret, J. F., Odin, F., Polverelli, M. and Cadet, J. (1990). 32P-postlabeling measurement of adenine N-1-oxide in cellular DNA exposed to hydrogen peroxide. Chem. Res. Toxicol. 3, 102-110.
- Naguib, K. (1972). Hydrogen peroxide treatment of milk for Domiati cheese. Australian J. Dairy Technol., June 1972, 56-60.
- Nahum, A., Wood, L. D. H., and Sznajder, J. I. (1989). Measurement of hydrogen peroxide in plasma and blood. Free Radical Biol. Med. 6, 479-484.
- Nakamura, S., Oda, Y., Shimada, T., Oki, I. and Sugimoto, K. (1987). SOS-inducing activity of chemical carcinogens and mutagens in *Salmonella typhimurium* TA1535/pSK1002: examination with 151 chemicals. Mutation Res. 192, 239-246.
- Nakane, K. and Kosaka, A. (1980). Applications of a hydrogen peroxide electrode in clinical chemistry. Rinsho Kagaku Shimpoujumu 20, 85-89 [Japanese].
- Nassi-Calo, L., Mello-Filho, A. C., and Meneghini, R. (1989). o-Phenanthroline protects mammalian cells from hydrogen peroxide-induced gene mutation and morphological transformation. Carcinogenesis 10, 1055-1057.
- Nettel, A., Jacob, P. and Klockow, D. (1984). Measurements of hydrogen peroxide in polar ice samples. Nature 311, 43-45.

- Neftel, A., Jacob, P. and Klockow, D. (1986). Long-term record of H_2O_2 in polar ice cores. *Tellus* 38B, 262-270.
- Nicholls, P. (1972). Contributions of catalase and glutathione peroxidase to red cell peroxide removal. *Biochim. Biophys. Acta* 279, 306-309.
- NIOSH (US National Institute for Occupational Safety and Health) (1990). NIOSH pocket guide to chemical hazards. US Government Printing Office, Washington, DC.
- Noni, H. and Hegner, D. (1978). Do mitochondria produce oxygen radicals *in vivo*. *Eur. J. Biochem.* 82, 563-567.
- Oberst, F. W., Comstock, C. C. and Hackley, E. B. (1954). Inhalation toxicity of ninety per cent hydrogen peroxide vapour. *AMA Arch. Ind. Hyg. Occup. Med.* 10, 319-327.
- Ogata, M. (1991). Acatlasemia. *Hum. Genet.* 86, 331-340.
- Olson, M. J. (1988). DNA Strand breaks induced by hydrogen peroxide in isolated rat hepatocytes. *J. Toxicol. Environ. Health* 23, 407-423.
- Olson, D. L. and Christensen, G. M. (1980). Effects on water pollutants and other chemicals on fish acetylcholinesterase (*in vitro*). *Environ. Res.* 21, 327-335.
- Olszyna, K. J., Meagher, J. F. and Bailey, E. M. (1988). Gas-phase, cloud and rain-water measurements of hydrogen peroxide at a high-elevation site. *Atmospheric Env.* 22, 1699-1706.
- OSHA (US Occupational Safety and Health Administration, Dep. of Labor) (1989a). Industrial exposure and control technologies for OSHA regulated hazardous substances. PB89-210199. NTIS, Springfield, VA, 1060-1062.
- OSHA (US Occupational Safety and Health Administration, Dep. of Labor) (1989b). Rules and regulations. Air contaminants. *Fed. Reg.* 54, 2332, 2940.
- Oshino, N., Oshino, R. and Chance, B. (1974). The properties of catalase "peroxidatic" reaction and its relationship to microsomal methanol oxidation. In: Thurman, R. G., Yonetani, T., Williamson, J. R. and Chance, B. (eds), Alcohol and aldehyde metabolizing systems. Academic Press, New York, 231-242.
- Oya, Y., Yamamoto, K. and Tonomura, A. (1986). The biological activity of hydrogen peroxide. I. Induction of chromosome-type aberrations susceptible to inhibition by scavengers of hydroxyl radicals in human embryonic fibroblasts. *Mutation Res.* 172, 245-253.
- Paugh, J. R., Brennan, N. A. and Efron, N. (1988). Ocular response to hydrogen peroxide. *Am. J. Optometry and Physiol. Optics* 65, 91-98.
- Petasne, R. G. and Zika, R. G. (1987). Fate of superoxide in coastal sea water. *Nature* 325, 516-518.
- Pharmacopée Européenne (1985). Hydrogenii peroxidum 3 per centum, solution de peroxyde d'hydrogène à 3 pour cent; Hydrogenii peroxidum 30 per centum, solution de peroxyde d'hydrogène à 30 pour cent. Conseil de l'Europe, Strasbourg, 395-396.
- Pilz, W. and Johann, I. (1974). Die Bestimmung kleinster Mengen von Wasserstoffperoxid in Luft. *Intern. J. Environ. Anal. Chem.* 3, 257-270.
- Pirie, A., Rees, J. R. and Holmberg, N. J. (1970). Diquat cataract: formation of the free radical and its reaction with constituents of the eye. *Exptl. Eye Res.* 9, 204-218.
- Possanzini, M., Di Palo, V. and Liberti, A. (1988). Annular denuder method for determination of hydrogen peroxide in the ambient atmosphere. *Sci. Tot. Environ.* 77, 203-214.
- Prise, K. M., Davies, S. and Michael, B. D. (1989). Cell killing and DNA damage in Chinese hamster V79 cells treated with hydrogen peroxide. *Int. J. Radiat. Biol.* 55, 583-592.
- Pumphrey, R. E. (1951). Hydrogen peroxide proctitis. *Amer. J. Surgery* 81, 60-62.
- Punte, C. L., Saunders, L. E. and Krakow, E. H. (1953). The inhalation toxicity of aerosols of 90% hydrogen peroxide. Chemical Corps Medical Laboratories, Res. Rep. 189. Army Chemical Center, Maryland, 1-11.
- Québec (1989). Règlement sur la qualité du milieu de travail. Editeur officiel du Québec, Québec, 41.

- Quimby, P. C., Jr. (1981). Preliminary evaluation of hydrogen peroxide as a potential herbicide for aquatic weeds. *J. Aquat. Plant Manage.* 19, 53-55.
- Rackoff, W. R. and Merton, D. F. (1990). Gas embolism after ingestion of hydrogen peroxide. *Pediatrics* 85, 593-594.
- Rees, T. D. and Orth, C. F. (1986). Oral ulcerations with use of hydrogen peroxide. *J. Periodontology* 57, 689-692.
- Regnier, J.-F. and Imbert, C. (1990). Evaluation of the EYETEX system as a screen for predicting the ocular irritancy potential of chemical products. Atochem, Paris-La-Défense. [submitted for publication in ATLA]
- Richards, L. W., Anderson, J. A., Blumenthal, D. L., McDonald, J. A., Kok, G. L. and Lazrus, A. L. (1983). Hydrogen peroxide and sulfur (IV) in Los Angeles cloud water. *Atmospheric Environment* 17, 911-914.
- Richards, L. W. (1989). Effects of air pollution on western forests. Air Pollution Control Association transaction series. Air and Waste Management Association, Pittsburgh, PA, 51-72.
- Roberts, N. and Philips, C. (1989a). The acute toxicity of 35% hydrogen peroxide to the mallard duck: HRC report BWZ6/841135, prepared by Huntingdon Research Centre. US EPA, Pesticide document management system (PDMS), report PDR20B04. [confidential]
- Roberts, N. and Philips, C. (1989b). The dietary toxicity (LC₅₀) of 35% hydrogen peroxide to the bobwhite quail: HRC report BWZ6/841135, prepared by Huntingdon Research Centre. US EPA, Pesticide document management system (PDMS), report PDR20B04. [confidential]
- Roberts, N. and Philips, C. (1989c). The dietary toxicity (LC₅₀) of 35% hydrogen peroxide to the mallard duck: HRC report BWZ4/841300, prepared by Huntingdon Research Centre. US EPA, Pesticide document management system (PDMS), report PDR20B04. [confidential]
- Roemer, F. G., Viljeer, J. W., van den Beld, L., Slangewal, H. J., Veldkamp, A. A. and Reijnders, H. F. R. (1985). *Atmospheric Env.* 19, 1847-1858.
- Romanovski, A., Murray, J. R. and Huston, M. J. (1960). Effects of hydrogen peroxide on normal and hypertensive rats. *Pharm. Acta. Helv.* 35, 354-357.
- Rondoni, P. and Cudkowicz, G. (1953). Hydrogen peroxide in tumours; its possible significance in carcinogenesis. *Experientia* 9, 348-349.
- Roots, R. and Okada, S. (1975). Estimation of life times and diffusion distances of radicals involved in X-ray-induced DNA strand breaks or killing of mammalian cells. *Radiat. Res.* 64, 306-320.
- Rowlatt, C., Chesterman, F. and Sheriff, M. (1976). Lifespan, age changes and tumor incidence in an aging C57B1/mouse colony. *Lab. Animals* 10, 419-442.
- Rowlatt, C., Franks, L., Sheriff, M. and Chesterman, F. (1969). Naturally occurring tumors and other lesions of the digestive tract in untreated C57B1 mice. *J. Nat. Cancer Institute* 43, 1353-1364.
- Ruch, R. J., Crist, K. A. and Klaunig, J. E. (1989). Effects of culture duration on hydrogen peroxide-induced hepatocyte toxicity. *Toxicol. Appl. Pharmacol.* 100, 451-464.
- Ruiz-Rubio, M., Alejandro-Durán, E. and Pueyo, C. (1985). Oxidative mutagens specific for A-T base pairs induce forward mutations to L-arabinose resistance in *Salmonella typhimurium*. *Mutation Res.* 147, 153-163.
- Sacks, L. E. and MacGregor, J. T. (1982). The *B. subtilis* multigene sporulation test for mutagens: detection of mutagens inactive in the *Salmonella* his reversion test. *Mutation Res.* 95, 191-202.
- Saito, T., Kurasaki, M., Kaji, M. and Saito, K. (1984). Deficiency of erythrocyte superoxide dismutase and catalase activities in patients with malignant lymphoma and acute myeloid leukemia. *Cancer Lett.* 24, 141-146.
- Sakugawa, H. and Kaplan, I. R. (1987). Atmospheric hydrogen peroxide measurement: comparison of cold trap method with impinger bubbling method. *Atmospheric Env.* 21, 1791-1798.
- Sakugawa, H. and Kaplan, I. R. (1989). Hydrogen peroxide in the atmosphere of Los Angeles and its vicinity: factors controlling their formation and their role as oxidants of SO₂. *J. Geophys. Res.* 94, 12,957-12,973.
- Sakugawa, H., Kaplan, I. R., Tsai, W., Cohen, Y. (1990). Atmospheric hydrogen peroxide. *Environ. Sci. Technol.* 24, 1452-1462.

- Saladino, A. J., Willey, J. C., Lechner, J. F., Grafstrom, R. C., LaVeck, M. and Harris, C. C. (1985). Effects of formaldehyde, acetaldehyde, benzoyl peroxide, and hydrogen peroxide on cultured normal human bronchial epithelial cells. *Cancer Res.* 45, 2522-2526.
- Sanford, K. K., Parshad, R. and Ganitt, R. (1986). Responses of human cells in culture to hydrogen peroxide and related free radicals generated by visible light: relationship to cancer susceptibility. In: Liss, A. T. (ed.). *Free radicals, aging, and degenerative diseases*, 373-394.
- Sasaki, M., Sugimura, K., Yoshida, M. A. and Abe, S. (1980). Cytogenetic effects of 60 chemicals on cultured human and Chinese hamster cells. *Kromosomo* 20, 574-584.
- Sawada, M., Sofuni, T. and Ishidate, M. (1988). Induction of chromosomal aberration in active oxygen-generating system II: A study with hydrogen peroxide-resistant cells in culture. *Mutation Res.* 197, 133-140.
- Schildknecht, H. and Holubek, K. (1961). Die Bombardierkaefer und ihre Explosionschemie. *Angew. Chem.* 73, 1-6.
- Schirmann, J. -P., Delavarenne, S. Y. (1979). *Hydrogen peroxide in organic chemistry. Edition et Documentation Industrielle (EDI)*, Paris.
- Schoeneich, J. (1967). The induction of chromosomal aberrations by hydrogen peroxide in strains of ascites tumors in mice. *Mutation Res.* 4, 384-388.
- Schraufstatter, I. U., Hinshaw, D. B., Hyslop, P. A., Spragg, R. G. and Cochrane, C. G. (1986). Oxidant injury of cells. DNA strand-breaks activate polyadenosine diphosphate-ribose polymerase and lead to depletion of nicotinamide adenine dinucleotide. *J. Clin. Invest.* 77, 1312-1320.
- Schumb, W. C., Satterfield, C. N., Wentworth, R. L. (eds) (1955). *Hydrogen Peroxide*. Reinhold Publishing, New York, 64-68, 181, 458-464, 535-547.
- Shamberger, R. J. (1972). Increase of peroxidation in carcinogenesis. *J. Nat. Cancer Inst.* 48, 5, 1491-1496.
- Shapiro, M., Brat, V. and Ershoff, B. H. (1960). Induction of dental caries and pathological changes in periodontium of rat with hydrogen peroxide and other oxidizing agents. *J. Dental Res.* 39, 332-343.
- Shaw, A., Cooperman, A. and Fusco, F. (1967). Gas embolism produced by Hydrogen Peroxide. *New England J. of Med.*, 277, 238-241.
- Sheehan, J. F. and Brynjolfsson, G. (1960). Ulcerative colitis following hydrogen peroxide enema: Case report and experimental production with transient emphysema of colonic wall and gas embolism. *Lab. Invest.* 9, 150-168.
- Shurtleff, L. E. (1989). Interox America sodium percarbonate and hydrogen peroxide - Acute toxicity to the freshwater invertebrate *Daphnia pulex*. Burlington Research, Burlington NC, 1-40.
- Sies, H. and Chance B. (1970). The steady state level of catalase compound I in isolated hemoglobin-free perfused rat liver. *FEBS Letters* 11, 172-176.
- Sies, H. (1974). Biochemistry of the peroxisome in the liver cell. *Angew. Chem. Int. Edit.* 13, 706-718.
- Sies, H. (1981). Measurement of hydrogen peroxide formation in situ. In: Jakoby, W. B. (ed.). *Methods in Enzymology* 77, detoxification and drug metabolism, conjugation and related systems. Academic Press, New York, 15-20.
- Sies, H. (1985). Oxidative stress: introductory remarks. In: Sies, H. (ed.), *Oxidative stress*. Academic Press, London, 1-8.
- Slanetz, L. W. and Brown, E. A. (1949). Studies on the numbers of bacteria in the mouth and their reduction by the use of oral antiseptics. *J. Dental Res.* 28, 313-323.
- Slomr, F., Harris, G. W., Hastie, D. R., Mackay, G. I. and Schiff, H. I. (1986). Measurement of gas phase hydrogen peroxide in air by tunable diode laser absorption spectroscopy. *J. of Geophys. Res.* 91, 5371-5378.
- Solvay (1984a). Material safety data sheet, products N° 61, hydrogen peroxide, with medical appendix. Solvay, Brussels, 1-5.
- Solvay (1984b). Fiche de sécurité, produits N° 61, peroxide d'hydrogene. Solvay, Bruxelles, 1-3.
- Solvay (1991). Concentrations en phase vapeurs pour des solutions de peroxyde d'hydrogene. memo process engineering H₂O₂, N°10. Solvay, Bruxelles, 7 p.

- Solyom, P. (1973). Examination of the impact of H_2O_2 on biological treatment of municipal waste water in the activated sludge process. report IVL. Stockholm. Eka Nobel, Surte. S [Swedish].
- Southgate, B. A. (1950). Polluting effects of sewage and industrial wastes, toxicity of substances to fish. Report for the year 1950. Water Pollution Research Board.
- Spain, J. C., Milligan J. D., Downey, D. C. and Slaughter, J. K. (1989). Excessive bacterial decomposition of H_2O_2 during enhanced biodegradation. *Ground Water* 27, 163-167.
- Spector, A. and Garner, W. H. (1981). Hydrogen peroxide and human cataract. *Exp. Eye Res.* 33, 673-681.
- Speit, G. (1986). The relationship between the induction of SCEs and mutations in Chinese hamster cells. I. Experiments with hydrogen peroxide and caffeine. *Mutation Res.* 174, 21-26.
- Speit, G., Vogel, W., and Wolf, M. (1982). Characterization of sister chromatid exchange induction by hydrogen peroxide. *Environmental Mutagenesis* 4, 135.
- SRI International (1988). Hydrogen peroxide. summary. United States and Canada, Japan, Western Europe. In: Heydorn, B., Mulach, R., Yoshikawa, S. (eds), *Chemical Economics Handbook*, CEH product review. SRI International, Menlo Park, CA, 741, 5000 D, F. E.
- Stevens, S. E., Jr., Pat Paterson, C. O. and Myers, J. (1973). The production of hydrogen peroxide by blue-green algae: a survey. *J. Phycol.* 9, 427-430.
- Stich, H. F. and Dunn, B. P. (1986). Relationship between cellular levels of beta-carotene and sensitivity to genotoxic agents. *Int. J. Cancer* 38, 713-717.
- Stich, H. F., Wei, L. and Lam, P. (1978). The need for a mammalian test system for mutagens: action of some reducing agents. *Cancer Letters* 5, 199-204.
- Stockwell, W. R. (1986). A homogeneous gas phase mechanism for use in a regional acid deposition model. *Atmos. Environ.* 20, 1615-1632.
- Suenaka, T., Akasaka, S. and Hirata, M. (1984). Proceedings of Osaka Prefectural Institute of Public Health. *Industrial Health*, N°22, 9-13. [Japanese]
- Svirbely, J. L., Dobrogorski, O. J. and Stokinger, H. M. (1961). Enhanced toxicity of ozone-hydrogen peroxide mixtures. *Amer. Ind. Hyg. Assoc. J.*, Feb. 1961, 21-26.
- Tachon, P. (1990). Intracellular iron mediates the enhancing effect of histidine on the cellular killing and clastogenicity induced by H_2O_2 . *Mutation Res.* 228, 221-228.
- Tachon, P. and Giacomoni, P. U. (1989). Histidine modulates the clastogenic effect of oxidative stress. *Mutation Res.*, 211, 103-109.
- Takahashi, M., Hasegawa, R., Furukawa, F., Toyoda, K., Sato, H. and Hayashi, Y. (1986). Effects of ethanol, potassium metabisulfite, formaldehyde and hydrogen peroxide on gastric carcinogenesis in rats after initiation with N-methyl-N'-nitro-N-nitrosoguanidine. *Gann* 77, 118-124.
- Tamura, H., Iida, T., Watanabe, T. and Suga, T. (1990). Long-term effects of peroxisome proliferators on the balance between hydrogen peroxide-generating and scavenging capacities in the liver of Fischer-344 rats. *Toxicol.* 63, 199-213.
- Tanner, R. L., Markovits, G. Y., Ferreri, E. M. and Kelly, Th. J. (1986). Sampling and determination of gas-phase hydrogen peroxide following removal of ozone by gas-phase reaction with nitric oxide. *Anal. Chem.* 58, 1857-1865.
- Test, S. T. and Weiss, S. J. (1984). Quantitative and temporal characterization of the extracellular H_2O_2 pool generated by human neutrophils. *J. Biol. Chem.* 259, 399-405.
- Tetra Pak (1991). Measurement of hydrogen peroxide in air in a plant for aseptic packaging. Personal communication by H. Moeller, Tetrapak. Eka Nobel, Surte, S. 1.
- Thacker, J. and Parker, W. F. (1976). The induction of mutation in yeast by hydrogen peroxide. *Mutation Research* 38, 43-52.
- Thielmann, H. W. and Gersbach, H. (1978). The nucleotide-permeable *Escherichia coli* cell, a sensitive DNA repair indicator for carcinogens mutagens, and antitumor agents binding covalently to DNA. *Z. Krebsforsch.* 92, 177-214.

- Thompson, A., Owens, M. A. and Stewart, R. W. (1988). Sensitivity of tropospheric hydrogen peroxide to global chemical and climate change. *Geophys. Res. Lett.* 16, 53-56.
- Thompson, J. D. (1989). New health food hazard - 35% hydrogen peroxide. *Vet. Hum. Toxicol.* 31, 339.
- Thompson, A. M., Owens, M. A. and Stewart, R. W. (1989). Sensitivity of tropospheric hydrogen peroxide to global chemical and climate change. *Geophys. Res. Lett.* 16, 1, 53-56.
- Toyoda, M., Ito, Y., Iwaida, M. and Fujii, M. (1982). Rapid procedure for the determination of minute quantities of residual hydrogen peroxide in food by using a sensitive oxygen electrode. *J. Agric Food Chem.* 30, 346-349.
- Troll, W. and Wiesner, R. (1985). The role of oxygen radicals as a possible mechanism of tumor promotion. *Ann. Rev. Pharmacol. Toxicol.* 25, 509-528.
- Tsuda, H. (1981). Chromosomal aberrations induced by hydrogen peroxide in cultured mammalian cells. *Jpn. J. Genet.* 56, 1-8.
- Tucker, J. D., Taylor, R. T., Christensen, M. L., Strout, C. L., Hanna, L. M. and Carrano, A. V. (1989). Cytogenetic response to 1,2-dicarbonyls and hydrogen peroxide in Chinese hamster ovary AUXB1 cells and human peripheral lymphocytes. *Mutation Res.* 224, 269-279.
- Uenluetuerk, A. and Turantas, F. (1987). Bactericidal effect of hydrogen peroxide on *Salmonella typhimurium* in liquid whole egg. *J. Applied Bacteriology* 62, 25-28.
- UK HSE (Health and Safety Executive) (1991). Occupational exposure limits 1991, EH40/91. HMSO, London, 18.
- Urschel, H. C. (1967). Progress in cardiovascular surgery: Cardiovascular effects of hydrogen peroxide: current status. *Dis. Chest* 51, 180-192.
- Ursini, F., Maiorino, M. and Gregolin, C. (1985). The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochem. Biophys. Acta* 839, 62-70.
- US DOT (US Department of Transport) (1987). Emergency response guidebook 1987. In: NLM (US National Library of Medicine) database print-out of 90. 01. 15. US Govt. Print. Off., Washington, DC, G45-G60.
- US FDA (Food and Drug Administration) (1974). Scientific literature reviews on generally recognized as safe (GRAS) food ingredients, hydrogen peroxide, rep. FDABF-GRAS-261. NTIS, Springfield, VA, PB-241 957, 1-137.
- US FDA (Food and Drug Administration) (1983). Hydrogen peroxide, proposed affirmation of GRAS status as a direct human food ingredient with specific limitations. *Fed. Reg.* 48, 223, 52323-52333.
- US FDA (Food and Drug Administration) (1988). Production of hydrogen peroxide in irradiated foods. *Fed. Reg.* 53, 251, 53198.
- US FDA (Food and Drug Administration) (1990). Maximum usage levels permitted [in food]. *Code of Fed. Reg.* 1, 4-1-90 ed., 452-453.
- Vala, M. (1965). Léčení rohovkových atekcí 20% roztokem peroxidu vodíku [Treatment of corneal diseases with 20% solution of peroxide of hydrogen]. *Cs. Oftalmologie.* 21, 357-359. [Czech]
- Van Baalen, C. and Marier, J. E. (1966). Occurrence of hydrogen peroxide in sea water. *Nature* 5052, 951.
- Van Valin, C. C., Ray, J. D., Boatman, J. F. and Gunter, R. L. (1987). Hydrogen peroxide in air during winter over the south-central United States. *Geophys. Res. Lett.* 14, 11, 1146-1149.
- Van Valin, C. C., Luria, M., Ray, J. D. and Boatman, J. F. (1990). Hydrogen peroxide and ozone over the Northeastern United States in June 1987. *J. Geophys. Res.* 95, 5689-5695.
- Varma, S. D. and Devamanoharan, P. S. (1991). Hydrogen peroxide in human blood. *Free Rad. Comms.* 14, 125-131.
- Von der Hude, W., Behm, C., Guertler, R. and Basler, A. (1988). Evaluation of the SOS chromotest. *Mutation Res.* 203, 81-94.
- Vuillaume, M. (1987). Reduced oxygen species, mutation, induction and cancer initiation. *Mutation Res.* 186, 43-72.
- Wales, R. G., White, I. G. and Lamond, D. R. (1959). The spermicidal activity of hydrogen peroxide *in vitro* and *in vivo*. *J. Endocrin.* 18, 236-244.

- Wangenheim, J. and Bolcsfoldi, G. (1988). Mouse lymphoma L5178Y thymidine kinase locus assay of 50 compounds. *Mutagenesis* 3, 193-205.
- Warm, E. and Laties, G. G. (1982). Quantification of hydrogen peroxide in plant extracts by the chemiluminescence reaction with luminol. *Phytochemistry* 21, 827-831.
- Wayne, R. P. (1988). The photochemistry of ozone. In: Hutzinger, O. (ed.), *The handbook of environmental chemistry 2E*, reactions and processes. Springer, Berlin, 10.
- Weiberg, O. and Leuchtenberger, W. (1978). Feuer- und Explosions-gefahren bei der Handhabung von Wasserstoffperoxid und organischen Peroxiden. In: Weigert, W. (ed.), *Wasserstoff und seine Derivate*. Chemie und Anwendung. Huethig, Heidelberg, 199-205.
- Weiner, M. L., Freeman, C., McCarty, J. D., Aguinaldo, E. R. and Fletcher, M. J. (1990). Eye irritation studies on three concentrations of hydrogen peroxide, acute toxicity data. *J. Am. Coll. Tox.* part B, 49-50.
- Weitzman, S. A., Weitberg, A. B., Stossel, T. P., Schwartz, J. and Shklar, G. (1986). Effects of hydrogen peroxide on oral carcinogenesis in hamsters. *J. Periodontol.* 57, 685-688.
- Wilcox, P., Naidoo, A., Wedd, D. J. and Gatehouse, D. G. (1990). Comparison of *Salmonella typhimurium* TA102 with *Escherichia coli* WP2 tester strains. *Mutagenesis* 5, 285-291.
- Williams, M. D., Leigh, J. S. and Chance, B. (1982). Hydrogen peroxide in human breath and its probable role in spontaneous breath luminescence. *Annals New York Academy of Sciences* 386, 478-483.
- Wilmer, J. W. G. M. and Natarajan, A. T. (1981). Induction of sister-chromatid exchanges and chromosome aberrations by γ -irradiated nucleic acid constituents in CHO cells. *Mutation Res.* 88, 99-107.
- Winquist, L., Rannug, U., Rannug, A. and Ramel, C. (1984). Protection from toxic and mutagenic effects of H_2O_2 by catalase induction in *Salmonella typhimurium*. *Mutation Res.* 141, 145-147.
- Winterbourn, C. C. and Stern, A. (1987). Human red cell scavenge extracellular hydrogen peroxide and inhibit formation of hydrochlorous acid and hydroxyl radical. *J. Clin. Invest.* 80, 1486-1491.
- Yamagata, S., Seino, S. and Nakao, T. (1952). Studies on blood and plasma catalase; a new micromethod of estimation of blood and plasma catalase. *Tohoku J. Exp. Med.* 57, 85-92.
- Yamagata, S. and Seino, S. (1953). Studies on blood and plasma catalase; the blood and plasma catalase content in blood diseases. *Tohoku J. Exp. Med.* 57, 101-107.
- Yoshizumi, K., Aoki, K., Nouchi, I., Okita, T., Kobayashi, T., Kamakura, S., Tajima, M. (1984). Measurements of the concentration in rainwater and of the Henry's Law constant of hydrogen peroxide. *Atmospheric Environment* 18, 395-401.
- Yosphe, Purer and Eylan (1968). Cited in: Vanbrabant, R. (1976), Disinfection by hydrogen peroxide of effluents of wastewater purification plants. Thesis, Katholieke Universiteit te Leuven, Leuven, B, 8-9.
- Zabelin, A. I. and Karbovnichii, P. N. (1983). Optimum conditions for the determination of trace amounts of hydrogen peroxide with a phenolphthalein leuco base. *Energetik*, 6, 26-27 [Russian; Chem. Abstr. 99, 76526r].
- Zecevic, D. and Gasparec, Z. (1979). Death caused by hydrogen peroxide. *Z. Rechtsmed.* 84, 57-59.
- Zepp, R. G., Skurlatov, Y. I. and Pierce, J. T. (1987). Algal-induced decay and formation of hydrogen peroxide in water: its possible role in oxidation of anilines by algae. In: *Photochemistry of Environmental Aquatic Systems*. American Chemical Society, symp. ser. 327, 215-224.
- Zhou, R., Sun, X. and Tang, D. (1991). Iron-mediated induction of the SOS responses by hydrogen peroxide. *Bull. Environ. Contam. Toxicol.* 46, 613-617.
- Ziegler-Skylakakis, K. and Andrae, U. (1987). Mutagenicity of hydrogen peroxide in V79 Chinese hamster cells. *Mutation Res.* 192, 65-67.
- Zika, R. G. and Saltzman, E. S. (1982). Interaction of ozone and hydrogen peroxide in water, implications for analysis of H_2O_2 in air. *Geophys. Res. Lett.* 9, 231-234.
- Zika, R. G., Saltzman, E. S., Chameides, W. L. and Davis, D. D. (1982a). H_2O_2 levels in rainwater collected in south Florida and the Bahama Islands. *J. Geophys. Res.* 87, C7, 5015-5017.

Zika, R. G., Moffett, J. W., Pestane, R. G., Cooper, W. J. and Saltzman, E. S. (1985a). Spatial and temporal variations of hydrogen peroxide in Gulf of Mexico waters. *Geochimica et Cosmochimica Acta* 49, 1173-1184.

Zika, R. G., Saltzman, E. S. and Cooper, W. J. (1985b). Hydrogen peroxide concentrations in the Peru upwelling area. *Marine Chemistry* 17, 265-275.

MEMBERS OF THE TASK FORCE

Ch. DE ROOIJ (Chairman)

SOLVAY

B - Brussel

J. BAKES

ATOCHEM

F - Paris

A. BALAY

CHEMOXAL

F - Chalon/Saone

W. J. BROCK

DUPONT

USA - Wilmington

I. DALIN

EKA NOBEL

S - Bohus

G. MALINVERNO

MONTEDISON

I - Milano

A. MAYR

DEGUSSA

D - Hanau

J. NAGAI

MITSUBISHI GAS CHEMICAL

J - Niigata

J. -F. REGNIER

ATOCHEM

F - Paris

M. WEINER (Consultant)

FMC CORPORATION

USA - Princeton

H. VRIJHOF (Secretary)

ECETOC

B - Brussel

MEMBERS OF THE ECETOC SCIENTIFIC COMMITTEE (Peer Review Committee)

W. F. TORDOIR (Chairman), Head of Occupational Health and Toxicology Division	SHELL NL - Den Haag
H. VERSCHUUREN, (Vice-Chairman) Head of Toxicology Department	DOW CHEMICAL CH - Horgen
O. C. BØCKMAN, Scientific Advisor†	NORSK HYDRO N - Porsgrunn
N. G. CARMICHAEL, Toxicology Director Worldwide	RHÔNE-POULENC F - Lyon
H. DE HENAU, European Technical Centre Professional and Regulatory Services	PROCTER & GAMBLE B - Brussels
A. DE MORSIER, Head, Ecotoxicology	CIBA-GEIGY CH - Basel
P. A. GILBERT, Head, Environmental Relations	UNILEVER UK - Port Sunlight
I. J. GRAHAM-BRYCE, Head of Environmental Affairs	SHELL NL - Den Haag
B. HILDEBRAND, Director, Experimental Toxicology	BASF AG D - Ludwigshafen
J. R. JACKSON, Director Medicine and Health Science	MONSANTO EUROPE B - Brussels
K. KÜNSTLER, Head Biological Research	HENKEL D - Duesseldorf
H. LAGAST, Chief Medical Officer†	SOLVAY B - Brussels
E. LÖSER, Head of Institute of Industrial Toxicology	BAYER D - Wuppertal
R. MILLISCHER, Chief Toxicologist†	ATOCEM F - Paris
I. F. H PURCHASE, Director, Central Toxicology Laboratory	ICI UK - Macclesfield

† Stewards - responsible for primary peer review

Joint Assessment of Commodity Chemicals No. 22, Hydrogen Peroxide

Errata (4 October 1994)

- Page 7, Table 1: 45% should read 35%, density should read 1,133.
Density: decimal points should be read as comma's.
- Page 16, Section 4.1.1 Surface water. The second and third reaction should be equated electrically: $\text{Org}^{\bullet} + \text{O}_2 \rightarrow \text{Org}^{\bullet+} + \cdot\text{O}_2^-$
- $$2 \cdot\text{O}_2^- + 2\text{H}^+ \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$$
- In seawater, similarly, the reactions should read:
- $$\text{Fe}^{2+} + \text{O}_2 \rightarrow \text{Fe}^{3+} + \cdot\text{O}_2^-$$
- $$\text{Fe}^{2+} + \cdot\text{O}_2^- + 2\text{H}^+ \rightarrow \text{Fe}^{3+} + \cdot\text{O}_2$$
- Page 36, Table 11: Concentrations measured by Kaelin *et al* without ventilation other than the windows were 41 mg/m³ on the floor and 12 mg/m³ close to the machine; with new ventilation the respective values were 4.5 mg/m³ and 1.5 mg/m³.
- Page 40, Table 13: µg/l should read mg/l
- Page 33, Section 5.2.1: The second sentence should be replaced by:
- "It is generally only permitted ... chemical means. The residual level in food immediately after aseptic packaging should not exceed 0.5 ppm (US FDA, 1990). Under the GRAS regulation, the maximum treatment levels for H₂O₂ in food applications can be up to 1.25% depending on the use (US FDA, 1992)."
- References:
- US FDA (US Food and Drug Administration) (1992). Maximum usage levels permitted [in food]. CFR 21, 184.1366c.
- US FDA (US Food and Drug Administration) (1990). Substances utilized to control the growth of microorganisms. CFR 21, 178.1005d [unchanged in 1993 edition].
- Page 54/55, Table 20 Please insert the attached table.
- Page 69, Gavage: calculated, based on 0.5% should read 5% solution
- Page 140: ATOCHEM should read ELF ATOCHEM and A. MAYR should read W. MAYR.
- Encl. Table 20

Table 20: Enzymatic Activity in Organs and Tissues of Rats (Matkovichs and Novak, 1977)

Organ or tissue	Enzymatic activity (U/g wet weight)					
	Superoxide dismutase	Peroxidase	Controls	H ₂ O ₂ -treated	Catalase	H ₂ O ₂ -treated
	Controls	H ₂ O ₂ -treated			Controls	
Organs						
Liver	4,000 ± 600	10,408 ± 750	0	0	4.80 ± 0.43	25,250 ± 2,455
Kidney	1,120 ± 151	1,692 ± 240	33	105	0.36 ± 0.03	3,300 ± 0.295
Spleen	560 ± 50	470 ± 240	91	418	2.40 ± 0.21	0.730 ± 0.062
Testes	960 ± 63	1,100 ± 53	41	-	0.41 ± 0.03	0.150 ± 0.011
Whole brain	240 ± 24	397 ± 38	120	13	0.04 ± 0.004	0.012 ± 0.001
Lung	210 ± 20	895 ± 85	86	201	0.26 ± 0.02	0.520 ± 0.050
Pancreas	310 ± 31	352 ± 33	13	46	0.195 ± 0.010	0.255 ± 0.023
Muscles						
Heart	480 ± 47	950 ± 93	300	155	0.245 ± 0.023	0.725 ± 0.065
Skeletal	300 ± 27	530 ± 50	10	69	0.11 ± 0.009	0.092 ± 0.009
Haemolysate*	696 ± 68	767 ± 80	11,666 ± 1,000	17,900 ± 1,340	4.12 ± 0.04	6.98 ± 0.07

* U/ml wet weight; n = 10; catalase activity-values in U/g wet weight