


## 6. Safety

Hydrogen peroxide has a high energy content:



Furthermore, its high oxygen content (47 wt%) is available for combustion reactions and the oxidation of organic compounds. This means that special safety conditions apply in handling hydrogen peroxide. Safety studies can be divided into three groups:

- 1) examination of the pure vapor phase,
- 2) examination of the pure liquid (aqueous) phase, and
- 3) examination of mixtures or solutions of organic compounds with or in aqueous hydrogen peroxide.

Results are summarized below, but the literature must be consulted for details. In specific cases, especially when hydrogen peroxide is used with organic compounds, the intended work area must be checked for hazards if it has not already been certified as safe by a safety examination.

**Pure Hydrogen Peroxide—Vapor Phase** [5], [116]–[119]. Explosive vapor mixtures are formed at atmospheric pressure when the hydrogen peroxide concentration in the vapor phase exceeds 26 mol%. Figure 16 shows the explosive




Figure 16. Dependence of the explosive range of the vapor phase (> 26 mol%  $\text{H}_2\text{O}_2$ ) on the temperature and the hydrogen peroxide concentration of liquid phase (at atmospheric pressure)




Figure 17. Dependence of the explosive range of vapor phase on pressure

range as a function of temperature and hydrogen peroxide concentration.

For the industrial concentration of hydrogen peroxide solutions, it is important to note that at reduced pressure the critical hydrogen peroxide concentration of the vapor phase may exceed 26 mol% (Fig. 17).

**Pure Hydrogen Peroxide—Liquid (Aqueous) Phase** [120], [121]. The explosive risk of pure, highly concentrated hydrogen peroxide solutions has been examined by various methods, but the sometimes contradictory findings are difficult to interpret consistently.

Results are influenced primarily by test conditions and not by hydrogen peroxide concentration. For example, 90.7 wt% hydrogen peroxide in a tube with an internal diameter of 2.67 cm (1.05 inches) could not be exploded by the detonation shock of a tetryl booster, whereas explosion occurred in a tube with an internal diameter of 4.09 cm (1.61 inches). Similarly, 86% hydrogen peroxide in a tube of 4.09 cm internal diameter was exploded by raising the




Figure 18. Explosive range (hatched area) of hydrogen peroxide-acetone-water mixtures  
Dashed line, indicates the stoichiometry  
 $8 \text{H}_2\text{O}_2 + \text{CH}_3\text{COCH}_3 \rightarrow 3 \text{CO}_2 + 11 \text{H}_2\text{O}$

temperature above  $50^\circ\text{C}$  [120], [121]. The decisive factors for explosion of pure aqueous hydrogen peroxide seem to be the degree of occlusion, the insulation, and the energy of detonation.

#### Mixtures or Solutions of Aqueous Hydrogen Peroxide with Organic Compounds [122]–[124].

The hazards associated with mixtures of hydrogen peroxide and organic chemicals are important for its industrial production and, especially, for its uses. As an example, Figure 18 shows results obtained with the hydrogen peroxide-acetone-water system [122]. Many other organic compounds give similar results. The size of the explosive region depends on the organic compound and the test conditions.

For practical use, safety tests of the intended working area are very important. When systems such as hydrogen peroxide-formic acid-water are used, other products may be formed (in this case performic acid), that are more dangerous than the original mixture [123].

### 7. Uses

Development of the AO process permitted large-scale production of hydrogen peroxide, which is now used widely in almost all industrial areas. Its main use is in bleaching ( $\rightarrow$  Bleaching, A4, p. 194). Uses in the chemical industry and in environmental protection are increasing because its great advantage is that the degradation product is water.

Hydrogen peroxide is used in the textile industry for bleaching cotton, linen, bast fibers, wool, silk, polyester fiber, and polyurethane fiber. In the pulp and paper industry, it is used to bleach sulfate and sulfite cellulose, wood pulp, and wastepaper, and to brighten wood veneers and wooden structures.

The chemical industry employs hydrogen peroxide for the production of peroxy compounds such as sodium perborate, sodium percarbonate, metallic peroxides, or percarboxylic acids. Hydrogen peroxide is very important in organic chemistry for epoxidation and hydroxylation (manufacture of plasticizers and stabilizers for the plastics industry), oxidation (manufacture of amine oxides as washing-up liquids), oxidation, and initiation of polymerization.

Sulfuric acid solutions of hydrogen peroxide are used for the pickling and chemical polishing of copper, brass, and other copper alloys, as well as for etching and cleaning printed circuit boards. Highly purified hydrogen peroxide is used in the manufacture of silicon semiconductor chips to clean silicon disks and to remove photoresist layers. It is also used for in situ leaching in underground uranium mining.

Hydrogen peroxide is increasingly used in environmental protection to detoxify effluents containing formaldehyde, phenols, or cyanide (e.g., wastewater from mines and tempering works, galvanizer concentrate, photochemical effluents), and to deodorize sulfur-containing effluents. Smoke and exhaust gases containing sulfur dioxide can be completely detoxified with hydrogen peroxide.

Hydrogen peroxide is a highly efficient disinfectant (especially for packaging materials); it is also used as a bleaching agent in hair preparations and as a propellant in space technology.

### 8. Toxicology and Occupational Health

**Toxicity in Humans.** In humans, brief contact of hydrogen peroxide with the skin leads to irritation and whitening (cutaneous emphysema), the severity of which depends on concentration. Longer contact or higher concentration can lead to burns.

Contact with the eyes also leads to serious injury. Hydrogen peroxide vapor or aerosol causes irritation or damage of the upper respira-

tory tract and serious lung injuries [125], [126]. The human reaction to the irritating effect of hydrogen peroxide on the mucous membranes and skin is far more sensitive than that of the rat. The threshold concentration for acute irritative effects of gaseous hydrogen peroxide on the respiratory tract is 60 mg/m<sup>3</sup> in rats but only 10 mg/m<sup>3</sup> in humans; the corresponding values for skin are 110 mg/m<sup>3</sup> for rats and 20 mg/m<sup>3</sup> for humans [127].

Accidental ingestion of hydrogen peroxide (33 wt% solution) in five cases led to stomach and chest pain, respiratory depression, foaming at the mouth, and loss of consciousness. Muscle and nerve disturbances and fever were observed as added complications. However, all affected persons recovered after several weeks [128].

Hydrogen peroxide has not been found to produce teratogenic or carcinogenic effects in humans; mutagenic or chromosomal effects have not been observed. The MAK and TLV-TWA values (1987) are 1 ppm (1.4 mg/m<sup>3</sup>).

#### Toxicity in Animals. Acute toxicities of hydrogen peroxide in animals follow [128]:

LD <sub>50</sub> (mouse, oral)	2538 mg/kg
LD <sub>50</sub> (mouse, oral)	2000 mg/kg
LD <sub>50</sub> (rat, oral)	4060 mg/kg
LD <sub>50</sub> (mouse, dermal)	1.2 × 10 <sup>4</sup> mg/kg
LC <sub>50</sub> (rat, inhalation, 4 h)	2000 mg/m <sup>3</sup>

Acute dermal toxicity depends on hydrogen peroxide concentration. With 90 wt% hydrogen peroxide, the dermal LD<sub>50</sub> in rabbit is 630 mg/kg and in rat 4800 mg/kg [129]. So far no indication of a toxic effect on reproduction in animals has been found [130].

Hydrogen peroxide has a mutagenic effect on fungi and bacteria (e.g., Ames test), but not on insects or mammalian cells in vitro [130].

Oral administration of hydrogen peroxide produced tumors in the small intestine of mice [131], [132]. According to the criteria of the International Agency for Research on Cancer (IARC), however, only "limited evidence of the carcinogenicity of hydrogen peroxide to experimental animals" exists [130]. When hydrogen peroxide was given orally to rats, no significant differences occurred between the two test groups and the control group [133]. No evidence of tumor formation was found one year after exposure of mouse skin to 5% hydrogen peroxide [134].

## 9. References

### General References

- [1] *Chemlin*, System no. 3, Sauerstoff.
- [2] *Kirk-Othmer*, 3rd ed., 13, 12, 38.
- [3] W. Machu: *Das Wasserstoffperoxid und die Verbindungen*, 2nd ed., Springer, Wien 1951.
- [4] *Hydrogen Peroxide*, Mannl, Chem. Assoc. Chem. Sal., Data Sheet SD-53 (1969).
- [5] W. C. Schumb, C. N. Sauterfield, R. L. Wennerth: *Hydrogen Peroxide*, Reinhold Publ. Co., New York 1955.
- [6] *Elinum*, 4th ed., 17, 694.
- [7] W. M. Weigert: *Wasserstoffperoxid und seine Derivate*, Hüthig, Heidelberg 1978.
- [8] *Winnacker-Kitchler*, 4th ed., *Anorganische Technologie* 1, 2, 563.

### Specific References

- [9] L. J. Thernard, *Ann. Chim. Phys.* 8 (1818) 306.
- [10] H. Medinger, *Ann. Chem. Pharm.* 88 (1853) 57.
- [11] H. Berthelot, C. R. *Hebd. Seances Acad. Sci.* 86 (1878) 71.
- [12] K. Elbs, O. Schönher, *Z. Elektrochem.* 1 (1895) 417, 468.
- [13] W. M. Weigert, H. Delle, G. Kätsch, *Chem. Ztg.* 99 (1975) 101.
- [14] W. Marchot, *Liebigs Ann. Chem.* 314 (1901) 177, 316 (1901) 318, 331.
- [15] J. H. Walton, G. W. Flisun, *J. Am. Chem. Soc.* 54 (1932) 3228.
- [16] O. v. Schösch, *Chem. Ing. Tech.* 32 (1960) 462.
- [17] J. Rivoire, *Chem. Week* 140 (1987) Sept. 16, 38.
- [18] W. R. Busch, H. A. Levy, *J. Chem. Phys.* 42 (1965) 3054.
- [19] G. Satchard, G. M. Karanagh, L. B. Ticknor, *J. Am. Chem. Soc.* 74 (1952) 3715.
- [20] W. M. Latimer: *The Oxidation States of the Elements and their Potentials in Aqueous Solutions*, 2nd ed., Prentice Hall, New York 1952.
- [21] Degussa, DE 1 039 500, 1956 (G. Kätsch, E. Richter, A. Gross).
- [22] I. G. Farbenindustrie AG, GB 508 081, 1939.
- [23] Degussa, DE-AS 1945750, 1969 (G. Kätsch, R. Trübe).
- [24] EDOGAWA, DE 1112051, 1957 (K. Hiratsuka, S. Matsumura, M. Matsuda, K. Takagi, T. Abe).
- [25] Laporte Chemicals Ltd., DE 953 790, 1953 (C. W. Lefevre).
- [26] Degussa, DE 1261 838, 1963 (G. Kätsch).
- [27] E. I. du Pont de Nemours & Co., DE 888 840, 1950 (C. R. Harris, J. W. Sprau).
- [28] BECCO, US 2537 655, 1950 (L. H. Dawsey, C. K. Muehlhauser, R. R. Umhoefer).
- [29] BECCO, US 2537 516, 1950 (L. H. Dawsey, C. K. Muehlhauser, R. R. Umhoefer).
- [30] Laporte Chemicals Ltd., DE 933 088, 1953 (C. W. Lefevre).
- [31] E. I. du Pont de Nemours & Co., BE 819 676, 1975 (L. G. Vaughan).
- [32] Degussa, DE 2018 686, 1970 (G. Gisselmann, G. Schreyer, W. Weigert).
- [33] BMC Corp., US 4394 369, 1983 (R. Wayne).
- [34] I. G. Farbenindustrie AG, US 2158 525, 1936 (H. I. Riedl, G. Pleiderer).

- [35] E. I. du Pont de Nemours & Co., US 2,689,169, 1949 (W. S. Hinegardner).
- [36] Degussa, DE 1,063,581, 1956 (L. R. Darbee, D. F. Kreuz).
- [37] EDOGAWA, DE-AS 1,085,861, 1957 (K. Hiratsuka et al.).
- [38] Laporte Chemicals Ltd., GB 741,444, 1953 (W. R. Holmes, C. W. Lefevre, R. Lait, W. S. Wood).
- [39] Solvay & Cie., DE 1,041,928, 1955 (M. Glonze, E. Lehlion).
- [40] Degussa, DE 1,051,257, 1957 (G. Kabisch, E. Richter).
- [41] E. I. du Pont de Nemours & Co., US 2,673,140, 1951 (J. W. Sprauer).
- [42] EDOGAWA, FR 1,319,025, 1962.
- [43] EDOGAWA, DE-OS 2,141,234, 1971 (H. Shin et al.).
- [44] Degussa, DE 1,273,499, 1964 (G. Schreyer, O. Weiberg, G. Giesseimann).
- [45] BASF, DE 801,840, 1951 (G. Pfeiderer, H.-J. Riedl, W. Deuschel).
- [46] FMC Corp., US 2,720,532, 1952 (R. R. Umhoefer).
- [47] FMC Corp., US 2,730,533, 1952 (R. R. Umhoefer).
- [48] Kali-Chemie, DE-AS 1,137,719, 1960 (V. Hauschild, M. Stahl).
- [49] Degussa, DE 1,542,089, 1965 (G. Kabisch, H. Herzog).
- [50] Degussa, GB 1,096,260, 1965.
- [51] Laporte Chemicals Ltd., GB 718,306, 1953 (C. A. Morgan).
- [52] Laporte Chemicals Ltd., GB 718,307, 1952 (J. A. Williams, C. W. Lefevre).
- [53] FMC Corp., US 3,009,782, 1958 (D. H. Porter).
- [54] Degussa, DE 1,072,228, 1959 (L. R. Darbee, D. F. Kreuz).
- [55] Degussa, DE 1,079,604, 1959 (D. H. Porter).
- [56] Degussa, DE 2,003,268, 1970 (M. Liebert, H. Delle, G. Kabisch).
- [57] *Chem. Age (London)* 82 (1958) 895.
- [58] *Chem. Ind. (London)* (1959) 76.
- [59] *Chem. Process Eng. (London)* 40 (1959) no. 1, 5.
- [60] *Br. Chem. Eng.* 4 (1959) 88.
- [61] *Ind. Chem.* 35 (1959) 9.
- [62] Allied Chemical Corp., US 2,902,347, 1954 (J. N. Cosby, R. L. Harris, E. H. Rissmiller).
- [63] Pittsburgh Plate Glass Co., DE 977,130, 1965 (J. R. Cox, T. E. Brown).
- [64] Pittsburgh Plate Glass Co., US 3,055,838, 1957 (J. W. Moore).
- [65] E. I. du Pont de Nemours & Co., US 2,739,875, 1951 (J. W. Sprauer, T. V. Williams).
- [66] Laporte Chemicals Ltd., GB 928,784, 1959 (B. K. Howe).
- [67] Laporte Chemicals Ltd., DE-AS 1,138,743, 1959 (B. K. Howe, S. Andrus, G. N. Davis).
- [68] E. I. du Pont de Nemours & Co., CA 649,850, 1961 (W. S. Hinegardner, H. R. Eisenhauer).
- [69] Pittsburgh Plate Glass Co., FR 1,342,577, 1962 (M. M. Korach).
- [70] Laporte Chemicals Ltd., GB 991,413, 1961 (B. K. Howe).
- [71] Columbia-Southern Chemical Corp., GB 795,433, 1955 (J. R. Cox, M. Korach).
- [72] Pittsburgh Plate Glass Co., GB 924,625, 1958 (M. Q. Freeland, P. R. Johnston).
- [73] E. I. du Pont de Nemours & Co., DE-AS 1,036,225, 1956 (J. W. Sprauer).
- [74] Laporte Chemicals Ltd., GB 841,323, 1956 (H. B. Hopkins).
- [75] FMC Corp., US 3,043,666, 1960 (G. W. Swinski).
- [76] Laporte Chemicals Ltd., DE-AS 1,467,091, 1964 (S. D. Williams).
- [77] E. I. du Pont de Nemours & Co., US 3,016,290, 1956 (A. A. Elston, W. S. Hinegardner).
- [78] Oxyntbise, FR 1,539,843, 1967 (P. Thirion).
- [79] Laporte Chemicals Ltd., US 3,073,755, 1963 (R. H. Banfield, A. M. Hildon).
- [80] Columbia-Southern Chemical Corp., US 2,960,448, 1960 (W. S. Straub, C. J. Sindlinger).
- [81] Degussa, US 3,060,405, 1962 (J. Müller).
- [82] FMC Corp., US 3,152,052, 1964 (T. M. Jenney, D. H. Porter, R. H. Rice).
- [83] E. I. du Pont de Nemours & Co., US 2,479,111, 1949 (C. R. Harris).
- [84] Shell Development Co., US 2,871,104, 1951 (E. F. Rust).
- [85] N. V. De Bataafse Petroleum Maatschappij, GB 708,339, 1954.
- [86] A. E. Brodskii, V. I. Franchuk, M. M. Alexankin, V. A. Lunok-Burmakina, *Dokl. Akad. Nauk SSSR* 123 (1953) no. 1, 117.
- [87] N. V. De Bataafse Petroleum Maatschappij, BE 1,002,295, 1953 (J. R. Skinner, S. E. Steimle).
- [88] *Chem. Eng. News* 35 (1957) no. 48, 26.
- [89] *Chem. Week* 125 (1979) no. 18, 13.
- [90] E. I. Preva, *Chimicheskaja Promyshlennost* 9 (1970) 19.
- [91] P. Askenasy, R. Rose, *Z. Anorg. Allg. Chem.* 189 (1930) 1, 10.
- [92] J. Lawrence, R. B. Pascock, C. W. Lefevre, *Bios Final Report* 1432, item No. 22 (1946).
- [93] D. I. Campbell, E. F. Lefson, US 2,497,812, 1950.
- [94] Atlantic Richfield Co., NL 7407,533, 1974.
- [95] Sun Oil Co., DE-OS 2,100,784, 1971 (J. O. Turner).
- [96] Wacker-Chemie, US 3,124,211, 1964 (W. Lohringer, J. Sixt).
- [97] Celanese Corp., US 3,341,297, 1967 (A. F. MacLean, A. L. Stautzenberger).
- [98] Elektrochem. Werke München, US 2,015,040, 1935 (A. Pretsch).
- [99] Imperial Chem. Ind. Ltd., GB 1,056,126, 1967 (G. W. Hooper).
- [100] Air Products & Chem. Inc., DE-OS 2,710,279, 1977 (P. M. Dyer, F. Mosley).
- [101] Shell Oil Co., US 4,007,236, 1977 (L. Kim, G. W. Schoenthal).
- [102] FMC Corp., US 4,347,231, 1982 (R. C. Michaelson).
- [103] Kymene AB, DE 1,054,970, 1954 (R. M. R. Örnheim).
- [104] Kymene AB, DE 1,036,829, 1955 (R. M. R. Örnheim).
- [105] Kymene AB, DE 1,063,131, 1956 (R. M. Rafaelson).
- [106] M. Traube, *Ber. Dtsch. Chem. Ges.* 15 (1882) 2434.
- [107] F. Fischer, O. Priess, *Ber. Dtsch. Chem. Ges.* 46 (1913) 698.
- [108] E. Berl, *Trans. Electrochem. Soc.* 76 (1939) 359.
- [109] F. Berl, DE 648,964, 1928.
- [110] B. Kastning, W. Faul, *Chem. Ing. Tech.* 49 (1977) no. 11, 911.
- [111] Kernforschungsanlage Jülich, DE-OS 2,501,342, 1976 (W. Faul, B. Kastning).
- [112] Laporte Chemicals Ltd., GB 893,069, 1959 (P. V. Rutledge, D. S. Bell).

- [113] Columbia-Southern Chemical Corp., US 2904 517, 1959 (P. H. Baker).
- [114] Shell Oil Company, US 3035 634, 1962 (D. B. Luten, R. E. Meeker).
- [115] Shell Oil Company, US 3053 632, 1962 (R. F. Meeker).
- [116] C. N. Satterfield, G. M. Kavanaugh, H. Resnick, *Ind. Eng. Chem.* 43 (1951) no. 11, 2507.
- [117] I. M. Monger, H. J. Baumgartner, G. C. Hood, C. E. Sanborn, *Journ. Chem. Eng. Data* 9 (1964) no. 1, 119.
- [118] E. S. Shandley, P. F. Strong, L. R. Darbee, N. D. Lee, *Ind. Eng. Chem. Fundam.* 9 (1970) no. 4, 574.
- [119] C. N. Satterfield, P. I. Cocconi, A. H. R. Feldbrugge, *Ind. Eng. Chem.* 47 (1955) 1040.
- [120] *Bur. Mines Inf., Chm.* 8387 (1968).
- [121] E. S. Hauser, *Explosionsstoffe* 617 (1953) 64.
- [122] E. S. Shanley, F. P. Greenspan, *Ind. Eng. Chem.* 39 (1947) no. 12, 1536.
- [123] I. M. Monger, H. Selbo, D. C. Lehwald, *J. Chem. Eng. Data* 6 (1961) no. 1, 23.
- [124] *Bur. Mines Rep. Invest.* 5817 (1961).
- [125] Merkblatt der Berufsgenossenschaft der chemischen Industrie, *Wasserstoffperoxid*, 11009, 7/84, FRG.
- [126] F. Haenschel, R. Ludewig, H. Mühlberg, "Corrosive action of hydrogen peroxide," *Nahrung-Schmidteberg's Arch. Exp. Pathol. Pharmacol.* 235 (1958) 51 62.
- [127] M. T. Budagovskiy, M. K. Vodaekhoriyu, A. I. Desyatov, "Poisoning with hydrogen peroxide," *oravi, peritidolem voen. Med. Zh.* 9 (1971) 79-81.
- [128] P. P. Ljarskij et al., "Toxikologisch-Hygienische Charakteristik von Desinfektionsmitteln auf der Basis von Wasserstoffperoxid und seinen Derivaten," *Gig. Sanit.* 1983 no. 5, 28-31.
- [129] Food and Drug Administration, *Evaluation of the Health Aspects of Hydrogen Peroxide as a Food Ingredient*, U.S. Department of Commerce, National Technical Information Service, PB 80-104 607, 1979.
- [130] *IARC Monogr. Eval. Carcinog. Risk Chem. Mon* 36 (1985) 285-314.
- [131] A. Ito, H. Watanabe, M. Naito, Y. Naito, "Induction of duodenal tumors in mice by oral administration of hydrogen peroxide," *Gann.* 72 (1981) 174 175.
- [132] A. Ito et al., "Correlation between induction of duodenal tumor by hydrogen peroxide and catalase activity in mice," *Gann.* 75 (1984) 17-21.
- [133] T. Takayama, S. Takizawa, "Carcinogenicity study of hydrogen peroxide in rats," *Annual Report of Cancer Research*, Ministry of Health and Welfare, Japan 1980.
- [134] Y. Kurokawa et al., "Studies on the promoting and complete carcinogenic activities of some oxidizing chemicals in skin carcinogenesis," *Cancer Lett.* 24 (1984) 299-304.

#### Hydrogen Selenide → Selenium and Selenium Compounds Hydrogen Storage Alloys → Hydrides

