

5) Hileman, R. E. and J. D. Astwood (2001). Additional characterization of the Cry3Bb1 protein produced in corn event MON 863. MSL-17137, an unpublished study conducted by Monsanto Company.

Study Title

Additional Characterization of the Cry3Bb1 Protein
Produced in Corn Event MON 863

Authors

Ronald E. Hileman, Ph.D. and James D. Astwood, Ph.D.

Study Completed On

April 13, 2001

Performing Laboratory

Monsanto Company
Product Safety Center
Biotechnology Regulatory Sciences
700 Chesterfield Parkway North
St. Louis, MO 63198

Laboratory Project ID

MSL Number: 17137



AA050317

©01 Proprietary Information of Monsanto Company

[REDACTED]

Information claimed confidential on the basis of its falling within the scope of FIFRA section 10(d)(1)(A), (B) or (C) has been removed to a confidential appendix, and is cited by cross reference number in the body of the report.

"We submit this material to the United States Environmental Protection Agency specifically under the requirements set forth in FIFRA as amended, and consent to the use and disclosure of this material by EPA strictly in accordance with FIFRA. By submitting this material to EPA in accordance with the method and format requirements contained in PR Notice 86-5, we reserve and do not waive any rights involving this material that are or can be claimed by the company notwithstanding this submission to EPA."

Company: Monsanto Company

Company Agent: Dennis P. Ward

Title: Regulatory Affairs Manager

Signature: [Signature] Date: 7 May 2001

Statement of Compliance

The research described in this report was not conducted under the Good Laboratory Practice standards as specified in 40 CFR Part 160.



Submitter

Date: 8 May 2001

Ravinder S Sidhu

Sponsor

Date: April 13, 2001



Director, Product Safety Center

Date: April 13, 2001

RW E. H. H. H.

Author

Date: 13 APR 2001

Summary of Quality Control Review

This report was reviewed to ensure that it accurately reflects the supporting documentation. The supporting documentation was audited for compliance to the Monsanto Company Guidelines for Keeping Research Records (GRR 10/1/99), and where applicable, to Monsanto SOP's.

Joan M. Rejda-Heath
Joan M. Rejda-Heath, Ph.D.
Quality Assurance Specialist
Monsanto Regulatory
Monsanto Company

Date: Apr 13, 2001

MSL Number: 17137

Title: Additional Characterization of the Cry3Bb1 Protein Produced in Corn Event MON 863

Facility: Monsanto Company
Product Safety Center
Biotechnology Regulatory Sciences
700 Chesterfield Parkway North
Chesterfield, Missouri 63198

Sponsor: Ravinder S. Sidhu

Authors: Ronald E. Hileman
James D. Astwood

Report Completion Date: April 13, 2001

Records Retention: All report related supporting documentation and final report will be retained at Monsanto-St. Louis.

Signatures of Final Report Approval:

R. E. Hileman
Author

Date: 13 APR 2001

J. D. Astwood
Director, Product Safety Center

Date: April 13, 2001

Ravinder S. Sidhu
Sponsor

Date: April 13, 2001

Table of Contents

| Section | Page |
|--|------|
| Title Page..... | 1 |
| Statement of Data Confidentiality Claims..... | 2 |
| Statement of Compliance | 3 |
| Summary of Quality Control Review..... | 4 |
| Signatures of Final Report Approval..... | 5 |
| Table of Contents | 6 |
| Abbreviations and Definitions | 8 |
| 1.0 Summary | 9 |
| 2.0 Background | 10 |
| 3.0 Molecular Characterization of the <i>cry3Bb1</i> Insert in Corn Event MON 863..... | 11 |
| 4.0 Characterization of the Cry3Bb1 Protein Produced in Corn Event MON 863 | 12 |
| 5.0 Conclusions | 12 |
| 6.0 References | 13 |

Tables

| | | |
|----------|--|----|
| Table 1. | Amino Acid Diversity of Cry3Bb1 Variants..... | 10 |
| Table 2. | Specific nucleotide modifications and resulting amino acid changes used to produce the Cry3Bb1.11098 and Cry3Bb1.11231 proteins in <i>Bacillus thuringiensis</i> | 15 |
| Table 3. | Summary of the amino acid differences among the Cry3Bb1 protein variants produced in corn..... | 15 |

Figures

| | | |
|-----------|---|----|
| Figure 1. | Summary of the origin of the Cry3Bb1 variant proteins present in corn events described in food, feed and environmental evaluations..... | 16 |
| Figure 2. | Sequences of the Cry3Bb1 protein variants | 17 |

Abbreviations and Definitions

| | |
|----------------|---|
| aa | amino acid |
| <i>B.t.</i> | <i>Bacillus thuringiensis</i> |
| Cry3Bb1 | <i>Bacillus thuringiensis</i> derived insecticidal protein |
| <i>cry3Bb1</i> | DNA that encodes the Cry3Bb1 protein |
| IUPAC-IUB | International Union of Pure and Applied Chemistry - International Union of Biochemistry |
| kb | kilobase |
| MALDI-TOF | Matrix assisted laser desorption ionization time of flight |
| <i>npt II</i> | DNA that encodes for the neomycin phosphotransferase II (NPT II) protein |

1.0 Summary

Corn plants protected from corn rootworm feeding damage were designed by insertion of a gene encoding a modified *Bacillus thuringiensis* Cry3Bb1 protein into the corn genome. Corn event MON 863, which produces the Cry3Bb1 protein referred to as Cry3Bb1.11098, is the focus of this report. Other corn events producing the Cry3Bb1.11098 protein as well as a separately modified Cry3Bb1 protein have been previously reported and are also described for completeness.

Initial modifications of the wild type *Bacillus thuringiensis* (*B.t.*) *cry3Bb1* gene (Donovan *et al.*, 1992) were performed using a *B.t.* cloning vector and expression system (English *et al.*, 2000). Bioactivity analyses were used to select variant *cry3Bb1* coding sequences that produced Cry3Bb1 proteins in *B.t.* with increased activity toward the larvae of a target insect, Southern corn rootworm. The modified *cry3Bb1* coding sequence contained in *B.t.* strain EG11098 (producing the Cry3Bb1.11098 protein) was selected for corn transformation as part of this extensive process.

The Cry3Bb1.11098 protein produced in *B.t.* differs from the wild type Cry3Bb1 protein by 5 amino acids. When this *B.t.* gene was modified for expression in corn, the resulting gene was predicted to encode an additional two amino acid changes relative to the *B.t.* produced Cry3Bb1.11098 protein sequence. Thus the Cry3Bb1.11098 protein produced in corn event MON 863 differs from the *B.t.* produced Cry3Bb1.11098 protein by 2 amino acids, and 7 amino acids relative to the wild type Cry3Bb1 protein.

Additional characterization of the DNA sequence of the *cry3Bb1* coding region in corn event MON 863 and the Cry3Bb1 protein produced in corn event MON 863 confirmed these described amino acid differences relative to the wild type Cry3Bb1 protein sequence deduced from the wild type *cry3Bb1* gene. The exact DNA and encoded protein sequence of this specific Cry3Bb1 allele is described in this report. Specific amino acid substitutions are shown in Table 1.

Table 1. Amino Acid Diversity of Cry3Bb1 Variants.

| Cry3Bb1 Allele | Amino acid substitutions relative to wild type Cry3Bb1 protein ^a | | | | | |
|---|--|-------|-------|-------|-------|-------------|
| <i>B.t.</i> strain EG11098 Cry3Bb1.11098 protein | D165G | H231R | S311L | N313T | E317K | |
| <i>B.t.</i> strain EG11231Cry3Bb1.11231 protein | | H231R | S311L | N313T | E317K | |
| Corn event MON 863 Cry3Bb1.11098 protein ^b | A2 | D166G | H232R | S312L | N314T | E318K Q349R |
| Corn event MON 853 Cry3Bb1.11231 protein ^b | A2 | | H232R | S312L | N314T | E318K |

^a wild type Cry3Bb1 protein corresponds to GenBank Accession No. M89794.

^b Amino acid positions are shifted upwards by one amino acid relative to the *B.t.* proteins due to the introduction of an alanine at position 2.

2.0 Background

A modified gene encoding a variant of the *Bacillus thuringiensis* (*B.t.*) Cry3Bb1 protein was used to design corn event MON 863. A separate but similar *cry3Bb1* gene variant was used to design corn event MON 853 (Cavato *et al.*, 1999). Although the Cry3Bb1 protein produced in corn event MON 853 is not the focus of this report, DNA and protein sequences are provided for completeness because initial food, feed and environmental evaluations were performed using this protein variant. Plants producing these Cry3Bb1 protein variants are resistant to feeding damage from the coleopteran insect, corn rootworm.

The wild type *cry3Bb1* gene (GenBank Accession No. M89794) was modified initially utilizing a *B.t.* plasmid and expression system (English *et al.*, 2000). Proteins encoded by two modified *cry3Bb1* genes were observed to have increased bioactivity towards corn rootworm when expressed in *B.t.* These were the *cry3Bb1.11098* and *cry3Bb1.11231* genes, contained in *B.t.* strains EG11098 and EG11231 respectively, which produced the corresponding Cry3Bb1.11098 and Cry3Bb1.11231 proteins. The specific modifications that yielded these Cry3Bb1 protein variants are described in Table 1. The Cry3Bb1.11098 protein produced in *B.t.* strain EG11098 differs from the wild type Cry3Bb1 protein by 5 amino acids. The Cry3Bb1.11231 protein produced in *B.t.* strain EG11231 differs from the wild type Cry3Bb1 protein by 4 amino acids.

DNA encoding these Cry3Bb1 protein variants were then modified to design genes compatible for cloning and expression in plants prior to transforming corn. Corn events that have been previously described are summarized in Figure 1. These modifications resulted in an additional change to the predicted amino acid sequences of these variant proteins. That is, three nucleotides (GCC for the *cry3Bb1.11098* allele or GCA for the *cry3Bb1.11231* allele, at positions 4, 5 and 6) were inserted at the 5' end of the coding sequence to create an *Nco* I restriction endonuclease site and facilitate cloning of the *cry3Bb1* gene into a plant transformation vector. This manipulation resulted in the introduction of an alanine (A) residue at position 2 of the predicted plant protein sequences.

In addition to the *cry3Bb1* gene, other genetic elements were included for gene expression and selection, but are not described in this report. For example, the neomycin phosphotransferase II (*nptII*) gene was used as the selectable marker (Cavato *et al.*, 2001), which enabled selection of cells in tissue culture containing the *cry3Bb1* gene.

Recent DNA sequencing of the *cry3Bb1* coding region and protein analyses of the Cry3Bb1 protein produced in corn event MON 863 have shown that the plant expressed 11098 gene encodes a Q349R substitution in the expected 11098 protein. The Cry3Bb1.11098 protein produced in *B.t.* strain EG11098 differs from the wild type Cry3Bb1 protein by 5 amino acids (Tables 2 and 3). When the coding region of this *B.t.* gene was modified for expression in corn, the resulting gene is predicted to encode an additional two changes relative to the *B.t.* strain EG11098 Cry3Bb1.11098 protein sequence. Thus, corn event MON 863 Cry3Bb1.11098 protein (as well as all other corn events producing the Cry3Bb1.11098 protein described in Figure 1) differs from the *B.t.* strain EG11098 Cry3Bb1.11098 protein by 2 amino acids, or 7 amino acids relative to the wild type Cry3Bb1 protein. The Q349R substitution is absent in corn events producing the Cry3Bb1.11231 protein.

3.0 Molecular Characterization of the *cry3Bb1* Insert in Corn Event MON 863

Additional molecular analyses have been recently performed on corn event MON 863 (Cavato and Lirette, 2001). Analysis of the insert confirmed the organization of the elements within the insert, as well as the complete DNA sequence of the insert in corn rootworm event MON 863 using PCR and DNA sequence analyses. The sequence of the 1.96 kb *cry3Bb1* DNA sequence and deduced 653 amino acid protein sequence is shown in Figure 2.

The deduced Cry3Bb1.11098 protein sequence in corn event MON 863 differs from the *B.t.* produced Cry3Bb1.11098 protein by 2 amino acids, or 7 amino acids relative to the wild type Cry3Bb1 protein. The deduced amino acid sequence of the Cry3Bb1.11098

protein in corn events MON 852, MON 858, MON 859 MON 862 and 96.040 is predicted to contain the same changes because the same *cry3Bb1.11098* gene was integrated into the transformation vectors used to create these corn events. Similarly, the deduced amino acid sequence of the Cry3Bb1.11231 protein in corn events MON 853, MON 854, MON 855 and MON 860 is predicted to contain the same changes because the same *cry3Bb1.11231* gene was integrated into the transformation vectors used to create these corn events. The deduced Cry3Bb1.11231 protein sequence in corn events producing the Cry3Bb1.11231 protein differs from the *B.t.* produced Cry3Bb1.11231 protein by 1 amino acid, or 5 amino acids relative to the wild type Cry3Bb1 protein. These amino acid changes are summarized in Table 3.

4.0 Characterization of the Cry3Bb1 Protein Produced in Corn Event MON 863

Additional analyses were also performed on the Cry3Bb1 protein produced in corn event MON 863. Cry3Bb1 protein was isolated from grain using immunoaffinity chromatography and analyzed by N-terminal sequencing. Peptide fragments produced by tryptic digestion were analyzed using matrix assisted laser desorption ionization time of flight (MALDI-TOF) mass spectrometry (Thoma *et al.*, 2001). Approximately 38% of the entire Cry3Bb1 protein sequence was identified by a combination of mass matching and N-terminal sequencing (Figure 2), confirming the amino acid sequence deduced from the DNA. Moreover, the mass spectral data were used to verify the presence of the alanine residue at position 2 and the Q349R substitution in corn event MON 863.

5.0 Conclusions

Previous reports did not indicate the presence of the intended Q349R substitution in the Cry3Bb1.11098 plant produced protein sequence. Recent additional analyses performed on both DNA and protein isolated from corn event MON 863 confirm that this substitution exists. The Q349R substitution is present in the predicted Cry3Bb1 protein sequence of all corn events producing the 11098 variant. This substitution was not intended for, and is therefore absent in corn events producing the Cry3Bb1.11231 protein variant.

Large quantities of Cry3Bb1.11098 and Cry3Bb1.11231 proteins were prepared from fermentation of *B.t.* (Hileman *et al.*, 2001) for mammalian and ecotoxicology evaluations as well as to serve as reference proteins in immunodiagnostic assays. These *B.t.* produced Cry3Bb1 protein variants were shown to be physicochemically and functionally equivalent to the respective corn produced proteins (Holleschak *et al.*, 2001a; Holleschak *et al.*, 2001b). Thus the Q349R substitution found in the plant produced Cry3Bb1.11098 protein is not expected to have changed the physicochemical and functional characteristics of this protein.

6.0 References

- Cavato, T. A., and Lirette, R. P. (2001). PCR analysis and DNA sequence of the insert in corn rootworm event MON 863. (Monsanto Company, St. Louis, MO) Study 01-01-39-02, Technical Report MSL-17108.
- Cavato, T. A., Mittanck, D. W., Albee, L. D., Rigden, E. C., Borevkova, I. G., Sanders, P. R., and Lirette, R. P. (1999). Molecular analysis of corn event MON 853. (Monsanto Company, St. Louis, MO) Study 98-01-39-17, Technical Report MSL-16031.
- Cavato, T. A., Rigden, E. C., Mittanck, D. W., and Lirette, R. P. (2001). Amended report for MSL-16505: Molecular analysis of corn event MON 863. (Monsanto Company, St. Louis, MO) Study 99-01-39-27, Technical Report MSL-17152.
- Donovan, W. P., Rupar, M. J., Slaney, A. C., Malvar, T., Gawron-Burke, M. C., and Johnson, T. B. (1992). Characterization of two genes encoding *Bacillus thuringiensis* insecticidal crystal proteins toxic to *Coleoptera* species. *Appl Environ Microbiol* 58:3921-3927.
- English, L. H., Brussock, S. M., Malvar, T. M., Bryson, J. W., Kulesza, C. A., Walters, F., Slatin, S. L., Von Tersch, M. A., and Romano, C. (2000). Nucleic acid segments encoding modified *Bacillus thuringiensis* coleopteran-toxic crystal proteins (United States Patent # 6,060,594: Ecogen, Inc., Langhorn, PA; Monsanto Company, St. Louis, MO).
- Hileman, R. E., Pyla, P. D., Lee, T. C., and Astwood, J. D. (2001). Amended report for MSL-15531: Characterization of *B.t.* protein 11098 and *B.t.* protein 11231 produced by fermentation. (Monsanto Company, St. Louis, MO) Study 98-01-39-05, Technical Report MSL-17219.
- Holleschak, G., Hileman, R. E., and Astwood, J. D. (2001a). Amended report for MSL-16596: Assessment of the physicochemical equivalence of Cry3Bb1.11098 and NPTII proteins in corn event MON 863 to microbial sources. (Monsanto Company, St. Louis, MO) Study 99-01-39-31, Technical Report MSL-17220.

Holleschak, G., Lee, T. C., Pyla, P. D., Hileman, R. E., and Astwood, J. D. (2001b). Amended report for MSL-15835: Assessment of the equivalence of *B.t.* protein 11098, *B.t.* protein 11231 and NPTII protein expressed in corn events MON 853 and MON 860 to microbial sources. (Monsanto Company, St. Louis, MO) Study 98-01-39-18; Technical Report MSL-17222.

Thoma, R. S., Holleschak, G., and Astwood, J. D. (2001). Primary structural protein characterization of Cry3Bb1.11098 protein from insect protected corn event MON 863 using MALDI time of flight mass spectrometric and N-terminal sequencing techniques. (Monsanto Company, St. Louis, MO) Technical Report MSL-17154.

Table 2. Specific nucleotide modifications and resulting amino acid changes used to produce the Cry3Bb1.11098 and Cry3Bb1.11231 proteins in *Bacillus thuringiensis*.

| Cry3Bb1 Allele | <i>cry3Bb1</i> DNA modifications ^a | Resulting Cry3Bb1 aa modifications ^a |
|-------------------------------|---|---|
| <i>B.t.</i> Strain EG11098 | A494G, T687C, A692G, C932T, D165G A938C, T942G, G949A, T954C | H231R, S311L, N313T, E317K |
| <i>B.t.</i> Strain EG11231 | T687C, A692G, C932T, A938C, T942G, G949A, T954C | H281R, S311L, N313T, E217K |

^a Standard biochemical nomenclature is shown using IUPAC-IUB single letter code. The first letter corresponds to the wild type base or amino acid, the following number corresponds to the position, and the second letter corresponds to the changed base or amino acid. The base and amino acid positions are shown relative to the wild type sequence.

Table 3. Summary of the amino acid differences among the Cry3Bb1 protein variants produced in corn.

| Cry3Bb1 Allele | Cry3Bb1 aa modifications ^a |
|-------------------------------------|--|
| Corn event MON 863 Cry3Bb1.11098 | A2, D166G, H232R, S312L, N314T, E318K, Q349R |
| Corn event MON 853 Cry3Bb1.11231 | A2, -, H232R, S312L, N314T, E318K, - |

^a Standard biochemical nomenclature is shown using IUPAC-IUB single letter code as described in Table 2. Amino acid positions are shifted upwards by one amino acid relative to the *B.t.* proteins (Table 2) due to the introduction of an alanine at position 2.




Figure 1. Summary of the origin of the Cry3Bb1 variant proteins present in corn events described in food, feed and environmental evaluations.

This placeholder page is in place of the following pages and can be found in **Cross Reference Number 1** of the **Confidential Attachment**.

| | |
|---------------------|---|
| Page | MSL-17137 Pages 17-22 of 22 |
| Page Title | Figure 2. Sequences of the Cry3Bb1 protein variants |
| Reason for Deletion | Discloses commercial information |
| FIFRA Reference | 10(d)(1)(A) |

MSL-17137 Pages 17-22 of 22 are found in the **Confidential Attachment**, Pages 3-8 of 8.

~~CONFIDENTIAL~~

ATTACHMENT